



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(51) International Patent Classification ⁶ : C12N 15/12, 15/57, 15/86, 7/01, 9/64, C07K 14/755, A61K 38/36, 38/37, 48/00		A2	(11) International Publication Number: WO 98/00542
			(43) International Publication Date: 8 January 1998 (08.01.98)
(21) International Application Number: PCT/US97/11785		(72) Inventors: JOLLY, Douglas, J.; 277 Hillcrest Drive, Leucadia, CA 92024 (US). BARBER, Jack, R.; 1168 Carlotta Street, San Diego, CA 92129 (US). CHANG, Stephen, M., W.; 12912 Camino del Valle, Poway, CA 92064 (US). RESPESS, James, G.; 4966 Lamont Street, San Diego, CA 92109 (US). ALLEN, John, R.; 8998 Adobe Buffs Drive, San Diego, CA 92129 (US). BODER, Mordechai; 3480 Juniper Street, San Diego, CA 92104 (US). CHONG, Kimberly; 12014 Dappie Way, San Diego, CA 92128 (US). DE LA VEGA, Dan, Jr.; 4220 Ampudia Street, San Diego, CA 92103 (US). DePOLO, Nicholas, J.; 964 Santa Estella, Solano Beach, CA 92075 (US). HSU, David, Chi-Tang; 7387 Mannix Court, San Diego, CA 92122 (US). IBANEZ, Carlos, E.; 13592 Millpond Way, San Diego, CA 92129 (US). MITTELSTAEDT, Denice, M.; 13687 Quinton Road, San Diego, CA 92129 (US). PRUSSAK, Charles, E.; 5121 Camino Playa Malaya, San Diego, CA 92124 (US). GREENGARD, Judith; 9053 Mesa Woods Avenue, San Diego, CA 92126 (US).	
(22) International Filing Date: 2 July 1997 (02.07.97)			
(30) Priority Data: 08/645,601 3 July 1996 (03.07.96) US 08/696,381 13 August 1996 (13.08.96) US Not furnished 4 June 1997 (04.06.97) US			
(71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US).			
		(74) Agents: KRUSE, Norman, J. et al.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US).	
		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
		Published Without international search report and to be republished upon receipt of that report.	

(54) Title: METHODS FOR ADMINISTRATION OF RECOMBINANT GENE DELIVERY VEHICLES FOR TREATMENT OF HEMOPHILIA


(57) Abstract

Methods are provided for obtaining measurable levels of a protein, nucleic acid molecule, or enzymatic product in a bodily fluid or cells of a human, comprising the step of administering to a human a recombinant retroviral preparation having a titer on HT1080 cells of greater than 10^5 cfu/ml, wherein the recombinant retroviral preparation is capable of directing the expression of a protein, nucleic acid molecule, or enzyme which generates an enzymatic product, such that measurable levels of the protein, nucleic acid molecule, or enzymatic product may be obtained in the bodily fluid or cells of the human.

RETROVIRAL BACKBONE (N2-derived)



CROSS-LESS RETROVIRAL BACKBONE: pBA-5



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS FOR ADMINISTRATION OF RECOMBINANT GENE DELIVERY VEHICLES FOR TREATMENT OF HEMOPHILIA

Field of the Invention

The present invention relates to methods of administration of recombinant gene transfer vehicles for the treatment of hemophilia, thrombosis, and other diseases. The present invention also relates generally to recombinant retroviruses, and more specifically, to high titer recombinant retroviral particle preparations suitable for a variety of applications.

Background of The Invention

Hemophilia is a genetic disease characterized by a severe blood clotting deficiency. As such, it will be amenable to treatment by gene therapy. In hemophilia A, a X-chromosome linked genetic defect disrupts the gene encoding factor VIII, a trace plasma glycoprotein which acts as a cofactor in conjunction with factor IXa in the activation of factor X. In humans, the factor VIII gene codes for 2,351 amino acids. The protein has six domains, designated from amino to carboxy terminus as A1, A2, B, A3, C1, and C2, respectively (Wood *et al.*, 1984, *Nature* 312:330; Vehar *et al.*, 1984, *Nature* 312:337; and Toole *et al.*, 1984, *Nature* 312:342) with a deduced molecular weight of about 280 kilo Daltons (kD). The 980 amino acid B domain is deleted in the activated procoagulant form of the protein. Additionally, in the native protein two polypeptide chains, the heavy and light chain, flanking the B domain, are bound to a divalent calcium cation.

The genetic defect causing hemophilia A affects about one in every 10,000 males. Due to the resultant clotting deficiency, those afflicted with the disease suffer severe bleeding episodes due to small injuries, internal bleeding, and joint hemorrhage, which leads to arthropathy, the major cause of morbidity in hemophilia. Normal levels of factor VIII average between 50 to 200 ng/ml of blood plasma (Mannucci, P.M. in *Practical Laboratory Hematology*, ed. Koepke, J.A., Churchill Livingstone, N.Y., pp:347-371, 1990); however, patients suffering from mild to moderate hemophilia A typically have plasma levels well below 2 - 60 ng/ml, while levels below about 2 ng/mL result in severe hemophilia.

Previously, therapy for hemophilia A involved repeated administration of human factor VIII purified from blood products pooled in lots from over 1000 donors. However, because of the low levels of circulating factor VIII, resulting pharmaceutical products using the natural protein typically were highly impure, with an estimated purity by weight (factor VIII to total protein) of approximately 0.04%. Due to the frequency of administration and inability to remove various human pathogens from such preparations, more than 90% of those suffering from hemophilia A were infected in the 1980s with the human immunodeficiency virus (HIV) as a result of their therapy. Many of these HIV infected patients and other HIV negative hemophiliacs have also been infected by Hepatitis B in the same way. Fortunately, recent advances in genetic engineering have lead to the commercial availability of a recombinant form of the protein free from contamination with human pathogens, except for those potentially derived from tissue culture origin of the proteins, or from human serum albumin used in formulation of the recombinant protein. However, this form of therapy is expensive and chronic, and a large proportion of hemophiliacs continue to rely on plasma-derived products due to expense and or shortages of the recombinant product. In addition, most hemophilia A patients in the United States do not presently receive factor VIII maintenance therapy, but instead only receive the polypeptide prior to activities or events which might cause bleeding, such as surgery, or to treat spontaneous bleeding. Interestingly, this is despite evidence showing that hemophilic arthropathy can be prevented by administering from an early age prophylactic amounts of factor VIII, typically 24 - 40 IU per kilogram bodyweight, three times a week. Such therapy kept factor VIII concentrations from falling below 1% of normal (Nillson, *et al.*, 1992, *J. Internal Med.* 232:23). For these reasons, a genetic therapy affording continuous, long-term therapeutically effective expression levels or amounts of factor VIII, *i.e.*, to decrease the severity of or eliminate the clotting disorder associated with hemophilia A, would be of great benefit.

A condition clinically indistinguishable from Hemophilia A is Hemophilia B, resulting from the deficiency of clotting factor IX. The incidence of this condition is about 5-fold lower than that of hemophilia A, and presents many of the same therapeutic

challenges and difficulties. For similar reasons, it would be of great benefit to provide a gene therapy to these patients.

Factor X deficiency results in a rare but serious bleeding disorder affecting 1 in 500,000 known as Stuart disease. Le *et al.*, 1997, *Blood* 89:1254-9, describes therapeutic levels of functional human factor X in rats after retroviral mediated hepatic gene therapy. As in the case of hemophilia A and B, a genetic therapy affording continuous, long term therapeutically effective expression levels or amounts of factor X, *i.e.*, to decrease the severity of or eliminate the clotting disorder associated with hemophilia B, would be of great benefit.

The instant invention also relates to the production and use of high titer recombinant retroviruses. Since the discovery of DNA in the 1940s and continuing through the most recent era of recombinant DNA technology, substantial research has been undertaken in order to realize the possibility that the course of disease may be affected through interaction with the nucleic acids of living organisms. Most recently, a wide variety of methods have been described for altering or affecting genes, including for example, viral vectors derived from retroviruses, adenoviruses, vaccinia viruses, herpes viruses, and adeno-associated viruses (see Jolly, 1994, *Cancer Gene Therapy* 1(1):51-64), as well as direct transfer techniques such as lipofection (Felgner *et al.*, 1989, *Proc. Natl. Acad. Sci. USA* 84:7413-7417), direct DNA injection (Acsadi *et al.*, 1991, *Nature* 352:815-818), microprojectile bombardment (Williams *et al.*, 1991, *PNAS* 88:2726-2730), liposomes of several types (*see, e.g.*, Wang *et al.*, 1987, *PNAS* 84:7851-7855) and administration of nucleic acids alone (PCT Patent Publication No. WO 90/11092).

Of these techniques, recombinant retroviral gene delivery methods have been most extensively utilized, in part due to: (1) the efficient entry of genetic material (the vector genome) into cells; (2) an active, efficient process of entry into the target cell nucleus; (3) relatively high levels of gene expression; (4) the potential to target particular cellular subtypes through control of the vector-target cell binding and the tissue-specific control of gene expression; (5) a general lack of pre-existing host immunity; and (6) substantial knowledge and clinical experience which has been gained with such vectors.

Briefly, retroviruses are diploid positive-strand RNA viruses that replicate through an integrated DNA intermediate. In particular, upon infection by the RNA virus, the retroviral genome is reverse-transcribed into DNA by a virally encoded reverse transcriptase that is carried as a protein in each retrovirus. The viral DNA is then integrated pseudo-
5 randomly into the host cell genome of the infecting cell, forming a "provirus" which is inherited by daughter cells.

Wild-type retroviral genomes (and their proviral copies) contain three genes (the *gag*, *pol* and *env* genes), which are preceded by a packaging signal (ψ), and two long terminal repeat (LTR) sequences which flank both ends. Briefly, the *gag* gene encodes the
10 internal structural (nucleocapsid) proteins. The *pol* gene codes for the RNA-dependent DNA polymerase which reverse transcribes the RNA genome, and the *env* gene encodes the retroviral envelope glycoproteins. The 5' and 3' LTRs contain *cis*-acting elements necessary to promote transcription and polyadenylation of retroviral RNA.

Adjacent to the 5' LTR are sequences necessary for reverse transcription of the
15 genome (the tRNA primer binding site) and for efficient encapsidation of retroviral RNA into particles (the ψ sequence). Removal of the packaging signal prevents encapsidation (packaging of retroviral RNA into infectious virions) of genomic RNA, although the resulting mutant can still direct synthesis of all proteins encoded in the viral genome.

Recombinant retroviruses and various uses thereof have been described in numerous
20 references including, for example, Mann *et al.*, 1983, *Cell* 33:153; Cane and Mulligan, 1984, *Proc. Natl. Acad. Sci. USA* 81:6349; Miller *et al.*, 1990, *Human Gene Therapy* 1:5-14; U.S. Patent Nos. 4,405,712; 4,861,719; 4,980,289 and PCT Patent Publication Nos. WO 89/02468; WO 89/05349 and WO 90/02806. Briefly, a foreign gene of interest may be incorporated into the retrovirus in place of the normal retroviral RNA. When the retrovirus
25 injects its RNA into a cell, the foreign gene is also introduced into the cell, and may then be integrated into the host's cellular DNA as if it were the retrovirus itself. Expression of this foreign gene within the host results in expression of the foreign protein by the host cell.

One disadvantage, however, of recombinant retroviruses is that they principally infect only replicating cells, thereby making efficient direct gene transfer difficult. Indeed,
30 some scientists have suggested that other, more efficient methods of gene transfer, such as

direct administration of pure plasmid DNA, be utilized (Davis *et al.*, 1993, *Human Gene Therapy* 4:733-740).

In order to increase the efficacy of recombinant retroviruses, methods which have been suggested for increasing the efficacy of recombinant retroviruses have principally been aimed at inducing the desired target cells to replicate, thereby allowing the retroviruses to infect the cells. Such methods have included, for example chemical treatment with 10% carbon tetrachloride in mineral oil (Kaleko *et al.*, 1991, *Human Gene Therapy* 2:27-32). Alternatively, others have suggested excising large portions of the liver (e.g., 70% in Rettinger *et al.*, 1994, *PNAS* 91:1460-1464; 70% in Moscioni *et al.*, 1993, *Surgery* 113:304-311) in order to stimulate the rapid division of hepatocytes and thereby increase the infection of such cells.

One further disadvantage of recombinant retroviruses, is that serum from primates (e.g., humans) is known to cause inactivation by an antibody independent complement lysis method. In particular retroviruses of avian, murine, feline, and simian origin are all inactivated and lysed by normal human serum. (Welsh *et al.*, 1975, *Nature* 257:612-614; Welsh *et al.*, 1976, *Virology* 74:432-440; Banapour *et al.*, 1986, *Virology* 152:268-271; and Cooper *et al.*, Immunology of the Complement System, Pub., American Press, Inc., pp. 139-162, 1986). The scientific literature has also reported that replication competent murine amphotropic retroviruses injected intravenously into primates are cleared within 15 minutes and that the disappearance is mediated, wholly or in part, by primate complement. (Cornetta *et al.*, 1991, *Human Gene Therapy* 2:5-14; Cornetta *et al.*, 1990, *Human Gene Therapy* 1:15-30; and Banapour *et al.*, 1986, *Virology* 152:268-271)

In order to increase the affect of recombinant retroviruses that are delivered *in vivo*, the present invention provides recombinant retrovirus compositions which are capable of surviving inactivation in human serum. In addition, the present invention provides high titer recombinant retrovirus compositions which allow delivery of therapeutics or palliatives by routes not previously deemed possible, and without the need to induce replication of cells by chemical treatment or by excision of a target organ such as the liver. The present invention provides these, as well as other related advantages.

Summary of the Invention

This invention provides for preparations of replication defective recombinant retrovirus expressing human factor VIII protein, wherein the recombinant retrovirus is capable of infecting human cells and is resistant to degradation by human complement. The invention also provides for preparations of replication defective recombinant retrovirus expressing human factor VIII protein in which the recombinant retrovirus preparation is resistant to degradation by human complement and is capable of inducing long term systemic expression of human factor VIII when administered intravenously to a human afflicted with hemophilia A. The wherein said long term systemic expression results in a measurable level of recombinant human factor VIII protein being produced in the blood of said human for a period of at least 30 days after the administration of said recombinant retroviral vector preparation and more preferably for at least six months after injection, and yet more preferably for longer periods of time as described herein.

Pharmaceutical compositions and therapeutic methods of the above-described retroviral vectors expressing factor VIII protein are also provided herein as are therapeutic methods for treatment of hemophilia A by intravenous injection of these retroviral vectors. The retroviral vectors of the invention can express a B domain-deleted form of factor VIII, which in one embodiment can be the SQN mutation of factor VIII. The retroviral vectors of the invention can have a titer on HT1080 cells of greater than 10^6 , more preferably 10^7 cfu/ml and more preferably at least 10^8 cfu/ml, more preferably 10^9 cfu/ml, more preferably at least 10^{10} cfu/ml, and most preferably 10^{11} cfu/ml.

In addition, the present invention provides high titer compositions comprising recombinant retroviruses, as well as methods for utilizing these compositions. Within one aspect of the present invention, methods are provided for obtaining measurable levels of a protein, nucleic acid molecule, or enzymatic product in a bodily fluid or cells of a human, comprising the step of administering to a human a recombinant retroviral preparation having a titer on HT1080 cells of greater than 10^5 cfu/ml, wherein the recombinant retroviral preparation is capable of directing the expression of a protein, nucleic acid molecule, or enzyme which generates an enzymatic product, such that measurable levels of the protein, nucleic acid molecule, or enzymatic product may be obtained in the bodily fluid or cells of

said human. Within certain embodiments, the titer may be greater than 10^6 cfu/ml, 10^7 cfu/ml, 10^8 cfu/ml, 10^9 cfu/ml, 10^{10} cfu/ml, or 10^{11} cfu/ml.

Within other aspects of the invention, methods are provided for obtaining measurable levels of a protein, nucleic acid molecule, or enzymatic product in a bodily fluid or cells of a human, comprising the steps of administering to a human a recombinant retroviral preparation having a titer in human serum and on HT1080 cells equivalent to its' titer in heat-inactivated serum and on HT1080 cells, wherein the recombinant retroviral preparation is capable of directing the expression of a protein, nucleic acid molecule, or enzyme which generates an enzymatic product, such that measurable levels of the protein, nucleic acid molecule, or enzymatic product may be obtained in the bodily fluid or cells of said human.

As utilized within the context of the present invention, "measurable levels" of a protein, nucleic acid molecule, or enzymatic product refers to a statistically significant level of detection over background, utilizing any suitable technique (e.g., antibody-mediated detection of a protein, PCR analysis for the presence of a nucleic acid molecule, or visualization of enzymatic products). Further, as utilized within the context of the present invention, "equivalent" titers are deemed to be those which are substantially the same within a given assay, generally, within about three-fold of each other.

Within certain embodiments of the invention, the recombinant retrovirus is administered to a site such as the cerebral spinal fluid, bone marrow, joints, arterial endothelial cells, rectum, buccal/sublingual, vagina, the lymph system, to an organ selected from the group consisting of lung, liver, spleen, skin, blood and brain, or to a site selected from the group consisting of tumors and interstitial spaces. Within other embodiments, the recombinant retrovirus may be administered intraocularly, intranasally, sublingually, orally, topically, intravesically, intrathecally, topically, intravenously, intraperitoneally, intracranially, intramuscularly, or subcutaneously.

These and other aspects of the present invention will become evident upon reference to the following detailed description and attached drawings. In addition, various references are set forth below which describe in more detail certain procedures or compositions (e.g.,

plasmids, etc.), and are therefore incorporated by reference in their entirety as if each were individually noted for incorporation.

Brief Description of the Drawings

- Figure 1 is a schematic illustration of p31N2R5(+).
- 5 Figure 2 is a schematic illustration of pN2R3(-).
- Figure 3 is a schematic illustration of p31N25 (+).
- Figure 4 is a schematic illustration of pN2R3(+).
- Figure 5 is a schematic illustration of pN2R5(-).
- Figure 6 is a schematic illustration of p31N25 (+).
- 10 Figure 7 is a schematic illustration of pTK A.
- Figure 8 is a schematic illustration of pPrTK A.
- Figure 9 is a schematic illustration of pTK-1 and pTK-3.
- Figure 10 is a bar graph which illustrates the effect of Ganciclovir on CT26, CT26 bgal and CT26TK Neo cells.
- 15 Figure 11 is a graph which illustrates the effect of tumor volume over time in a Ganciclovir dose study of mice injected with CT26TK Neo.
- Figure 12 is a graph illustrating the effect of Ganciclovir in CT26 versus CT26TK Neo cells.
- Figure 13 is a graph demonstrating retention of viral activity upon reconstitution of a representative recombinant retrovirus lyophilized in a formulation buffer containing mannitol.
- 20 Figure 14 is a graph demonstrating retention of viral activity upon reconstitution of a representative recombinant retrovirus lyophilized in a formulation buffer containing lactose.
- Figure 15 is a graph demonstrating retention of viral activity upon reconstitution of a representative recombinant retrovirus lyophilized in a formulation buffer containing trehalose.
- 25 Figures 16A-16D are representative graphs comparing stability of liquid non-lyophilized recombinant retrovirus stored at -80°C versus lyophilized formulated recombinant retrovirus stored at -20°C, using various saccharides.

Figure 17 is a bar graph which depicts the results of a reverse transcriptase assay on samples sliced from a gel. Slice 1 is from the lowest part of the gel, and slice 8 from the highest.

Figure 18 depicts a 8% to 25% gradient polyacrylamide gel. Lane 1 is DA/ β gal S-5 500 purified; Lane 2 is DA/ β gal crude supernatant; Lane 3 is HIV-IT crude supernatant; lane 4 is DAC 6A3 crude supernatant; Lane 5 is HBc/SA2 crude supernatant; and Lane 5 is molecular weight markers.

Figure 19 is a schematic illustration of pDHF811.

Figures 20A-D are graphs which indicate the total quantity of lactate in low seed and10 high seed cultures (Figures 20A and 20B, respectively) and the level of lactate production per day in low seed and high seed cultures (Figures 22C and 22D, respectively).

Figure 21 is a bar graph which depicts the titer of the cell line 2X- β -gal under different initial seeding conditions.

Figure 22: Factor VIII antigen was determined in citrated blood samples drawn on15 the indicated days. Antigen determined by a standard curve obtained using normal pooled human plasma is plotted on the left y axis; the per cent of normal human levels is indicated on the right y axis.

Figure 23: Factor VIII expression in rabbits. Juvenile rabbits were injected with a total of 10^9 cfu of the factor VIII vector B-del-1 in a total of 9 injections over days 0,1, and20 2. Citrated plasma samples were obtained at the indicated times and human factor VIII antigen levels determined by ELISA. Rabbits 93 and 95 (controls) received formulation buffer in lieu of vector.

Figure 24: Serum samples were obtained from rabbits on indicated days, and human growth hormone antigen was measured using ELISA .

25 Figure 25: Serum samples were obtained from mice on indicated days and human growth hormone antigen was determined by ELISA. Since the animals were too small on Day 14 to yield sufficient sample for the ELISA measurement, sera from test animals 1-5, test animals 6-10, and control animals 11 and 12 were respectively pooled prior to measurement. Data from subsequent days were obtained from individual animals.

Figure 26: Sera were obtained from adult mice on the indicated days. Human growth hormone antigen was measured by ELISA. Each point represents an individual animal. Negative controls had been injected with formulation buffer; positive controls with 1E7 cfu retroviral vector; "600 ug CsA" with both vector and cyclosporin A as described in the text.

Figure 27: PCR localization of vector in B-del-1 treated rabbits. Rabbits #82 and 95 shown in figure 22, herein were sacrificed on Day 511. DNA was extracted from indicated organs, subjected to 40 cycles of PCR, and dot blotted, and hybridized to 32-P labeled probe as described in text. Several replicate PCR reactions were run. Lanes indicated "spike" had 5 copies of amplicon added to the template.

Figure 28: Quantitative PCR of beta-galactosidase vector in organs of transduced mice. Juvenile mice, transduced as described in the text, were sacrificed on Day 59. DNA was obtained from indicated organs and subjected to quantitative PCR as described.

Figure 29: Human growth hormone antigen in organs of transduced mice. Indicated organs were lysed and analyzed by ELISA as described in the text. Results are shown normalized to the amount of total protein recovered from the organ.

Figure 30: Expression of human factor VIII antigen in transduced normal dogs. Juvenile dogs (#47 and #55) were injected with a total of 4E9 cfu of factor VIII vector B-del-1 in a total of 9 injections over days 0,1, and 2. Control dog #39 received formulation buffer on the same schedule. Dog #47 was boosted with the same amount of vector on days 90, 91, and 92. At indicated times, cuitrated plasma was obtained and analyzed by ELISA for human factor VIII antigen. Percent human level is calculated based on a level of 200 ng/ml factor VIII in pooled in normal human plasma.

Figure 31: Transfer of expression of human growth hormone following adoptive transfer of splenocytes from transduced donor mice to lethally irradiated syngeneic recipients. Juvenile mice were transduced with growth hormone retroviral vector as described in the text. On day 59, Donors #1-5 were sacrificed, and splenocytes from each were injected into two lethally irradiated recipients (A and B, numbered according to donor number). Every 14 days subsequent to reconstitution, serum from the recipients was monitored for growth hormone levels.

Figure 32: Whole blood clotting times in hemophiliac dogs treated with factor VIII vector. Two hemophiliac dogs (#28 and #33) were injected with B-del-1 as described in the text. At indicated times, citrated whole blood samples were obtained and clotting times were determined following recalcification. Clotting times of two untreated normal littermates are included for reference.

Figure 33 is a schematic illustration of pKT1.

Figure 34 is a description of all modifications carried out on retroviral vector as shown in A), resulting in the cross-less retroviral vector shown in B). The cross-less retroviral backbone cloned into a prokaryotic vector is called pBA-5.

Figure 35: Schematic representation of pKm201-80L.

Figure 36: Schematic representation of pKm201-90-H.

Figure 37: Schematic representation of pSVF8-t β 2.

Figure 38: Nucleotide and predicted amino acid sequence linking the light and heavy chains of Factor VIII in pSVF8-t β 2.

Figure 39: Linker sequences for pSVF8-500 and pc1/1 AG α F8-500.

Figure 40: Schematic representation of pSVF8500 and pc1/1 AG α F8-500.

Figure 41: Linker sequence for pSVF8-500B.

Detailed Description of the Invention

Prior to setting forth the invention, it may be helpful to an understanding thereof to set forth definitions of certain terms that will be used hereinafter.

"Vector construct", "retroviral vector", "recombinant vector", and "recombinant retroviral vector" refers to a nucleic acid construct which carries, and within certain embodiments, is capable of directing the expression of a nucleic acid molecule of interest. The retroviral vector must include at least one transcriptional promoter/enhancer or locus defining element(s), or other elements which control gene expression by other means such as alternate splicing, nuclear RNA export, post-translational modification of messenger, or post-transcriptional modification of protein. Such vector constructs must also include a packaging signal, long terminal repeats (LTRs) or portion thereof, and positive and negative strand primer binding sites appropriate to the retrovirus used (if these are not already present

in the retroviral vector). Optionally, the recombinant retroviral vector may also include a signal which directs polyadenylation, selectable markers such as Neomycin resistance, TK, hygromycin resistance, phleomycin resistance, histidinol resistance, or DHFR, as well as one or more restriction sites and a translation termination sequence. By way of example, such
5 vectors typically include a 5' LTR, a tRNA binding site, a packaging signal, an origin of second strand DNA synthesis, and a 3' LTR or a portion thereof.

"Recombinant retrovirus", "retroviral gene delivery vehicle" and "retroviral vector particle" as utilized within the present invention refers to a retrovirus which carries at least one gene of interest. The retrovirus may also contain a selectable marker. The recombinant
10 retrovirus is capable of reverse transcribing its genetic material into DNA and incorporating this genetic material into a host cell's DNA upon infection.

"Factor VIII" is a nonenzymatic cofactor found in blood in an inactive precursor form. Precursor factor VIII is converted to the active cofactor, factor VIIIa, through limited proteolysis at specific sites by plasma proteases, notably thrombin and factor IXa. The
15 majority of factor VIII circulates as a two-chain heterodimer most likely due to intracellular or pericellular processing of the single chain gene product. The two chains are noncovalently associated in a metal ion dependent manner.

The "biological activity" of factor VIII refers to a function or set of functions performed by the polypeptide or fragments thereof in a biological system or in an *in vitro*
20 facsimile thereof. The biological activity of factor VIIIa is characterized by its ability to form a membrane binding site for factors IXa and X in a conformation suitable for activation of the latter by the former. This includes standard assays of factor IX or X activation, binding to phospholipids, von Willebrand factor, or specific cell surface molecules, and susceptibility to thrombin, factor IXa, activated protein C, or other specific
25 proteases under appropriate conditions. More particularly biologically active factor VIII corrects the coagulation defect in plasma derived from individuals afflicted with hemophilia A.

A "factor VIII cDNA molecule" is one encoding a full-length factor VIII polypeptide, a B domain deleted factor VIII protein, or other forms of factor VIII protein

with biological activity. The human full-length factor VIII coding region is 7,056 nucleotides.

"Biologically active factor IX" encompasses those forms of factor IX which are capable of correcting the coagulation defect in plasma derived from individuals afflicted with hemophilia B as measured in a standard *in vitro* clotting assay.

As noted above, the present invention provides for methods of administration of recombinant gene delivery vehicles for treatment of hemophilia and a variety of other disorders by gene therapy techniques. The present invention provides high titer complement resistant recombinant retroviral preparations which are suitable for administration to humans. Such preparations provide the unexpected result of providing efficacious gene therapy for a variety of diseases (and by a variety of routes) that were previously not amenable for gene therapy. In addition, the present invention provides recombinant retroviruses which are capable of surviving inactivation in human serum, thereby allowing more efficient gene transfer over prolonged periods of time.

A. PREPARATION OF RECOMBINANT GENE DELIVERY VEHICLES:
GENE DELIVERY VEHICLES

The therapeutic proteins of the present invention may be utilized in a wide variety of gene delivery vehicles. The gene delivery vehicle may be of either viral or non-viral origin (See generally, Jolly, 1994, *Cancer Gene Therapy* 1:51-64; Kimura, 1994, *Human Gene Therapy* 5:845-852; Connelly, 1995, *Human Gene Therapy* 6:185-193; and Kaplitt, 1994, *Nature Genetics* 6:148-153). Gene therapy vehicles for delivery of constructs including a coding sequence of a therapeutic of the invention can be administered either locally or systemically. These constructs can utilize viral or non-viral vector approaches in *in vivo* or *ex vivo* modality. Expression of such coding sequence can be induced using endogenous mammalian or heterologous promoters. Expression of the coding sequence *in vivo* can be either constitutive or regulated as is described in detail below.

1. Retroviral vectors

As noted above, the present invention provides recombinant retroviruses which are constructed to carry or express a selected nucleic acid molecule of interest. Briefly, numerous retroviral gene delivery vehicles may be utilized within the context of the present invention, including for example those described in EP 0,415,731; WO 90/07936; WO 94/03622; WO 93/25698; WO 93/25234; U.S. Patent No. 5,219,740; WO 93/11230; 5 WO 93/10218; Vile and Hart, 1993, *Cancer Res.* 53:3860-3864; Vile and Hart, 1993, *Cancer Res.* 53:962-967; Ram *et al.*, 1993, *Cancer Res.* 53:83-88; Takamiya *et al.*, 1992, *J. Neurosci. Res.* 33:493-503; Baba *et al.*, 1993, *J. Neurosurg.* 79:729-735 (U.S. Patent No. 4,777,127, GB 2,200,651, EP 0,345,242 and WO 91/02805). Particularly preferred 10 recombinant retroviruses include those described in WO 91/02805.

Retroviral gene delivery vehicles of the present invention may be readily constructed from a wide variety of retroviruses, including for example, B, C, and D type retroviruses as well as spumaviruses and lentiviruses (see RNA Tumor Viruses, Second Edition, Cold Spring Harbor Laboratory, 1985). Preferred retroviruses for the preparation or construction 15 of retroviral gene delivery vehicles of the present invention include retroviruses selected from the group consisting of Avian Leukosis Virus, Bovine Leukemia Virus, Murine Leukemia Virus, Mink-Cell Focus-Inducing Virus, Murine Sarcoma Virus, Reticuloendotheliosis virus and Rous Sarcoma Virus. Particularly preferred Murine Leukemia Viruses include 4070A and 1504A (Hartley and Rowe, 1976, *J. Virol.* 19:19-25), 20 Abelson (ATCC No. VR-999), Friend ATCC No. VR-245), Graffi, Gross (ATCC No. VR-590), Kirsten, Harvey Sarcoma Virus and Rauscher (ATCC No. VR-998), and Moloney Murine Leukemia Virus (ATCC No. VR-190). Such retroviruses may be readily obtained from depositories or collections such as the American Type Culture Collection ("ATCC"; Rockville, Maryland), or isolated from known sources using commonly available 25 techniques.

Any of the above retroviruses may be readily utilized in order to assemble or construct retroviral gene delivery vehicles given the disclosure provided herein, and standard recombinant techniques (*e.g.*, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press, 1989; Kunkle, 1985, *PNAS* 82:488). 30 In addition, within certain embodiments of the invention, portions of the retroviral gene

delivery vehicles may be derived from different retroviruses. For example, within one embodiment of the invention, retrovector LTRs may be derived from a Murine Sarcoma Virus, a tRNA binding site from a Rous Sarcoma Virus, a packaging signal from a Murine Leukemia Virus, and an origin of second strand synthesis from an Avian Leukosis Virus.

5 Within preferred aspects of the present invention, recombinant retroviruses may be made by introducing a vector construct as discussed above, into a cell (termed a "packaging cell") which contains those elements necessary for production of infectious recombinant retrovirus which are lacking in the vector construct. A wide variety of retrovector constructs may be utilized within the present invention in order to prepare recombinant
10 retroviruses. For example, within one aspect of the present invention retrovector constructs are provided comprising a 5' LTR, a tRNA binding site, a packaging signal, one or more heterologous sequences, an origin of second strand DNA synthesis and a 3' LTR, wherein the vector construct lacks gag/pol or env coding sequences. Briefly, Long Terminal Repeats ("LTRs") are subdivided into three elements, designated U5, R and U3. These elements
15 contain a variety of signals which are responsible for the biological activity of a retrovirus, including for example, promoter and enhancer elements which are located within U3. LTRs may be readily identified in the provirus due to their precise duplication at either end of the genome. As utilized herein, a 5' LTR should be understood to include a 5' promoter element and sufficient LTR sequence to allow reverse transcription and integration of the
20 DNA form of the vector. The 3' LTR should be understood to include a polyadenylation signal, and sufficient LTR sequence to allow reverse transcription and integration of the DNA form of the vector.

 The tRNA binding site and origin of second strand DNA synthesis are also important for a retrovirus to be biologically active, and may be readily identified by one of
25 skill in the art. For example, retroviral tRNA binds to a tRNA binding site by Watson-Crick base pairing, and is carried with the retrovirus genome into a viral particle. The tRNA is then utilized as a primer for DNA synthesis by reverse transcriptase. The tRNA binding site may be readily identified based upon its location just downstream from the 5' LTR. Similarly, the origin of second strand DNA synthesis is, as its name implies, important for

the second strand DNA synthesis of a retrovirus. This region, which is also referred to as the poly-purine tract, is located just upstream of the 3' LTR.

In addition to a 5' and 3' LTR, tRNA binding site, and origin of second strand DNA synthesis, certain preferred retrovector constructs which are provided herein also comprise a packaging signal, as well as one or more nucleic acid molecules (e.g., heterologous sequences), each of which is discussed in more detail below.

Within one aspect of the invention, retrovector constructs are provided which lack both *gag/pol* and *env* coding sequences. As utilized herein, the phrase "lacks *gag/pol* or *env* coding sequences" should be understood to mean that the retrovector does not contain at least 20, preferably at least 15, more preferably at least 10, and most preferably at least 8 consecutive nucleotides which are found in *gag/pol* or *env* genes, and in particular, within *gag/pol* or *env* expression cassettes that are used to construct packaging cell lines for the retrovector construct.

Packaging cell lines suitable for use with the above-described retrovector constructs may be readily prepared (see PCT Patent Publication No. WO 95/30763; see also WO 92/05266), and utilized to create producer cell lines (also termed vector cell lines or "VCLs") for the production of recombinant vector particles. Within particularly preferred embodiments of the present invention packaging cell lines are made from human (e.g., HT1080 cells) or mink parent cell lines, thereby allowing production of recombinant retroviruses that are capable of surviving inactivation in human serum.

Utilizing the methods of the present invention as disclosed herein, packaging cell lines that produce greater than recombinant retroviral particles at titers greater than 10^6 or 10^7 cfu/ml (in crude supernatant) may readily be obtained. In addition, it should be noted that such titers are generally obtained from titer assays on HT1080 cells, which produce a three-fold lower titer than titers obtained on murine 3T3 cells.

2. Alphavirus and eucaryotic layered vector gene delivery vehicles

The present invention also provides a variety of alphavirus-based vectors which can function as gene delivery vehicles. Such vectors can be constructed from a wide variety of alphaviruses, including for example, Sindbis viruses vectors, Semliki Forest virus (ATCC

VR-67; ATCC VR-1247), Ross River virus (ATCC VR-373; ATCC VR-1246) and Venezuelan equine encephalitis virus (ATCC VR923; ATCC VR-1250; ATCC VR-1249; ATCC VR-532).

As an example, the Sindbis virus, which is the prototype member of the alphavirus genus of the togavirus family is an unsegmented genomic RNA (49S RNA) of virus of approximately 11,703 nucleotides in length. This virus contains a 5' cap and a 3' polyadenylated tail, and displays positive polarity. Infectious enveloped Sindbis virus is produced by assembly of the viral nucleocapsid proteins onto the viral genomic RNA in the cytoplasm and budding through the cell membrane embedded with viral encoded glycoproteins. Entry of virus into cells is by endocytosis through clathrin coated pits, fusion of the viral membrane with the endosome, release of the nucleocapsid, and uncoating of the viral genome. During viral replication the genomic 49S RNA serves as template for synthesis of the complementary negative strand. This negative strand in turn serves as template for genomic RNA and an internally initiated 26S subgenomic RNA. The Sindbis viral nonstructural proteins are translated from the genomic RNA while structural proteins are translated from the subgenomic 26S RNA. All viral genes are expressed as a polyprotein and processed into individual proteins by post translational proteolytic cleavage. The packaging sequence resides within the nonstructural coding region, therefore only the genomic 49S RNA is packaged into virions.

A variety of different alphavirus vector systems may be constructed and utilized within the present invention. Representative examples of such systems include those described in U.S. Patent application Nos. 08/198,450, 08/405,627 and 08/679,640, U.S. Patent Nos. 5,091,309; 5,217,879 and 5,185,440, PCT Publication Nos. WO 92/10578, WO/94/21792, WO 95/27069, WO 95/27044 and WO 95/07994.

Particularly preferred alphavirus vectors for use within the present invention include those which are described within U.S. Application No. 08/198,450. Briefly, within one embodiment, alphavirus vector constructs are provided comprising a 5' sequence which is capable of initiating *in vitro* transcription of a alphavirus RNA, a nucleotide sequence encoding alphavirus non-structural proteins, a viral junction region which is active, modified to reduce viral transcription of the subgenomic fragment, or inactivated such that

viral transcription of the subgenomic fragment is prevented, and an alphavirus RNA polymerase recognition sequence.

In still further embodiments, the vector constructs described above contain no alphavirus structural proteins in the vector constructs. The selected heterologous sequence may be located downstream from the viral junction region; in the vector constructs having a second viral junction, the selected heterologous sequence may be located downstream from the second viral junction region, where the heterologous sequence is located downstream, the vector construct may comprise a polylinker located between the viral junction region and said heterologous sequence, and preferably the polylinker does not contain a wild-type Sindbis virus restriction endonuclease recognition sequence.

In addition, within one embodiment of the invention the gene delivery vehicles is a eukaryotic layered expression systems (ELVS) (see PCT Patent Publication Nos. WO 95/07994 and WO 96/17072 for a detailed description of eukaryotic layered expression systems). Although initially developed for alphavirus vectors, ELVS vectors may also be developed using other viral and non-viral nucleotide sequences (see PCT Patent Publication Nos. WO 95/07994 and WO 96/17072).

3. Adeno-associated viral vectors

Gene delivery vehicles of the present invention also include parvovirus such as adenovirus associated virus (AAV) vectors. Representative examples of such vectors include the AAV vectors disclosed by Srivastava in WO 93/09239, Samulski *et al.*, 1989, *J. Vir.* 63:3822-3828; Mendelson *et al.*, 1988, *Virol.* 166:154-165; Flotte *et al.*, 1993, *PNAS* 90(22):10613-10617. Particularly preferred AAV vectors comprise the two AAV inverted terminal repeats in which the native D-sequences are modified by substitution of nucleotides, such that at least 5 native nucleotides and up to 18 native nucleotides, preferably at least 10 native nucleotides up to 18 native nucleotides, most preferably 10 native nucleotides are retained and the remaining nucleotides of the D-sequence are deleted or replaced with non-native nucleotides. The native D-sequences of the AAV inverted terminal repeats are sequences of 20 consecutive nucleotides in each AAV inverted terminal repeat (*i.e.*, there is one sequence at each end) which are not involved in HP formation. The

non-native replacement nucleotide may be any nucleotide other than the nucleotide found in the native D-sequence in the same position. Other employable exemplary AAV vectors are pWP-19, pWN-1, both of which are disclosed in Nahreini, 1993, *Gene* 124:257-262.

Another example of such an AAV vector is psub201. See Samulski, 1987, *J. Virol.*

- 5 61:3096. Another exemplary AAV vector is the Double-D ITR vector. How to make the Double D ITR vector is disclosed in U.S. Patent No. 5,478,745. Still other vectors are those disclosed in Carter, U.S. Patent No. 4,797,368 and Muzyczka, U.S. Patent No. 5,139,941; Chartejee, U.S. Patent No. 5,474,935; and Kotin, PCT Patent Publication WO 94/288157. Yet a further example of an AAV vector employable in this invention is SSV9AFABTKneo, 10 which contains the AFP enhancer and albumin promoter and directs expression predominantly in the liver. Its structure and how to make it are disclosed in Su, *Human Gene Therapy* 7:463-470, 1996. Additional AAV gene therapy vectors are described in U.S. Patent Nos. 5,354,678; 5,173,414; 5,139,941; and 5,252,479.

4. Other Viral Vector Delivery Systems

- 15 In addition to retroviral vectors and alphavirus-based vectors, numerous other viral vectors systems may also be utilized as a gene delivery vehicle. For example, within one embodiment of the invention adenoviral vectors may be employed as a gene delivery vehicle. Representative examples of such vectors include those described by, for example, Berkner, 1988, *Biotechniques* 6:616-627; Rosenfeld *et al.*, 1991, *Science* 252:431-434; WO 20 93/9191; Kolls *et al.*, 1994, *PNAS* 91(1):215-219; Kass-Eisler *et al.*, 1993, *PNAS* 90(24):11498-502; Guzman *et al.*, 1993, *Circulation* 88(6):2838-48; Guzman *et al.*, 1993, *Cir. Res.* 73(6):1202-1207; Zabner *et al.*, 1993, *Cell* 75(2):207-216; Li *et al.*, 1993, *Hum. Gene Ther.* 4(4):403-409; Caillaud *et al.*, 1993, *Eur. J. Neurosci.* 5(10):1287-1291; Vincent *et al.*, 1993, *Nat. Genet.* 5(2):130-134; Jaffe *et al.*, 1992, *Nat. Genet.* 1(5):372-378; and 25 Levrero *et al.*, 1991, *Gene* 101(2):195-202; and WO 93/07283; WO 93/06223; and WO 93/07282. Exemplary known adenoviral gene therapy vectors employable in this invention include those described in the above referenced documents and in PCT Patent Publication No. WO 94/12649, WO 93/03769, WO 93/19191, WO 94/28938, WO 95/11984, WO 95/00655, WO 95/27071, WO 95/29993, WO 95/34671, WO 96/05320, WO 94/08026, WO

94/11506, WO 93/06223, WO 94/24299, WO 95/14102, WO 95/24297, WO 95/02697, WO 94/28152, WO 94/24299, WO 95/09241, WO 95/25807, WO 95/05835, WO 94/18922 and WO 95/09654. Alternatively, administration of DNA linked to killed adenovirus as described in Curiel, 1992, *Hum. Gene Ther.* 3:147-154, may be employed.

5 Gene delivery vehicles of the present invention also include herpes vectors. Representative examples of such vectors include those disclosed by Kit in *Adv. Exp. Med. Biol.* 215:219-236, 1989; and those disclosed in U.S. Patent No. 5,288,641 and EP 0176170 (Roizman). Additional exemplary herpes simplex virus vectors include HFEM/ICP6-LacZ disclosed in WO 95/04139 (Wistar Institute), pHSVlac described in Geller, 1988, *Science* 10 241:1667-1669, and in WO 90/09441 and WO 92/07945; HSV Us3::pgC-lacZ described in Fink, 1992, *Human Gene Therapy* 3:11-19; and HSV 7134, 2 RH 105 and GAL4 described in EP 0453242 (Breakefield), and those deposited with the ATCC as accession numbers ATCC VR-977 and ATCC VR-260.

Gene delivery vehicles may also be generated from a wide variety of other viruses, 15 including for example, poliovirus (Evans *et al.*, 1989, *Nature* 339:385-388; and Sabin, 1973, *J. Biol. Standardization* 1:115-118); rhinovirus; pox viruses, such as canary pox virus or vaccinia virus (Fisher-Hoch *et al.*, 1989, *PNAS* 86:317-321; Flexner *et al.*, 1989, *Ann. N.Y. Acad. Sci.* 569:86-103; Flexner *et al.*, 1990, *Vaccine* 8:17-21; U.S. Patent Nos. 4,603,112, 4,769,330 and 5,017,487; WO 89/01973); SV40 (Mulligan *et al.*, 1979, *Nature* 20 277:108-114); influenza virus (Luytjes *et al.*, 1989, *Cell* 59:1107-1113; McMichael *et al.*, 1983, *N. Eng. J. Med.* 309:13-17; and Yap *et al.*, 1978, *Nature* 273:238-239); SV40; HIV (Poznansky, 1991, *J. Virol.* 65:532-536); measles (EP 0 440,219); astrovirus (Munroe *et al.*, 1993, *J. Vir.* 67:3611-3614); and coronavirus, as well as other viral systems (*e.g.*, EP 0,440,219; WO 92/06693; U.S. Patent No. 5,166,057). In addition, viral carriers may be 25 homologous, non-pathogenic(defective), replication competent virus (*e.g.*, Overbaugh *et al.*, 1988, *Science* 239:906-910), and nevertheless induce cellular immune responses, including CTL.

4. Non-viral gene delivery vehicles

Other gene delivery vehicles and methods that may be employed such as, for example, nucleic acid expression vectors; polycationic condensed DNA linked or unlinked to killed adenovirus alone, for example see U.S. Serial No. 08/366,787, filed December 30, 1994, and Curiel, 1992, *Hum Gene Ther* 3:147-154; ligand linked DNA, for example see Wu, 5 1989, *J Biol Chem* 264:16985-16987; eucaryotic cell delivery vehicles cells, for example see U.S. Serial No. 08/240,030, filed May 9, 1994, and U.S. Serial No. 08/404,796; deposition of photopolymerized hydrogel materials; hand-held gene transfer particle gun, as described in US Patent No. 5,149,655; ionizing radiation as described in U.S. 5,206,152 and in WO 92/11033; nucleic charge neutralization or fusion with cell membranes. Additional 10 approaches are described in Philip, 1994, *Mol Cell Biol* 14:2411-2418, and in Woffendin, 1994, *Proc. Natl. Acad. Sci.* 91:1581-1585.

Particle mediated gene transfer may be employed, for example, see U.S. provisional application No. 60/023,867. Briefly, the sequence of interest can be inserted into conventional vectors that contain conventional control sequences for high level expression, 15 and then be incubated with synthetic gene transfer molecules such as polymeric DNA-binding cations like polylysine, protamine, and albumin, linked to cell targeting ligands such as asialoorosomucoid, as described in Wu and Wu, 1987, *J. Biol. Chem.* 262:4429-4432, insulin as described in Hucked, 1990, *Biochem Pharmacol* 40:253-263, galactose as described in Plank, 1992, *Bioconjugate Chem* 3:533-539, lactose or transferrin.

20 Naked DNA may also be employed. Exemplary naked DNA introduction methods are described in WO 90/11092 and U.S. Patent No. 5,580,859. Uptake efficiency may be improved using biodegradable latex beads. DNA coated latex beads are efficiently transported into cells after endocytosis initiation by the beads. The method may be improved further by treatment of the beads to increase hydrophobicity and thereby facilitate 25 disruption of the endosome and release of the DNA into the cytoplasm.

Liposomes that can act as gene delivery vehicles are described in U.S. Patent No. 5,422,120, PCT Patent Publication Nos. WO 95/13796, WO 94/23697, and WO 91/144445, and European Patent Publication No. 524,968. As described in U.S. provisional application No. 60/023,867, nucleic acid sequences can be inserted into conventional vectors that 30 contain conventional control sequences for high level expression, and then be incubated

with synthetic gene transfer molecules such as polymeric DNA-binding cations like polylysine, protamine, and albumin, linked to cell targeting ligands such as asialoorosomucoid, insulin, galactose, lactose, or transferrin. Other delivery systems include the use of liposomes to encapsulate DNA comprising the gene under the control of a variety of tissue-specific or ubiquitously-active promoters. Further non-viral delivery suitable for use includes mechanical delivery systems such as the approach described in Woffendin *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91(24):11581-11585. Moreover, the coding sequence and the product of expression of such can be delivered through deposition of photopolymerized hydrogel materials. Other conventional methods for gene delivery that can be used for delivery of the coding sequence include, for example, use of hand-held gene transfer particle gun, as described in U.S. Patent No. 5,149,655; use of ionizing radiation for activating transferred gene, as described in U.S. Patent No. 5,206,152 and PCT Patent Publication No. WO 92/11033.

Exemplary liposome and polycationic gene delivery vehicles are those described in U.S. Patent Nos. 5,422,120 and 4,762,915, in PCT Patent Publication Nos. WO 95/13796, WO 94/23697, and WO 91/14445, in European Patent Publication No. 524,968 and in Starrier, Biochemistry, pages 236-240 (1975) W.H. Freeman, San Francisco; Shokai, *Biochem Biophys Acct* 600:1, 1980; Bayer, 1979, *Biochem Biophys Acct* 550:464; Rivet, 1987, *Meth Enzyme* 149:119; Wang, 1987, *Proc Natl Acad Sci* 84:7851; Plant, 1989, *Anal Biochem* 176:420.

B. Construction of Recombinant Gene Delivery Vehicles Expressing Therapeutic Proteins

The gene delivery vehicles of the invention can be administered by intravenous delivery or by other methods described herein in order to induce long term *in vivo* expression of a variety of therapeutic proteins. As is demonstrated in Examples 18-24 herein, administration of a retroviral vector intravenously results in sustained, long-term systemic expression of therapeutic genes expressed by the retroviral vector. Thus, methods for obtaining long-term systemic expression *in vivo* of a variety of proteins known to one of

skill in the art are encompassed by the instant invention. Preferably, long-term *in vivo* systemic expression is obtained by intravenous delivery methods as is described in detail below (see Section H of this application). For long term expression from a retroviral vector *in vivo*, the action of human complement on the retroviral vector is suppressed. This can be done by a variety of techniques known to one of skill in the art. Preferably, human packaging cell lines are used in order to inhibit the action of human complement on the retroviral vector particles (see PCT publication number US 91/06852, entitled "Packaging Cells"). The terms "human complement resistant" or "resistance to human complement" when applied to a gene delivery vehicle such as a retroviral vector means that the vectors is at least 70% resistant, more preferably 80% resistant and most preferably at least 90% resistant to inactivation by human complement when tested in a standard serum inactivation assay as described in Example 11 herein.

The production of high titer preparations of recombinant retroviral vectors is described in detail herein. The term "high titer retroviral vector preparation" as used herein refers to a retroviral vector preparation that has a titer greater than 10^6 cfu/ml, more preferably greater than 10^7 cfu/ml, still more preferably greater than 10^8 cfu/ml, more preferably greater than 10^9 cfu/ml, yet more preferably greater than 10^{10} cfu/ml, and most preferably greater than 10^{11} cfu/ml when tested on HT1080 cells in the assay for colony forming units on HT1080 cells as is described herein. This assay depends on the expression of a marker protein to determine cfu/ml. The term "cfu" refers to "colony forming units" when vectors contain a selectable marker. The term refers to identifiable colonies when a phenotypically observable marker is used, such as blue colonies when the marker is beta-galactosidase. In some cases, for example for factor VIII vectors, when no marker gene is present, "cfu" refers to units measured by a PCR-based titer. Briefly, HT1080 cells are transduced with the markerless vector and number of proviral DNA copies are measured by quantitative PCR; HT1080 cells transduced with a retroviral vector containing the neomycin resistance gene are used as a standard (see Section F of the Detailed Description, herein).

The terms "long term systemic expression" or "sustained systemic expression" as used herein in reference to *in vivo* expression of protein encoded by a retroviral vector mean measurable or biologically active expression into the bloodstream for 30 days, more

preferably for 60 days, yet more preferably for 90 days, and more preferably for six months, still for preferably for 1 year, more preferably for at least 5 years after administration of the retroviral vector particle to a host. Most preferably, the expression would be for the life of the patient such that the retroviral vector would only need to be administered to the patient
5 once, or that "booster" injections would have to be delivered to the patient only a few times during the lifetime of the patient. The term "measurable expression" as used herein in reference to *in vivo* expression of a protein encoded by a retroviral vector means that the protein is produced in sufficient amounts so as to be detectable in biological fluids such as serum by an assay specific for the expressed protein. The term "systemic expression" as
10 used herein means that the proteins are expressed into the circulation and are thus useful for treatment of certain diseases. A variety of diseases discussed in detail below are amenable to treatment by this type of gene therapy.

For example, measurement of human growth hormone can be determined by an ELISA assay specific for human growth hormone protein as described in Example 19
15 herein. The term "biologically active expression" or "protein expression in biologically or therapeutically active amounts" when used herein in reference to *in vivo* expression of a protein encoded by a retroviral vector means that protein is produced in sufficient amounts so as to be detectable in a functional assay. For example, expression of factor VIII can be measured in a clotting assay as described in Example 18, herein. Similarly other expressed
20 proteins can be measured by specific assays for each particular protein that are known to those of skill in the art.

Long-term *in vivo* expression of a variety of proteins can be effected by the methods of the invention, preferably by *in vivo* administration of high titer retroviral vectors as described herein. A large variety of different proteins can be expressed for therapeutic applications in
25 a number of different disease states. Preferred proteins include, cytokines and immune system modulators, hormones, growth factors, vaccine antigens, and proteins for treatment of inherited diseases.

1. Gene Delivery Vehicles Expressing Human Factor VIII and Factor IX for Treatment of Hemophilia

Preparation of retroviral vectors expressing a B-domain deleted factor VIII protein are described in detail in the Example section of this application (see Examples 1, 2, 27 and 28 herein). In particular, these examples demonstrate the insertion of a particular factor VIII deletion called the SQN deletion the construction of which is described in Example 2 herein, and which is further described in detail in PCT WO 91/09122.

The B domain separates the second and third A domains of factor FVIII in the newly synthesized single-chain molecule. The B domain extends from amino acids 712 to 1648 according to Wood *et al.*, 1984, *Nature* 312:330-337. Proteolytic activation of factor VIII involves cleavage at specific Arg residues located at positions 372, 740, and 1689. Cleavages of plasma factor VIII by thrombin or Factor Xa at Arg 372 and Arg 1689 are essential for factor VIII to participate in coagulation. Therefore, activated factor VIII consists of a heterodimer comprising amino acids residues 1-372 (containing the A1 domain) and residues 373-740 (containing the A2 domain), and residues 1690-2332 (containing the A3-C1-C2 domain). An important advantage in using the B domain deleted FVIII molecule is that the reduced size appears to be less prone to proteolytic degradation and therefor, no addition of plasma-derived albumin is necessary for stabilization of the final product. The term "B domain deletion" as used herein with respect to factor VIII protein refers to a factor VIII protein in which some or all removal of some or all of the amino acids between residues 711 and 1694 have been deleted, and which still preserves a biologically active FVIII molecule.

A range of B domain deletions can exist depending on which amino acid residues are in the B domain is deleted and whereby the biological activity of the FVIII molecule is still preserved. A specific B domain deletion called the SQN exists which is created by fusing Ser 743 to Gln 1638 (Lind *et al.*, 1995, *Eur J. Biochem* 323:19-27, and PCT WO 91/09122) This deletes amino acid residues 744 to 1637 from the B domain creating a Ser-Glu-Asn (SQN) link between the A2 and A3 FVIII domains. When compared to plasma-derived FVIII, the SQN deletion of the B domain of FVIII did not influence its *in vivo* pharmacokinetics (Fijnvandraat, et. al., *P.R.Schattauer Verlagsgesellschaft mbH (Stuttgart)*

77:298-302, 1997). The terms "Factor VIII SQN deletion" or "SQN deletion" as used herein refer to this deletion and to other deletions which preserve the single S-Q-N tripeptide sequence and which result in the deletion of the amino acids between the two B-domain SQN sequences (See PCT WO 91/09122 for a description of this amino acid sequence).

There are number of other B-domain deleted forms of factor VIII. cDNA's encoding all of these B-domain deleted factor VIII proteins can be inserted into retroviral vectors by using standard molecular biology techniques similar to those described in Examples 1, 2, 27 and 28 herein. For example cDNA molecules encoding the following B-domain factor VIII deletions can be constructed as described below:

Eaton (1986) Biochemistry 25:8343	des 797-1562 deletion
Toole (1986) PNAS 83:5939	des 760-1639 (LA-FVIII)
Meutien (1988) Prot Eng 2:301	des 771-1666 (FVIII del II: missing one thrombin site)
Sarver (1987) DNA 6:553	des 747-1560
Mertens (1993) Br J Haematol 85:133	des 868-1562 des 713-1637 (thrombin resistant)
Esmon (1990) Blood 76:1593	des 797-1562
Donath (1995) Biochem J 312:49	des 741-1668
Webb (1993) BBRC 190:536	PCR cloned from mRNA
Lind (1995) Eur J Biochem 232:19	des 748-1648 (partially processed) des 753-1648(partially processed des 777-1648(partially processed) des 744-1637 (FVIII-SQ) des 748-1645 (FVIII-RH) des B-domain + 0, 1 ,2 Arg (partially processed) desB,+3Arg (FVIIIIR4) desB, +4Arg (FVIIIIR5)
Langner (1988) Behring Inst Mitt 16-25	des 741-1689 des 816-1598

Cheung (1996) Blood 88:325a	des 746-1639
Pipes (1996) Blood 88:441a	des 795-1688 (thrombin sites mutated)

A B domain deletion in which an IgG hinge region has been inserted can also be used. For instance, a deletion of this type can be obtained from plasmid PSVF8-t β 2. PSVF8-t β 2 is a plasmid, which was designed to link the heavy and light chains with a short hinge region from immunoglobulin A (see Figure 37). To obtain cleavage at the end of the heavy chain and to release the light chain, some residues of the β domain were included on either side of the hinge sequence. The 5' untranslated leader and signal peptide are from the human Factor VIII:C cDNA, with the Kozak consensus sequence at the initiation codon as in pSVF8-302. A description of this vector is included in Chapman *et al.*, U.S. Patent No. 5,595,886. The 3' untranslated region is the same fused Factor VIII and tPA sequence as found in pSVF8-80K.

The construction was completed in two steps: an oligomer with cohesive ends for EcoRI and BclI (117 bp) was cloned into a transfer vector, pF8GM7, the DNA sequence of the oligomer was checked by m13 subcloning and Sanger sequencing.

Next, the final plasmid was assembled by ligation of the following three fragments:

- (a) FspI-EcoRI fragment from pSVF8-92S;
- (b) EcoRI-NdeI fragment of the transfer vector pF8GM7 with oligomer; and
- (c) FspI-NdeI fragment of pSVF8-80K.

Descriptions of pSVF8-92S and pSVF8-80K are included in Chapman *et al.*, U.S. Patent No. 5,595,886. The sequence linking the heavy and light chains is shown in Figure 38.

Three additional B domain-deleted factor VIII constructs of particular interest for inclusion in the gene delivery vehicles of the invention can be prepared as follows. Plasmid pSVF8-500 encodes a factor VIII protein with amino acids 770 to 1656 of the full length Factor VIII deleted, according to the nomenclature used in Seq ID No. 45. In addition the threonine at position 1672 of the full-length factor VIII sequence of Seq ID No. 45 was also deleted. The following is a description of the construction of the vector.

The pSVF8-500 plasmid is a derivative of pSVF8-302 in which the regions coding for the 92K and 80K domains are fused with a small connecting β -region of 21 amino acids, retaining the natural proteolytic processing sites. This plasmid was constructed in the following manner:

- 5 (1) A Sall-KpnI fragment of 1984 bp containing the region coding for the 92K protein (except for the carboxyl terminal end) and BstXI-Sall fragment of 2186 bp containing the region coding for the carboxyl end of the 80K protein with 3' end untranslated region were isolated by gel electrophoresis after digestion of pSVF8-302 with restriction enzymes.
- 10 (2) A BclI-BstXI fragment of 1705 bp containing most of the region coding for the 80K protein was isolated after gel electrophoresis of the BamHI-XbaI fragment of pUC12F8. (pUCF812 is prepared from pF8-102 which is described in U.S. Patent No. 5,045,455. pF8-102 is digested with Bam-XbaI and ligated into vector pUC12 by *in vitro* mutagenesis at a BclI site using the following primer: 5' ACT ACT CTT CAA TCT GAT
15 CAA GAG GAA 3' (Seq ID No. 52).
- (3) A KpnI-EcoRI fragment containing the carboxyl end of the 92K protein and part of the β region (4 amino acids) was obtained by digestion of the Sall cassette from pSVF8-302 with KpnI and EcoRI.
- (4) Ligation of four pieces of synthetic DNA (shown in Figure 39) to the
20 fragments of steps (2) and (3) and digestion with KpnI.
- (5) Final ligation of fragments from steps (1) and (4); digestion with Sall and gel purification of the 6428 bp Sall cassette.
- (6) Ligation of the Sall cassette into pSV7d vector; transformation of HB101 and colony hybridization to isolate pSVF8-500 (Figure 40). The sequence of the junction region
25 coding for 92K- β -80K was verified by DNA sequence after cloning in M13.

Note that the sequence in the linker shown in Figure 39 is different from wild type human Factor VIII. The sequence was changed to incorporate unique NruI and MluI restriction sites without changing the amino acid sequence. These sites were also used to construct other two additional B-domain deleted vectors which are described below.

pSV500B Δ Thr was constructed from pSVF8-500. The threonine deletion at position 1672 was maintained. The synthetic linker shown in Figure 41 was used to construct pSV500B Δ Thr. The linker extends from a unique NruI site at Ser(765) to a unique MluI site at Ile(1659) in the pSVF8-500 vector. This linker was substituted for the corresponding region of pSVF8-500.

A third vector pSVF8-500B was constructed from pSV500B Δ Thr. This vector is identical to pSVF8-500B except that the codon for threonine 1672 was re-inserted using standard mutagenesis methods. The relationship between, pSVF8-500B, pSVF8-500B, is further illustrated in the table below. Amino acid sequence numbers in the table were determined by reference to full-length factor VIII sequence depicted in Seq. ID No. 45.

Name	Amino Acids Deleted	Thr at 1672 Deletion
pSVF8-500	770 to 1656	Yes
pSVF8-500B Δ Thr	779 to 1658	Yes
pSVF80-500B	779 to 1658	No

In all cases, the BglII-PfII 1.35 kb fragments of each modified cDNA listed above can be inserted into the retroviral vectors described herein using standard molecular biology procedures known to those of skill in the art and described herein. For instance, cDNA clones containing the B domain-deleted factor VIII proteins can be excised and used to replace the BglII to PfII fragment in the retroviral vectors described in examples 27 and 28 herein. A similar procedure can be used to construct other retroviral vectors, including those described in section A, above.

The full-length factor VIII cDNA can also be inserted into the retroviral vectors of the invention as described in Example 2 herein and in WO 96/21035 which is hereby incorporated by reference in its entirety. In addition, the full-length human factor VIII cDNA can be derived from the construct pSV7dF8-200 or pSV7dF8-300 described in Example 28, herein. This can be constructed similarly to the retroviral vector expressing the B-domain deleted factor VIII described in Example 28 where the cDNA fragment is cloned into the BglII/PfII linearized vector. For instance, an AccI to XbaI 5.9 kb fragment can replace the 3.15 kb AccI to XbaI fragment of the modified cDNA in F8:4213. The

construction of the retroviral vector can then essentially follow the steps described in Example 28. A variety of Factor VIII deletions, mutations, and polypeptide analogs of Factor VIII can also be introduced into the gene delivery vehicles of the invention including retroviral vectors by modifications of the procedures described herein. These analogs
5 include, for instance, those described in PCT Patent Publication Nos. WO 97/03193, WO 97/03194, WO 97/03195, and WO 97/03191, all of which are hereby incorporated by reference.

Other vectors expressing B domain deleted factor VIII or full-length factor VIII can be constructed using techniques known to those of skill in the art or described herein. For
10 instance, adenoviral vectors expressing B-domain deleted factor VIII molecules can be constructed as described in WO 94/29471. Gene delivery vehicles derived from adenovirus are also provided, in which the factor VIII gene, or its various derivatives as described herein, is inserted into a variety of adenovirus-derived vectors, and subsequently recombinant vector particles are generated as described. The adenovirus vectors of the
15 invention also include the following vectors constructed to express factor VIII proteins.

The prototype recombinant adenovirus vectors were deleted in the early region one, [E1a/E1b, (E1)] region, rendering them replication defective. Following insertion of a gene of interest into the deleted region, propagation of the recombinant E1-deleted adenovirus vector is accomplished in 293 cells, a complementing human embryonic kidney cell line
20 stably transformed with the Ad E1 region, which provide these Ad gene products in *trans*. Recombinant Ad vectors generated in this fashion can yield preparations with titers between 10^{11} to 10^{13} particles/ml (Reviewed in Berkner, 1988, *Biotechniques* 6:616-629). However there are several drawbacks to this prototype Ad vector system, including: (1) restriction of heterologous genetic material to approximately 4.5 to 5.0 kb, and (2) partial replication
25 competence of the E1-deleted Ad vectors (Rich, 1993, *Hum. Gen. Ther.* 4:461-476). This later point arises in part to a complementing "E1-like activity" that is expressed in human cells, and results in the expression of other viral gene products present in these vectors, including the highly immunogenic, "late", or structural gene products (e.g. penton protein). As a result of immune responses of the recipient to Ad-specific proteins expressed by the

E1-deleted vectors, expression of the heterologous gene, or transgene is transient and associated with the development of pathology at the site of gene transfer.

Thus, second generation Ad vectors have sought to further "cripple" the capacity of the vector to replicate, and express viral-specific gene products, and to increase the capacity of heterologous genetic material. Such vectors have been of three types: (1) E1 and E3 genes deleted [Bett, *PNAS*, 91:8802-8806, 1994], (2) E1 and E4 genes deleted (Wang, 1995, *Gene Ther.* 2:775-783, and PCT Patent Publication Nos. WO 96/22378 and WO 95/02697), and (3) deletion of all Ad viral genes, or "gutless" (Fisher, 1996, *Virology*, 217:11-22; Hardy, *J. Virol.*, 71:1842-1849, and Kochanek, 1996, *PNAS*, 93:5731-5736). The duration of transgene expression in animals inoculated with these second generation recombinant adenovirus vectors has been dramatically increased, as a result of the mitigation of the recipient's immune response.

As expected, increased deletion of viral-specific genes in the second generation Ad vectors has also resulted in an increased capacity for heterologous genetic material, thus extending the usefulness of this system for application to human gene transfer. This capacity for heterologous genetic material is approximately 8 kb in the E1/E3 and E1/E4 vectors, and is greater than 30 kb for the "gutless" Ad vectors, permitting the insertion of entire genes, including relevant gene expression control regions.

Generation of recombinant Ad vectors, including the E1/E3, E1/E4, and "gutless" vectors, harboring the full-length factor VIII gene, or its B-domain derived derivatives, can be accomplished according to methods well-known to those skilled in the art. For example: (1) the full-length factor VIII gene, or its B-domain derived derivatives can be inserted into plasmid pBHG11 (Bett, 1994, *PNAS* 91:8802-8806), and recombinant E1/E3-deleted Ad vectors are generated after transfection of 293 cells and subsequent intracellular homologous recombination, (2) the full-length factor VIII gene, or its B-domain derived derivatives can be first substituted into the E1 region of any of a variety of E1-deleted Ad vectors and co-transfected with Cla I digested H5dl1014, and recombinant E1/E4-deleted Ad vectors are generated after transfection of 293-E4 cells [Wang *Gene Ther.*, 2:775-783, (1995)], and subsequent intracellular homologous recombination, (3) the full-length factor VIII gene, or its B-domain derived derivatives is first inserted into the Δ Ad plasmid "

[Fisher, 1996, *Virology*, 217:11-22), along with appropriate amounts of "stuffer" sequence derived from, for example, bacteriophage lambda DNA, to permit efficient packaging of recombinant "gutless" adenovirus vector genomes, transfected onto 293 cells and infected with H5.CBALP helper virus (Yang, 1995, *Virol.* 69:2004-2015). Purification of
5 recombinant "gutless" adenovirus vector particles from helper virus can be accomplished, for example, by centrifugation over a cesium gradient, as a result of a buoyant density lower than that of helper virus.

In the case of adeno-associated viral vectors, preferably a factor VIII heavy chain and light chain are inserted into separate vectors, as is demonstrated in Examples 29 and 30
10 herein. Other adenoviral vectors including those described above and in co-owned U.S. serial number 60/025649 can be constructed to express factor VIII light chain and heavy chain using the techniques described in Examples 29 and 30. Heavy and light chain constructs for inclusion into the AAV vectors of the invention can be prepared for instance as described in Chapman *et al.*, U.S. Patent No. 5,595,886.

15 Hemophilia B can also be treated with systemically administered factor IX-expressing gene delivery vehicles including retroviral vectors. Human factor IX deficiency (Christmas disease or Hemophilia B) affects primarily males because it is transmitted as sex-linked recessive trait. It affects about 2000 people in the US. The human factor gene codes a 416 amino acids of mature protein.

20 The human factor IX cDNA can be obtained for instance by constructing plasmid pHfIX1, as described by Kurachi and Davie, 1982, *PNAS* 79(21):6461-6464. The cDNA sequence can be excised as a PstI fragment of about 1.5 kb, blunt ended using T4 DNA polymerase. The factor cDNA fragment can be readily inserted, for example into a SrfI site introduced into a retroviral vector. For instance the cDNA fragment can be inserted into the
25 SrfI site of a linearized pMBA retroviral vector using the procedure described in Examples 27 and 28 herein. In this case, the resulting retroviral vector: pMB-F9 will be produced from HAI cells and its expression will be determined using HT1080 fibroblasts as target cells.

Similarly, the full-length factor IX cDNA fragment can be introduced into other
30 gene delivery vehicles described herein, such as adenoviral vectors and AAV vectors. In the

case of adneoviral vectors, vectors expressing factor IX can be produced, for instance, as described in. For instance, adenoviral vectors expressing B-domain deleted factor VIII molecules can be constructed as described in WO 94/29471. Adenoviral vectors expressing factor IX protein can be produced by methods similar to those described above for factor VIII protein.

In the case of AAV vectors, a factor IX cDNA can be inserted into vectors using, for instance, procedures similar to those described in Examples 29 and 30 herein for factor VIII heavy and light chains. Other adenoviral vectors including those described above and in co-owned U.S. provisional application No. 60/025,649 can be constructed to express factor VIII light chain and heavy chain using the techniques similar to those of Examples 29 and 30.

2. Gene delivery vehicles expressing proteins for treatment of other hereditary disorders

There are a number of proteins useful for treatment of hereditary disorders that can be expressed *in vivo* by the methods of invention. Many genetic diseases caused by inheritance of defective genes result in the failure to produce normal gene products, for example, thalassemia, phenylketonuria, Lesch-Nyhan syndrome, severe combined immunodeficiency (SCID), hemophilia, A and B, cystic fibrosis, Duchenne's Muscular Dystrophy, inherited emphysema and familial hypercholesterolemia (Mulligan *et al.*, 1993, *Science* 260:926; Anderson *et al.*, 1992, *Science* 256:808; Friedman *et al.*, 1989, *Science* 244:1275). Although genetic diseases may result in the absence of a gene product, endocrine disorders, such as diabetes and hypopituitarism, are caused by the inability of the gene to produce adequate levels of the appropriate hormone insulin and human growth hormone respectively.

Gene therapy by the methods of the invention is a powerful approach for treating these types of disorders. This therapy involves the introduction of normal recombinant genes into somatic cells so that new or missing proteins are produced inside the cells of a patient. A number of genetic diseases have been selected for treatment with gene therapy, including adenine deaminase deficiency, cystic fibrosis, α_1 -antitrypsin deficiency,

Gaucher's syndrome, as well as non-genetic diseases. As an example of the present invention, a retroviral vector can be used to treat Gaucher disease. Gaucher disease is a genetic disorder that is characterized by the deficiency of the enzyme glucocerebrosidase. This enzyme deficiency leads to the accumulation of glucocerebroside in the lysosomes of all cells in the body. For a review see *Science* 256:794, 1992 and *The Metabolic Basis of Inherited Disease*, 6th ed., Scriver, et al., vol. 2, p. 1677).

As described in detail above, long term expression of Factor VIII or Factor IX is useful for treatment of blood clotting disorders, such as hemophilia. Different forms of Factor VIII, such as the B domain deleted Factor VIII construct described in Example 2 herein can be used to produce retroviral vectors expressing Factor VIII for use in the methods of the invention. In addition to clotting factors, there are a number of proteins which can be expressed in the retroviral vectors of the invention and which are useful for treatment of hereditary diseases. These include lactase for treatment of hereditary lactose intolerance, AD for treatment of ADA deficiency, and alpha-1 antitrypsin for treatment of alpha-1 antitrypsin deficiency. See F.D. Ledley, 1987, *J. Pediatrics* 110:157-174; I. Verma, *Scientific American* (Nov., 1987) pp. 68-84; and PCT Patent Publication WO 95/27512 entitled "Gene Therapy Treatment for a Variety of Diseases and Disorders" for a description of gene therapy treatment of genetic diseases.

One such disorder is familial hypercholesterolemia is a disease characterized clinically by a lifelong elevation of low density lipoprotein (LDL), the major cholesterol-transport lipoprotein in human plasma; Pathologically by the deposition of LDL-derived cholesterol in tendons, skin and arteries leading to premature coronary heart disease; and genetically by autosomal dominant inherited trait. Heterozygotes number about 1 in 500 persons worldwide. Their cells are able to bind cholesterol at about half the rate of normal cells. Their plasma cholesterol levels show two fold elevation starting at birth. Homozygotes number 1 in 1 million persons. They have severe cholesterolemia with death occurring usually before age 20. The disease (Arteriosclerosis) depends on geography. It affects 15.5 per 100,000 individuals in the U.S. (20,000 total) and 3.3 per 100,000 individuals in Japan. Gene delivery vehicles expressing the LDL receptor for treatment of disorders manifesting with elevated serum LDL can be constructed by techniques known to

those of skill in the art. An example of a retroviral vector expressing LDS receptor is shown in example 32 herein. Other retroviral vectors such as those described herein can readily be constructed using PCR amplication and restriction enzyme treatment methods similar to those described in Example 31.

- 5 Similarly, retroviral vectors expressing alpha-1 antitrypsin for treatment of alpha-1 antitrypsin deficiency can be constructed by variations of the procedure demonstrated in Example 32, herein.

Sequences which encode the above-described nucleic acid molecules may be obtained from a variety of sources. For example, plasmids which contain sequences that
10 encode altered cellular products may be obtained from a depository such as the American Type Culture Collection (ATCC, Rockville, Maryland), or from commercial sources such as Advanced Biotechnologies (Columbia, Maryland). Plasmids containing cytokine genes or immunodulatory genes can be digested with appropriate restriction enzymes, and DNA fragments containing the particular gene of interest can be inserted into the gene delivery
15 vector using standard molecular biology techniques. (See Sambrook *et al.*, Molecular Cloning: A Laboratory Manual (2nd Edition, Vols. 1-3, Cold Spring Harbor Laboratory (1989) or Ausbel, et al. ed. Current Protocols in Molecular Biology, Greene Publishing and Wiley-Interscience, New York (1987).

Alternatively, cDNA sequences for use with the present invention may be obtained
20 from cells which express or contain the sequences. Briefly, within one embodiment mRNA from a cell which expresses the gene of interest is reverse transcribed with reverse transcriptase using oligo dT or random primers. The single-stranded cDNA may then be amplified by PCR (*see* U.S. Patent Nos. 4,683,202, 4,683,195 and 4,800,159. See also *PCR Technology: Principles and Applications for DNA Amplification*, Erlich (ed.), Stockton
25 Press, 1989) utilizing oligonucleotide primers complementary to sequences on either side of desired sequences. In particular, a double-stranded DNA is denatured by heating in the presence of heat stable Taq polymerase, sequence-specific DNA primers, ATP, CTP, GTP and TTP. Double-stranded DNA is produced when synthesis is complete. This cycle may be repeated many times, resulting in a factorial amplification of the desired DNA.

Nucleic acid molecules which are carried and/or expressed by the recombinant retroviruses described herein may also be synthesized, for example, on an Applied Biosystems Inc. DNA synthesizer (*e.g.*, APB DNA synthesizer model 392 (Foster City, California)).

5 Within yet another aspect of the invention, the recombinant retrovirus directs the expression of a substance capable of activating an otherwise inactive precursor into an active inhibitor of a pathogenic agent, or a conditional toxic palliative, which are palliatives that are toxic for the cell expressing the pathogenic condition. As should be evident given the disclosure provided herein, a wide variety of inactive precursors may be converted into
10 active inhibitors of a pathogenic agent. For example, antiviral nucleoside analogues such as AZT or ddI are metabolized by cellular mechanisms to the nucleotide triphosphate form in order to specifically inhibit retroviral reverse transcriptase, and thus viral replication (Furmam *et al.*, 1986, *Proc. Natl. Acad. Sci. USA* 83:8333-8337). Recombinant retroviruses which direct the expression of a gene product (*e.g.*, a protein) such as Herpes Simplex Virus
15 Thymidine Kinase (HSVTK) or Varicella Zoster Virus Thymidine Kinase (VZVTK) which assists in metabolizing antiviral nucleoside analogues to their active form are therefore useful in activating nucleoside analogue precursors (*e.g.*, AZT or ddC) into their active form. AZT or ddI therapy will thereby be more effective, allowing lower doses, less generalized toxicity, and higher potency against productive infection. Additional nucleoside
20 analogues whose nucleotide triphosphate forms show selectivity for retroviral reverse transcriptase but, as a result of the substrate specificity of cellular nucleoside and nucleotide kinases are not phosphorylated, will be made more efficacious.

 Within one embodiment of the invention, the HSVTK gene may be expressed under the control of a constitutive macrophage or T-cell-specific promoter, and introduced into
25 macrophage or T-cells. Constitutive expression of HSVTK results in more effective metabolism of nucleotide analogues such as AZT or ddI to their biologically active nucleotide triphosphate form, and thereby provides greater efficacy, delivery of lower doses. less generalized toxicity, and higher potency against productive infection. Additional nucleoside analogues whose nucleotide triphosphate forms show selectivity for retroviral
30 reverse transcriptase but, as a result of the substrate specificity of cellular nucleoside and

nucleotide kinases are not phosphorylated, may also be utilized within the context of the present invention.

Within a related aspect of the present invention, gene delivery vehicles such as recombinant retroviruses are provided which direct the expression of a substance that
5 activates another compound with little or no cytotoxicity into a toxic product in the presence of a pathogenic agent, thereby effecting localized therapy to the pathogenic agent. In this case, expression of the gene product from the recombinant retrovirus is limited to situations wherein an entity associated with the pathogenic agent, such as an intracellular signal identifying the pathogenic state, is present, thereby avoiding destruction of nonpathogenic
10 cells. This cell-type specificity may also be conferred at the level of infection, by targeting the recombinant retrovirus to cells having or being susceptible to the pathogenic condition.

Within a related aspect of the present invention, gene delivery vehicles such as recombinant retroviruses are provided which direct the expression of a gene product(s) that activates a compound with little or no cytotoxicity into a toxic product. Briefly, a wide
15 variety of gene products which either directly or indirectly activate a compound with little or no cytotoxicity into a toxic product may be utilized within the context of the present invention. Representative examples of such gene products include HSVTK and VZVTk which selectively monophosphorylate certain purine arabinosides and substituted pyrimidine compounds, converting them to cytotoxic or cytostatic metabolites. More
20 specifically, exposure of the drugs ganciclovir, acyclovir, or any of their analogues (e.g., FIAC, DHPG) to HSVTK, phosphorylates the drug into its corresponding active nucleotide triphosphate form.

C. Tissue-specific promoters

Although not an absolute requirement for the practice of the invention, in a further
25 embodiment, the gene delivery vehicles of the invention can contain a liver specific promoter to maximize the potential for liver specific expression of the exogenous DNA sequence contained in the vectors. Preferred liver specific promoters include the hepatitis B X-gene promoter and the hepatitis B core protein promoter. These liver specific promoters are preferably employed with their respective enhancers. The enhancer element

can be linked at either the 5' or the 3' end of the nucleic acid encoding the therapeutic molecule. The hepatitis B X gene promoter and its enhancer can be obtained from the viral genome as a 332 base pair EcoRV-NcoI DNA fragment employing the methods described in Twu, 1987, *J. Virol.* 61:3448-3453. The hepatitis B core protein promoter can be obtained
5 from the viral genome as a 584 base pair BamHI-BglII DNA fragment employing the methods described in Gerlach, 1992, *Virol* 189:59-66. It may be necessary to remove the negative regulatory sequence in the BamHI-BglII fragment prior to inserting it. Other liver specific promoters include the AFP (alpha fetal protein) gene promoter and the albumin gene promoter, as disclosed in EP Patent Publication 0 415,731, the -1 antitrypsin gene
10 promoter, as disclosed in Rettenger, 1994, *Proc. Natl. Acad. Sci.* 91:1460-1464, the fibrinogen gene promoter, the APO-A1 (Apolipoprotein A1) gene promoter, and the promoter genes for liver transference enzymes such as, for example, SGOT, SGPT and -glutamyle transferase. See also PCT Patent Publications WO 90/07936 and WO 91/02805 for a description of the use of liver specific promoters in retroviral vectors.

15 For the retroviral vectors described herein, such as those described in Examples 1 and 27 herein, a liver specific promoter as described in this section can be introduced into the vector to operably linked to gene of interest (for example factor VIII) in order to induce more liver specific expression of the protein. In the case of AAV vectors, the promoter is operably linked to the nucleic acid encoding the therapeutic molecule upstream from the
20 latter and between the AAV vector sequences (for example between the inverted terminal repeats in psub201 or downstream of the Double D ITR sequence).

Examples of the construction of retroviral vectors expressing α interferon under the control of liver-specific promoters is shown in Example 31 herein. Similar techniques can be used to construct retroviral vectors expressing other proteins under control of liver-
25 specific promoters, for instance, by using a three-way ligation procedure analogous to the procedure described in Example 31.

D. Use of gene delivery vectors co-expressing therapeutic protein and a prodrug-converting enzyme:

The gene delivery vehicles and the therapeutic methods of delivery are useful for the long term expression of therapeutic proteins for treatment of a variety of disorders described herein. Particularly, because of the long term treatment made possible by the present invention, it can be desirable for the gene delivery vehicles of the invention to co-express a
5 prodrug converting enzyme which converts a non-cytotoxic compound into its cytotoxic counterpart. The prototype of such enzymes is herpes simplex thymidine kinase (HSVTK) which converts the prodrug gancyclovir into a toxic compound.

Techniques for introducing HSVTK or other prodrug converting enzymes into gene delivery vehicles, particularly retroviral vectors are well known to those of skill in the art.
10 (See eg. WO 91/02805, WO 90/07936, and WO 95/14091). Procedures for preparing gene delivery vehicles, particularly retroviral vectors which co-express a therapeutic protein and HSVTK or other prodrug converting enzymes are described herein and are also described in number of publications (See, eg. WO 93/10218.) These procedures can readily be used by those of skill in the art to introduce the HSVTK gene or a gene encoding another prodrug
15 converting enzyme into the gene delivery vehicles of the invention. In addition, the prodrug converting enzyme can be introduced into a second vector particle which is then co-administered with the gene delivery vehicle expressing the therapeutic protein. An example of this approach is described in WO 96/21015.

20 E. Pretreatment of target tissues:

Retroviral vectors are known to preferentially infect dividing cells. (See Miller *et al.*, 1990, *Molecular Cell Biol.* 10:4239.) There are variety of techniques that may be used to increase the number of dividing cells in target tissues and thereby enhance the efficiency of target cell infection by the retroviral vectors of the invention. For example, growth
25 factors may be used to stimulate target tissues to enter the portion of the cell cycle in which retroviral vector integration can take place. Such growth factors and their target tissues include, but are not limited to, the following:

Protein S and Gas6 acting on nervous system cells and smooth muscle cells;
thrombin acting on smooth muscle cells, gastrointestinal epithelium, liver fat storage cells,
30 dental pulp cells, fibroblasts, endothelial cells, mesangial cells, and astrocytes; coagulation

Factor Xa acting on smooth muscle cells; nerve growth factor acting on nervous system cells; CSF-1 acting on placenta and endometrium; IGF-1 acting on kidney, bone, skin, adipose tissue, airway smooth muscle cells, gastrointestinal epithelium, neural tissue, muscle, and follicular cells; insulin acting on gastrointestinal epithelium, skin, and adipose tissue; KGF acting on urothelium, mammary epithelium, skin, liver, and gastrointestinal epithelium; TGF acting on gastrointestinal epithelium, dental pulp, neural tissue, fibroblasts, connective tissue, inner ear sensory epithelium, colon, bone, pneumocyte type II cells, cornea, and smooth muscle cells; endothelin acting on kidney, smooth muscle cells, melanocytes, cardiac muscle, and astrocytes; PDGF acting on kidney, airway smooth muscle cells, gastrointestinal epithelium, neural tissue, and connective tissue; EGF acting on kidney, skin, neural tissue, inner ear sensory epithelium, connective tissue, fibroblasts, endometrium, liver, and intestine; HGF acting on liver, kidney, mammary epithelium, gastrointestinal epithelium, alveolar epithelium, melanocytes, placenta, and alveolar type II cells; PSA acting on prostate, breast, lung, colon, ovary, liver, and kidney; injurin and HGF-activators acting on liver, kidney, and mammary epithelium; FGF acting on neural cells, kidney, endothelium, fibroblasts, skin, skeletal muscle, connective tissue, melanocytes, cornea, bone marrow, dental pulp cells, liver, melanocytes, smooth muscle cells, and thyroid follicular cells; VEGF acting on endothelial cells; Arg-vasopressin acting on liver, kidney, and fibroblasts; thyroid hormones acting on bone; azoxymethane acting on the colon; prostaglandins acting on liver and dental pulp; IL1 acting on fibroblasts; IL2 acting on T cells; IL15 acting on muscle; triiodothyronine acting on liver; LIF acting on muscle and bone; amphiregulin acting on the skin; soluble thrombomodulin acting on fibroblasts; stem cell factor acting on erythroid progenitors; osteogenic protein 1 acting on chondrocytes, and bone; bone morphogenic protein acting on liver; MGF acting on melanocytes; MGSA acting on melanocytes; heregulins acting on mammary epithelium, keratinocytes, and Schwann cells; and melanotropin acting on melanocytes. Growth factors can also be used in combination, particularly but not limited to mixtures consisting of one or several of EGF, IGF, PDGF, FGF, or KGF. In particular, multiple growth factors known to stimulate cell division of a particular target tissue can be used in combination to increase the proportion of dividing cells in the tissue. The actions of the above and other growth

factors can also be potentiated with substances including but not limited to dextran sulfate, heparin, and other sulfated glycosaminoglycans, FBP, leukotrienes, prostaglandins, oleic acid, HGF activators, androgens, estrogens, ethanol, PF4, and TGF beta antagonists.

Treatment with growth factors or the other substances described above can occur by
5 administering the substances *in vivo* or can also be used in other treatment modalities such as *ex vivo* treatment.

The polypeptide growth factors described herein can be administered in a variety of forms including full-length growth factors, growth factor fragments, truncated growth factors, growth factor fusion proteins and growth factor analogues. Growth factor fusion
10 proteins include fusion proteins with the full-length growth factor, truncated growth factor, growth factor analogue, and growth factor fragment. As used herein the term "growth factor fragment" refers to any growth factor polypeptide that contains less than a full-length sequence, and which retains sufficient biological activity to be used in the methods of the invention. The term "growth factor analogue" as used herein refers to growth factors,
15 truncated growth factors and growth factor fragments with amino acid substitutions, deletions, additions, and modifications; and retaining the biological activity of the growth factor. Thus, the term "growth factor analogue" as used herein includes splice variants, truncations, variants, alleles and derivatives of the mature protein. Analogues possess one or more of the bioactivities of the full length protein. Thus, polypeptides that are identical
20 or contain at least 60%, preferably 70%, more preferably 80%, and most preferably 90% amino acid sequence homology to the amino acid sequence of the mature protein wherever derived, from human or nonhuman sources, are included within this definition.

Growth factor analogues include variants of the growth factors described herein. Growth factor variants contain amino acid substitutions, deletions, or insertions. The amino
25 acid substitutions can be conservative amino acid substitutions or substitutions to eliminate non-essential amino acid residues such as to alter a glycosylation site, a phosphorylation site, an acetylation site, or to minimize misfolding by substitution or deletion of one or more cysteine residues that are not necessary for function. Conservative amino acid substitutions are those that preserve the general charge, hydrophobicity/hydrophilicity and/or steric bulk
30 of the amino acid substituted, for example, substitutions between the members of the

following groups are conservative substitutions: Gly/Ala, Val/Ile/Leu, Asp/Glu, Lys/Arg, Asn/Gln, Ser/Thr/Cys and Phe/Trp/Tyr. The analogs herein further include peptides having one or more peptide mimics, also known as peptoids, that possess the bioactivity of the protein. Included within the definition are also polypeptides containing one or more analog amino acid (including, for example, unnatural amino acids, etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring. The term polypeptide also does not exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like.

10 A large variety of different forms of the various growth factors described above are known to those of skill in the art and can be used in the methods of the invention. For example, a variety of PDGF polypeptides are known in the art. For instance, nucleic acid vectors encoding naturally occurring PDGF and the production of recombinant PDGF are described in U.S. Patent No. 5,219,759, issued June 15, 1993, entitled "Recombinant DNA
15 encoding PDGF A-chain Polypeptide Expression Vectors". In addition to the naturally occurring PDGF polypeptides, a variety of PDGF analogues are known in the art and can be used in the methods described herein. (See eg. U.S. Patent No. 5,149,792, issued September 22, 1992, entitled "Platelet-Derived Growth Factor B Chain Analogues"; U.S. Patent No. 5,128,321, issued July 7, 1992, entitled "PDGF Analogues and Methods of Use"; and U.S.
20 Patent No. 4,849,407, issued July 18, 1989, and entitled "Biologically Active Mosaic Proteins".

 Different forms of IGF-1 and IGF-2 growth factor polypeptides are also well known the art. For example, recombinant human, as well as specific DNA sequences and expression vectors for the production of human IGF-1 are described in European Patent No.
25 0123228B1, grant published on September 19, 1993, entitled "Hybrid DNA Synthesis of Mature Insulin-like Growth Factors". A variety of other forms of IGF-1 polypeptides are known in the art and can readily be produced for use in the methods of the invention. See eg. U.S. Patent No. 5,019,500, entitled "59 Valine Insulin-Like Growth Factor I and Process for Production Thereof"; PCT Patent Publication No. WO 93/23067 entitled "IGF-1
30 Analogues"; European Patent No. EP0128733B1, entitled " Human Insulin-Like Growth

Factor (IGF) Produced from a Human Host, Process, Expression Vector and Recombinant Host Therefor., and IGF-containing Pharmaceutical Composition".

FGF growth factors encompass two families of growth factors found in a variety of different tissues and which are mitogenic factors for a number of different tissues. These two families of polypeptides have been termed acidic FGF and basic FGF. A large variety of different FGF polypeptides, analogues and fragments are known in the art and can be used in the methods of the present invention. See, *eg.* U. S. Patent No. 5,464,774, issued Nov. 7, 1995, entitled "Bovine Basic Fibroblast Growth Factor"; U.S. Patent No. 5,155,214, entitled Basic Fibroblast Growth Factor, and U. S. Patent No. 4,994,559, entitled "Human Fibroblast Growth Factor".

Different forms of keratinocyte growth factor that are useful in the methods of the invention are also known to those of skill in the art. See, for example, a preferred truncated form of KGF which is described in PCT Patent Publication No. 95/10434, published January 12, 1995, and entitled "A Truncated Form of KGF Having Increased Biological Activity". Similarly, different forms of EGF polypeptides are known in the art. See *eg.* PCT publication No. WO 90/08771 and U.S. Patent No. 5,096,825, issued March 17, 1992, entitled Gene for Human Epidermal Growth Factor and Synthesis of Expression Thereof.

The growth factor polypeptides, fragments and analogues used in the instant invention can be produced in a variety of ways, including isolation of PDGF polypeptide from naturally occurring sources, polypeptide chain synthesis by peptide synthesis methods and production of recombinant proteins. These methods are well known to those of skill in the art and are further described herein. For examples of the production of recombinant PDGF polypeptides and analogues, see U.S. Patent No. 5,045,633, issued Sept. 3, 1991, entitled "Expression of Biologically Active PDGF Analogues in Eucaryotic Cells"; U.S. Patent No. 4,769,328, issued Sept. 6, 1988, entitled "Expression of Biologically Active PDGF Analogues in Yeast"; and U.S. Patent No. 4,801,542, issued January 31, 1989, entitled "Expression of Biologically Active PDGF Analogues in Eucaryotic Cells". For examples of the production of FGF polypeptides see *eg.* U.S. Patent No. 5,229,501, entitled "Expression and Production of Human Fibroblast Growth Factor Receptor", U.S. Patent No. 5,331,095, entitled "Process for Purification and Production of Basic Fibroblast

Growth Factor"; and U.S. Patent No. 5,143,829, entitled "High Level Expression of Basic FGF Having a Homogeneous N-terminus".

Liver is an attractive organ for gene therapy because it is easily accessible via the circulation and is the source of a variety of proteins involved in genetic disorders, including factor VIII. Gene therapy targeting the liver using the retroviral vectors of the invention
5 can be performed with or without pretreatment to include benign hyperplasia of the liver. Pretreatment to induce benign liver hyperplasia can be effected, for example, by treatment with hepatocyte growth factor (HGF) and/or transforming growth factor alpha. (See eg. Liu et al., 1994, *Hepatology* 19:1521.)

10 A variety of different forms of HGF useful to induce liver cell proliferation are known in the art. See eg. European Patent Publication No. EP 461560, published December 12, 1991, and entitled "Recombinant Human Leukocyte Derived Hepatocyte Growth Factor- with DNA encoding it, Recombinant Expression Vectors, and Transformant Cells Expressing it." HGF can also be produced and administered to induce liver

15 proliferation in vivo as is described in Joplin et al., 1992, *J. Clin. Invest.* 90:1284.

Liver cells can also be stimulated by administration of agents that mediate or potentiate the activation of endogenous HGF. HGF is produced as a single chain protein that is inactive as a growth factor. Single chain HGF is subsequently cleaved into a two-chain form which is the biologically active growth factor. Enzymes which are shown to
20 convert single-chain HGF to its biologically active form are useful for inducing liver cell proliferation. Therefore, these enzymes can be administered either alone or in combination with exogenous HGF to enhance liver proliferation. Examples of such enzymes are coagulation factor XIIa (see Shiomura et al., 1995, *European J. of Biochem* 229:257); HGF activator (see Shiomura et al., 1992, *Cytotechnology* 8:219); HGF converting enzyme
25 (Mizuno et al., 1994, *Biochem. Biophys. Res. Comm.* 198:1161); and urokinase and tissue plasminogen activators (Mars et al., 1993, *Am J. Pathol.* 143:949). For example, urokinase can be co-administered with HGF. HGF and urokinase could be either be co-formulated or mixed immediately prior to injection. If HGF and urokinase were co-formulated, they could, for example, be stored at low pH in order to minimize the activity of urokinase.

The effects of either HGF or TGF α on liver stimulation have also been shown to be enhanced by prior administration of collagenase (Liu, 1994, *Hepatology* 19:1521). Agents that stimulate release of collagenase-like materials from monocytic cell types, such as LPS or endotoxin-like substances (Beauchamp, 1994, *Surgery* 116:637) may also be used. Nonenzymatic adjuncts and costimulants of growth factors may also be used, including estrogens with HGF (Ni, 1994, *Hepatology* 19:183), heparin or other sulfated glycosaminoglycans with HGF (Yamakazi, 1996, *Cytokine* 8:178; Matsumoto, 1996, *Biochem Biophys Res Commun* 227:455), spermine or spermidine with HGF (Higaki, 1994, *Gastroenterology* 106:1024), nicotinamide with EGF (Wu (1994) *Cancer Res* 54:5964), or choline (Tessitore, 1997, *Biochem J.* 322:151). The phosphofetuin/phospho- α -2-HS glycoprotein pathway downregulates liver regeneration. Inhibitors of this pathway may be used to upregulate liver proliferation. Such inhibitors include IL6, IL1 α , or anti-fetuin antibodies (Ohnishi, 1997, *Eur. J. Biochem.* 243:753). Combinations of these agents can also be applied along with the various growth factors described above.

In addition to growth factor therapy, liver cell growth can also be stimulated by nutritional manipulation. A variety of different nutritional regimens can be used. For instance, a period of protein deprivation followed by consumption of a high protein meal can be used to stimulate DNA synthesis and cell division. (See Mead *et al.*, 1990, *Cancer Res.* 50:7023-7030.)

Another example of stimulation of cell division in a particular tissue is the use of cyclooxygenase inhibitors, such as indomethacin, to induce hyperplasia in the gastric mucosa. In particular, indomethacin is known to increase DNA synthesis and cell clearance in duodenal and jejunal mucosa (see Uribe *et al.*, 1992, *Dig. Dis. Sci.* 37:403-408). Thus, pretreatment with indomethacin or other non-steroidal anti inflammatory drugs can be used to increase cell proliferation in the gastric mucosa. In addition, prostaglandins, in particular, prostaglandin E2 can be used after introduction of the gene therapy vehicle in order to increase the retention of the transduced intestinal cells.

Yet another example of a method to increase the number of dividing cells in gastric mucosa is the administration omeprazole. Omeprazole can be obtained and administered as described in Kakei *et al.*, 1995, *Biochem. Biophys. Res. Comm.* 214:861-867.

An additional class of drugs that is known to induce liver proliferation is the peroxisome proliferators. These are a set of structurally diverse drugs that cause an increase in the number of peroxisomes present in hepatocytes, but which also lead to hepatocyte mitosis mimicking regeneration through unknown mechanism(s). Some of these have found clinical utility as hypolipidemic drugs, particularly clofibrate (Reddy and Lalwai, 1983, *CRC Crit Rev Toxicol* 12:1-58), and act through activation of members of the superfamily of nuclear steroid receptors known as PPAR (peroxisome proliferator-activated receptors) and subsequent increased transcription of peroxisomal beta-oxidative enzymes (Isseman and Green, 1990, *Nature* 347:645-650), and possibly through down-regulation of the endoplasmic reticulum protein, BiP (Motojima & Goto, 1992; *FEBS Lett* 308:207-210). These drugs include clofibrate (Sigma Chemical Compan, St. Louis, MO, USA), WY 16463 (Chem Syn, Lenexa, KS, USA), DEHP (Aldrich, Milwaukee WI, USA), DHEA (Sigma), PFDA (Aldrich), PFOA (Aldrich), fenofibrate (Laboratories Fournier, Daix, France), gemfibrozil (Lopid, Warner-Lambert , Ann Arbor MI), (Sigma), clofibric acid (Sigma), bezafibrate (Sigma), methyl clofenapate, tibric acid, BR 931(4-chloro-6(2,3-xylylidino)2-pyrimidinylthio-(N-beta-hydroxy-ethyl)acetamide), DEHA (di(2ethylhexyl)adipate), nafenopin, clinofibrate.

Other classes of nonproteinaceous substances known to stimulate liver proliferation may be employed. These include 9-cis-retinoic acid (Ohmura, 1996, *Life Sciences* 58:PL211), the glycolipid, hepatopoietin B, (Michalopoulos, 1990, *FASEB J* 4:176), cyclosporine A (Masuhara, 1993, *Carcinogenesis* 14:1579), phosphatidylethanolamine N-methyltransferase inhibitors such as 3-deazaadenosine or other methylation inhibitors (Chiang, 1979, *Biochem Pharmacol* 28:1897), nonsteroidal antiinflammatory drugs that activate PPAR such as flufenamic acid, fenoprofen, and ibuprofen (Lehmann, 1997, *J. Biol. Chem.* 272:3406); or apoptosis-inducing molecules (Busch, 1984, *Carcinogenesis* 5:453; Colambano, 1985, *Lab Invest* 52:670) such as anti-Fas antibody, dexamethasone, etoposide, camptothecin, staurosporin, hypericin, S-nitrosoglutathione, taxol, 4-hydroxyphenyl retinamide, prostaglandin A2, delta-12-PGJ2, sulindac sulfone, actinomycin D, beta-lapachone, TPEN, vinblastine, vincristine, and A23187. Triiodothyronine can also be

administered as a stimulatory compound. Combinations of these substances can also be applied along with the peptide growth factors described herein.

F. CELL CULTURE

5 As noted above, the present invention provides high titer recombinant retroviral preparation suitable for administration to humans. In order to produce such high titer preparations, cell culture methods as described below are provided in order to enable the production of high titer recombinant retroviruses. Briefly, a wide variety of methods may be utilized, including for example, the use of fermenters or bioreactors, roller bottles, cell
10 hotels or cell factories, and hollow fiber culture.

 In particular, for bioreactors or fermenters, cells are preferably grown on microcarriers (*i.e.*, Cytodex 1 or Cytodex 2; Pharmacia, Piscataway, N.J. at concentrations ranging from 3 to 15 g/L microcarrier. Suitable media, and growth conditions are described by way of a representative illustration in Example 17.

15 For roller bottles, suitable conditions include those described above for bioreactors, with the exception that microcarrier beads are not utilized. Generally, cells are grown in 850 cm² roller bottles ("FALCON" Corning, Corning New York) containing DMEM media, along with 15%-20% FBS. Preferably, the bottles are sealed to avoid contamination, although "open" bottles may also be utilized under appropriate conditions (*e.g.*, 5% CO₂).
20 Generally, the roller bottles are incubated at a temperature of 37°C, with a rotation speed of 0.5 rpm/minute.

 Cell factories may also be utilized for the large scale cell culture and production of recombinant retroviruses. Briefly, cell factories (also termed "cell hotels") typically contain 2, 10, or 40 trays, are molded from virgin polystyrene, treated to provide a Nuclon D
25 surface, and assembled by sonic welding one to another. Generally, these factories have two port tubes which allow access to the chambers for adding reagents or removing culture fluid. A 10-layer factory provides 6000cm² of surface area for growing cells, roughly the equivalent of 27 T-225 flasks. Cell factories are available from a variety of manufacturers, including for example Nunc.

Most cell types are capable of producing high titer vector into the media for 3-6 days, allowing for multiple harvests. Each cell type is tested to determine the optimal harvest time after seeding the culture and optimal number of harvest days. Cells are typically initially grown in DMEM supplemented with 2-20% FBS in roller bottles until the
5 required number of cells for seeding a cell factory is obtained. Cells are then seeded into the factories and 2 liters of culture supernatant containing vector is harvested from each day for four days. Fresh DMEM/FBS is used to replenish the cultures.

Within other aspects of the present invention, hollow fiber culture methods are provided for the production of recombinant retroviruses. Briefly, high titer retroviral
10 production using hollow fiber cultures is based on increasing viral concentration as the cells are being cultured to a high density in a reduced volume of media. Cells are fed nutrients and waste products are diluted using a larger volume of fresh media which circulates through the lumen of numerous capillary fibers. The cells are cultured on the exterior spaces of the capillary fibers in a bioreactor chamber where cell waste products are
15 exchanged for nutrients by diffusion through 30,000 Dalton pores in the capillary fibers. Retroviruses which are produced from the cell lines are too large to pass through the 30,000 Dalton pore membrane, and thus concentrate in the hollow fiber bioreactor along side of the cells. The volume of media being cultured on the cell side is approximately 10 to 100 fold lower then volumes required for equivalent cell densities cultured in tissue culture dishes or
20 flasks. This decrease fold in volume inversely correlates with the fold induction of titer when hollow fiber retroviral titers are compared to tissue culture dishes or flasks. This 10-100 fold induction in titer is seen when an individual retroviral producer cell line is amiable to hollow fiber growth conditions. To achieve maximum cell density, the individual cells must be able to grow in very close proximity and on top of each other. Many cell lines will
25 not grow in this fashion and retroviral packaging cell lines based on these types of cell lines may not achieve 10 fold increases in titer. Cell lines which would grow very well would be non-adherent cell line and it is believed that a retroviral producer line based on a non-adherent cell line may reach 100 fold increases in titer compared to tissue culture dishes and flasks.

The harvest procedure, in its original design, is a procedure which uses syringes to evacuate and replace culture sups to harvest the produced vector. This syringe procedure has an associated high risk of possible contamination, which requires a significant number manual connections and disconnection's of media flow paths. However a convenient
5 procedure has been devised which will reduce the risk of contaminating the cultures, increase the daily volumes which can harvested and reduce the time required to handle the culture system. The harvesting procedure is now performed using a batch peristaltic pump drive to deliver precise volumes of fresh media to replace equal volumes of harvest material which is then delivered through thin line tubing into a collection bottle stored at 40C. The
10 pump batch sequence is activated by a timer which can be set at specific time points and the pump can be adjusted to harvest any set volume of harvest material, twenty-four hours a day. The collected supernatant can then be frozen, pooled with earlier harvests, or processed as described elsewhere. This collection procedure can be used for any hollow fiber system including the Cellco (Rockville, Maryland), Unisyn (Tustin, CA), or Cellex
15 (Coon Rapids, MN) systems including ceramic matrix high density culture systems.

G. CONCENTRATION AND PURIFICATION OF RECOMBINANT RETROVIRAL PARTICLES

As noted above, the present invention provides methods for concentrating and purifying recombinant retroviruses, in order to increase the purity of therapeutic preparation, as well as to increase the titer of recombinant retrovirus that may be given. A wide variety
20 of methods may be utilized for increasing viral concentration and purity, including for example, precipitation of recombinant retroviruses with ammonium sulfate, polyethylene glycol ("PEG") concentration, concentration by centrifugation (either with or without gradients such as PERCOLL, or "cushions" such as sucrose, use of concentration filters (e.g., Amicon filtration), and 2-phase separations. Each of these methods will be discussed
25 in more detail below.

Briefly, in order to accomplish concentration by precipitation of recombinant retroviruses with ammonium sulfate, ammonium sulfate is added slowly to an appropriate concentration, followed by centrifugation and removal of the ammonium sulfate either by dialysis or by separation on a hydrophobic column. One difficulty with this method

however, is that in addition to concentration of recombinant retroviruses, other proteinaceous debris may also be concentrated.

Alternatively, recombinant retroviruses may be concentrated from culture medium with PEG (Green *et al.*, 1970, *PNAS* 67:385-393; Syrewicz *et al.*, 1972, *Appl. Micro.* 24:488-494). Such methods are rapid, simple, and inexpensive. However, like ammonium sulfate precipitation, use of PEG also concentrates other proteins from solution.

Within other embodiments, recombinant retroviruses may be concentrated by centrifugation, and more particularly, low speed centrifugation. Briefly, low speed centrifugation allows concentration of recombinant retroviruses, while avoiding the difficulties associated with pelleting that accompanies high speed centrifugation (*e.g.*, virus destruction or inactivation). Particularly preferred methods for concentrating viruses by low-speed centrifugation are described below in more detail in Example 15.

Within yet other aspects of the invention, recombinant retroviruses may be concentrated by an aqueous two-phase separation method. Briefly, polymeric aqueous two-phase systems may be prepared by dissolving two different non-compatible polymers in water. Many pairs of water-soluble polymers may be utilized in the construction of such two-phase systems, including for example polyethylene glycol ("PEG") or methylcellulose, and dextran or dextran sulfate (see Walter and Johansson, 1986, *Anal. Biochem.* 155:215-242; Albertsson, "Partition of Cell Particles and Macromolecules" Wiley, New York, 1960). As described in more detail below in Example 13, utilizing PEG at concentrations ranging from 5% to 8% (preferably 6.5%), and dextran sulfate at concentrations ranging from 0.4% to 1% (preferably 0.4%), an aqueous two-phase system may be established suitable for purifying recombinant retroviruses. Utilizing such procedures, approximately 1.4 liters of crude research grade supernatant may be reduced to a 10 mL volume, while recovering approximately 50% of the total starting retrovirus.

For purposes of illustration, one representative concentration process which combines several concentration steps is set forth below. Briefly, recombinant retroviruses may be prepared either from roller bottles, cell factories, or bioreactors prior to concentration. Preferably, daily harvests of recombinant retroviruses from producer cells is preferred, followed by addition of fresh media. Removed media containing the recombinant

retrovirus may be frozen at -70°C, or more preferably, stored at 2°C to 8°C in large pooled batches prior to processing.

For material obtained from a bioreactor, the recombinant retrovirus pool is first clarified through a 0.8 um filter (1.2 um glass fiber pre-filter, 0.8 um cellulose acetate) connected in series with a 0.65 um filter (Sartorius). This filter arrangement provides approximately 2 square feet of filter, and allows processing of about 15-20 liters of pooled material before clogging. For material obtained from roller bottles or cell factories, a single 0.65um cartridge (2 sq. ft.) normally suffices for volumes up to 40 liters. For 80 liter cell factory processes, a 5 sq. ft. filter may be required.

Preferably, after clarification, the filter is rinsed with buffer (150 mM NaCl, 25 mM Tris, pH 7.2-7.5). This step has allowed recoveries of recombinant retrovirus ranging from 80% to 120%.

Following clarification, recombinant retroviruses are concentrated by tangential flow ultrafiltration utilizing Filtron units and Sigma Screen cassettes with a 300,000 mw cut off.

For bioreactor material (containing 12% to 16% FBS), 4 to 5 liters of material may be concentrated per cassette. For roller bottles or cell factories at 12-16% FBS, 5-6 liters of material may be concentrated per cassette. Finally, for cell factories containing 10% FBS, 8 to 9 liters of material may be concentrated per cassette. Utilizing a pressure differential of 2 psi between filtrate (8 psi) and retentate (10 psi), up to 80 liters of material may be concentrated to a volume of less than 500 ml in under two hours. This process also provides a yield of about 80%.

Following the ultrafiltration step, DNase is added to a concentration of 50 U/ml, and recirculated at a lower pump speed with the filtrate line closed for 30 minutes. If retroviruses have been trapped within a gel layer formed during the ultrafiltration, this step will break down trapped retrovirus, and improve recovery.

Discontinuous diafiltration is then accomplished by addition of 4 liters of additional buffer, and utilizing the same cross differential pressure set forth above. Generally, recovery after this step is approximately 70%.

Concentrated material is then subjected to column chromatography on a Pharmacia S-500 HG size exclusion gel, utilizing 50 mM NaCl and 25 mM Tris pH 7.2-7.5 as

minimum salt and ionic strength concentrations. Generally, recombinant retroviruses elute off in the first peak.

Tangential flow filtration may once again be utilized in order to further reduce the volume to under 200 ml. Finally, the concentrated material is sterilized by filtration through
5 a 0.2 um Millipore filter (PVDF, or Sterivex).

H. ASSAYS

Within other aspects of the present invention, methods are provided for quantitating retroviral particles utilizing non-denaturing gels (e.g., 4-15% gradient polyacrylamide gels),
10 along with methods for estimating or quantitating the resultant products such as, for example, staining with coomassie blue or silver stain, followed by densitometry scanning. Such methods, while not capable of discriminating between viable and non-viable vector particles, are advantageous because they are relatively simple and quick. One representative example of such methods is set forth below in Example 10 in more detail.

15 Within other aspects of the present invention, assays are provided for titering recombinant retrovirus in a sample. Typically, such assays may be based upon presence of a selectable marker, or formation of blue colonies. However, within certain embodiments recombinant retroviruses are provided which do not include a gene coding for a selectable marker. Therefore, antibody and PCR assays, the latter of which is described below, may be
20 employed in order to determine retrovirus titer. To use PCR to amplify sequences unique to the recombinant retroviruses described herein, various primers are required. Such primers can readily be designed by those skilled in the art and will depend on the retroviral vector backbone employed and the components thereof, the particular region(s) desired to be amplified, *etc.* Representative examples of particular primer pairs include those specific for
25 LTR sequences, packaging signal sequences or other regions of the retroviral backbone, include primers specific for the nucleic acid molecule (*i.e.*, non-heterologous sequence) of interest.

Briefly, within one embodiment of the invention a PCR titering assay is performed by growing a known number of cells, transduced with a recombinant retrovirus on 6-well
30 plates for at least 16 hr. before harvest. One well per plate is sacrificed for counting. Cells

from the other wells are lysed and their contents isolated. DNA is prepared using a QUIAmp DNA isolation kit (QUIAgen, Inc., Chatsworth, CA). DNAs are resuspended in 5×10^6 cell equivalents/ μ l per sample.

To calculate titer, a standard curve is generated using DNA isolated from 5×10^6 untransduced HT1080 cells (negative control) and 5×10^6 HT1080 cells transduced with a known vector and having one copy of that vector per cell genome (positive control), such as may be prepared from packaging cell lines transduced with a recombinant retrovirus encoding a selectable marker, *e.g.*, neomycin resistance. The standard curve is generated by combining different amounts of the positive and negative control DNA and amplifying specific sequences therefrom by PCR using primers specific to a particular region of the recombinant retrovirus. A representative group of mixtures for generating a standard curve is:

<u>Tube</u>	<u>100%</u>	<u>75%</u>	<u>50%</u>	<u>25%</u>	<u>10%</u>	<u>5%</u>	<u>0%</u>	<u>Blank</u>
15 Positive Control (μ l)	50	37.5	25	12.5	5	2.5	0	0
Negative Control (μ l) 0	12.5	25	37.5	45	47.5	50	0	
Distilled water (μ l)	0	0	0	0	0	0	0	50

Five microliters from each tube is placed into one of eight reaction tubes (duplicates are also prepared), with the remainder being stored at -20°C . Five microliters from each sample DNA preparations are placed into their own reaction tubes in duplicate. PCR reactions (50 μ l total volume) are then initiated by adding 45.0 μ l of a reaction mix containing the following components per tube to be tested: 24.5 μ l water, 5 μ l 10X reaction PCR buffer, 4 μ l of 25 mM MgCl_2 , 4 μ l dNTPs (containing 2.5 mM of each of dATP, dGTP, dCTP, and dTTP), 5 μ l of primer mix (100 ng of each primer), 0.25 μ L TaqStart monoclonal antibody (Clontech Laboratories, Inc., Palo Alto, CA), 1.00 μ L TaqStart buffer (Clontech Labs, Inc.), and 0.25 μ L AmpliTaq DNA polymerase (Perkin-Elmer, Inc., Norwalk, CN). Just prior to aliquoting the reaction mix to the reaction tubes, 1 μ L of α - ^{32}P dCTP (250 μCi ; 3000 C/mmol, 10 mCi/mL, Amersham Corp., Arlington Heights, IL) is added into the reaction mix. After aliquoting 45.0 μ L the reaction mix into each of the reaction tubes, the tubes are

capped and placed into a thermocycler. The particular denaturation, annealing, elongation times and temperatures, and number of thermocycles will vary depending on size and nucleotide composition of the primer pair used. 20 - 25 amplification thermocycles are then performed. 5 μ L of each reaction is then spotted on DE81 ion exchange chromatography paper (Whatman, Maidstone, England) and air dried for 10 min. The filter is then washed five times, 100 mL per wash, in 50 mM Na_2PO_4 , pH 7, 200 mM NaCl, after which it is air dried and then sandwiched in Saran Wrap. Quantitation is performed on a PhosphorImager SI (Molecular Dynamics, Sunnyvale, CA). Filters are typically exposed to a phosphor screen, which stores energy from ionizing radiation, for a suitable period, typically about 120 min. After exposure, the phosphor screen is scanned, whereby light is emitted in proportion to the radioactivity on the original filter. The scanning results are then downloaded and plotted on a log scale as cpm (ordinate) versus percent positive control DNA (abscissa). Titters (infectious units/mL) for each sample are calculated by multiplying the number of cells from which DNA was isolated by the percentage (converted to decimal form) determined from the standard curve based on the detected radioactivity, divided by the volume of recombinant retrovirus used to transduce the cells. As will be appreciated by those in the art, other methods of detection, such as colorimetric methods, may also be employed to label the amplified products.

20 I. FORMULATION

Within other aspects of the present invention, methods are provided for preserving an infectious recombinant retrovirus, such that the recombinant retrovirus is capable of infecting mammalian cells upon reconstitution (see U.S. Serial No. 08/153,342). Briefly, recombinant retrovirus which has been purified or concentrated as described above may be preserved by first adding a sufficient amount of a formulation buffer to the media containing the recombinant retrovirus, in order to form an aqueous suspension. The formulation buffer is an aqueous solution that contains a saccharide, a high molecular weight structural additive, and a buffering component in water. As utilized within the context of the present invention, a "buffering compound" or "buffering component" should be understood to refer

to a substance that functions to maintain the aqueous suspension at a desired pH. The aqueous solution may also contain one or more amino acids.

The recombinant retrovirus can also be preserved in a purified form. More specifically, prior to the addition of the formulation buffer, the crude recombinant retrovirus described above may be clarified by passing it through a filter, and then concentrated, such as by a cross flow concentrating system (Filtron Technology Corp., Northborough, MA). Within one embodiment, DNase is added to the concentrate to digest exogenous DNA. The digest is then diafiltrated to remove excess media components and establish the recombinant retrovirus in a more desirable buffered solution. The diafiltrate is then passed over a Sephadex S-500 gel column and a purified recombinant retrovirus is eluted. A sufficient amount of formulation buffer is added to this eluate to reach a desired final concentration of the constituents (see, *e.g.*, Example 9) and to minimally dilute the recombinant retrovirus, and the aqueous suspension is then stored, preferably at -70°C or immediately dried. As noted above, the formulation buffer is an aqueous solution that contains a saccharide, a high molecular weight structural additive, and a buffering component in water. The aqueous solution may also contain one or more amino acids.

The crude recombinant retrovirus can also be purified by ion exchange column chromatography. This method is described in more detail in U.S. Patent Application Serial No. 08/093,436. In general, the crude recombinant retrovirus is clarified by passing it through a filter, and the filtrate loaded onto a column containing a highly sulfonated cellulose matrix. The recombinant retrovirus is eluted from the column in purified form by using a high salt buffer. The high salt buffer is then exchanged for a more desirable buffer by passing the eluate over a molecular exclusion column. A sufficient amount of formulation buffer is then added, as discussed above, to the purified recombinant retrovirus and the aqueous suspension is either dried immediately or stored, preferably at -70°C.

The aqueous suspension in crude or purified form can be dried by lyophilization or evaporation at ambient temperature. Specifically, lyophilization involves the steps of cooling the aqueous suspension below the glass transition temperature or below the eutectic point temperature of the aqueous suspension, and removing water from the cooled suspension by sublimation to form a lyophilized retrovirus. Briefly, aliquots of the

formulated recombinant retrovirus are placed into an Edwards Refrigerated Chamber (3 shelf RC3S unit) attached to a freeze dryer (Supermodulyo 12K). A multistep freeze drying procedure as described by Phillips et al. (Cryobiology 18:414, 1981) is used to lyophilize the formulated recombinant retrovirus, preferably from a temperature of -40°C to -45°C.

- 5 The resulting composition contains less than 10% water by weight of the lyophilized retrovirus. Once lyophilized, the recombinant retrovirus is stable and may be stored at -20°C to 25°C, as discussed in more detail below.

- Within the evaporative method, water is removed from the aqueous suspension at ambient temperature by evaporation. Within one embodiment, water is removed through spray drying (EP 520,748). Within the spray drying process, the aqueous suspension is delivered into a flow of preheated gas, usually air, whereupon water rapidly evaporates from droplets of the suspension. Spray drying apparatus are available from a number of manufacturers (e.g., Drytec, Ltd., Tonbridge, England; Lab-Plant, Ltd., Huddersfield, England). Once dehydrated, the recombinant retrovirus is stable and may be stored at -20°C to 25°C. Within the methods described herein, the resulting moisture content of the dried or lyophilized retrovirus may be determined through use of a Karl-Fischer apparatus (EM Science Aquastar™ V1B volumetric titrator, Cherry Hill, NJ), or through a gravimetric method.

- The aqueous solutions used for formulation, as previously described, are composed of a saccharide, high molecular weight structural additive, a buffering component, and water. The solution may also include one or more amino acids. The combination of these components act to preserve the activity of the recombinant retrovirus upon freezing and lyophilization, or drying through evaporation. Although a preferred saccharide is lactose, other saccharides may be used, such as sucrose, mannitol, glucose, trehalose, inositol, fructose, maltose or galactose. In addition, combinations of saccharides can be used, for example, lactose and mannitol, or sucrose and mannitol. A particularly preferred concentration of lactose is 3%-4% by weight. Preferably, the concentration of the saccharide ranges from 1% to 12% by weight.

- The high molecular weight structural additive aids in preventing viral aggregation during freezing and provides structural support in the lyophilized or dried state. Within the

context of the present invention, structural additives are considered to be of "high molecular weight" if they are greater than 5000 m.w. A preferred high molecular weight structural additive is human serum albumin. However, other substances may also be used, such as hydroxyethyl-cellulose, hydroxymethyl-cellulose, dextran, cellulose, gelatin, or povidone.

5 A particularly preferred concentration of human serum albumin is 0.1% by weight. Preferably, the concentration of the high molecular weight structural additive ranges from 0.1% to 10% by weight.

The amino acids, if present, function to further preserve viral infectivity upon cooling and thawing of the aqueous suspension. In addition, amino acids function to further

10 preserve viral infectivity during sublimation of the cooled aqueous suspension and while in the lyophilized state. A preferred amino acid is arginine, but other amino acids such as lysine, ornithine, serine, glycine, glutamine, asparagine, glutamic acid or aspartic acid can also be used. A particularly preferred arginine concentration is 0.1% by weight. Preferably, the amino acid concentration ranges from 0.1% to 10% by weight.

15 The buffering component acts to buffer the solution by maintaining a relatively constant pH. A variety of buffers may be used, depending on the pH range desired, preferably between 7.0 and 7.8. Suitable buffers include phosphate buffer and citrate buffer. A particularly preferred pH of the recombinant retrovirus formulation is 7.4, and a preferred buffer is tromethamine.

20 In addition, it is preferable that the aqueous solution contain a neutral salt which is used to adjust the final formulated recombinant retrovirus to an appropriate iso-osmotic salt concentration. Suitable neutral salts include sodium chloride, potassium chloride or magnesium chloride. A preferred salt is sodium chloride.

Aqueous solutions containing the desired concentration of the components described

25 above may be prepared as concentrated stock solutions.

A particularly preferred method of preserving recombinant retroviruses in a lyophilized state for subsequent reconstitution comprises the steps of (a) combining an infectious recombinant retrovirus with an aqueous solution to form an aqueous suspension, the aqueous suspension including 4% by weight of lactose, 0.1% by weight of human serum

30 albumin, 0.03% or less by weight of NaCl, 0.1% by weight of arginine, and an amount of

tromethamine buffer effective to provide a pH of the aqueous suspension of approximately 7.4, thereby stabilizing the infectious recombinant retrovirus; (b) cooling the suspension to a temperature of from -40°C to -45°C to form a frozen suspension; and (c) removing water from the frozen suspension by sublimation to form a lyophilized composition having less than 2% water by weight of the lyophilized composition, the composition being capable of infecting mammalian cells upon reconstitution. It is preferred that the recombinant retrovirus be replication defective and suitable for administration into humans upon reconstitution.

As illustrated in Figures 1 and 2, mannitol and lactose lyophilized recombinant retrovirus formulations were assayed for preservation of viral activity under various storage temperatures as a function of time. Similarly, Figure 3 illustrates the results of assays of trehalose recombinant retrovirus formulations for preservation of viral activity under various storage temperatures as a function of time. Figure 4 depicts a comparison of the viral infectivity of frozen formulated recombinant retrovirus (-80°C) as a liquid and the viral infectivity of lyophilized recombinant retrovirus stored at -20°C. Mannitol formulations may lose considerable activity upon lyophilization (5-6 fold), but appear to remain stable subsequent to the lyophilization event. Although not preferable, such a loss is acceptable assuming sufficient amounts of retrovirus are present in the aqueous solution.

It will be evident to those skilled in the art given the disclosure provided herein that it may be preferable to utilize certain saccharides within the aqueous solution when the lyophilized retrovirus is intended for storage at room temperature. More specifically, it is preferable to utilize disaccharides, such as lactose or trehalose, particularly for storage at room temperature.

The lyophilized or dehydrated retroviruses of the subject invention may be reconstituted using a variety of substances, but are preferably reconstituted using water. In certain instances, dilute salt solutions which bring the final formulation to isotonicity may also be used. In addition, it may be advantageous to use aqueous solutions containing components known to enhance the activity of the reconstituted retrovirus. Such components include cytokines, such as IL-2, polycations, such as protamine sulfate, or other components which enhance the transduction efficiency of the reconstituted retrovirus. Lyophilized or

dehydrated recombinant retrovirus may be reconstituted with any convenient volume of water or the reconstituting agents noted above that allow substantial, and preferably total solubilization of the lyophilized or dehydrated sample.

J. ADMINISTRATION

5 As noted above, high titer recombinant retroviral particles of the present invention may be administered to a wide variety of locations including, for example, into sites such as the cerebral spinal fluid, bone marrow, joints, arterial endothelial cells, rectum, buccal/sublingual, vagina, the lymph system, to an organ selected from the group consisting of lung, liver, spleen, skin, blood and brain, or to a site selected from the group consisting of
10 tumors and interstitial spaces. Within other embodiments, the recombinant retrovirus may be administered intraocularly, intranasally, sublingually, orally, topically, intravesically, intrathecally, topically, intravenously, intraperitoneally, intracranially, intramuscularly, or subcutaneously. Other representative routes of administration include gastroscopy, ECRP and colonoscopy, which do not require full operating procedures and hospitalization, but
15 may require the presence of medical personnel.

Considerations for administering the compositions of the present invention include the following:

Intravenous (IV) administration can occur under a variety of protocols known to those of skill in the art. For instance, retroviral vector particles can be formulated for IV
20 administration as described above and administered as a single injection. Alternatively, the retroviral vector particles can be delivered in a multiple injection protocol. An example of a multiple injection protocol is administration for three times a day for several consecutive days or on alternate days. The multiple injection schedule can be carried out over a number of days for example a week or 10 days or two weeks. The injection schedule can also be
25 repeated. The total number of vector particles delivered can be dispersed in varying amounts of formulation buffer. Depending on the volume delivered, the retroviral vectors can be delivered as an injection or as an IV formulation such as an IV drip which can be delivered over a longer period of time. Similarly, the rate of administration can vary. Details of the administration protocol such as the single or multiple injection schedule and volume and

time of delivery can be determined experimentally by those of skill in the art, and will also vary depending on the particular gene of interest to be delivered. IV administration is a preferred route of administration for retroviral vectors expressing secretory proteins such as Factor VIII and human growth hormone (see e.g. examples 18-21 herein).

5 Oral administration is easy and convenient, economical (no sterility required), safe (over dosage can be treated in most cases), and permits controlled release of the active ingredient of the composition (the recombinant retrovirus). Conversely, there may be local irritation such as nausea, vomiting or diarrhea, erratic absorption for poorly soluble drugs, and the recombinant retrovirus will be subject to "first pass effect" by hepatic metabolism
10 and gastric acid and enzymatic degradation. Further, there can be slow onset of action, efficient plasma levels may not be reached, a patient's cooperation is required, and food can affect absorption. Preferred embodiments of the present invention include the oral administration of recombinant retroviruses that express genes encoding erythropoietin, insulin, GM-CSF cytokines, various polypeptides or peptide hormones, their agonists or
15 antagonists, where these hormones can be derived from tissues such as the pituitary, hypothalamus, kidney, endothelial cells, liver, pancreas, bone, hemopoetic marrow, and adrenal. Such polypeptides can be used for induction of growth, regression of tissue, suppression of immune responses, apoptosis, gene expression, blocking receptor-ligand interaction, immune responses and can be treatment for certain anemias, diabetes,
20 infections, high blood pressure, abnormal blood chemistry or chemistries (e.g., elevated blood cholesterol, deficiency of blood clotting factors, elevated LDL with lowered HDL), levels of Alzheimer associated amyloid protein, bone erosion/calcium deposition, and controlling levels of various metabolites such as steroid hormones, purines, and pyrimidines. Preferably, the recombinant retroviruses are first lyophilized, then filled into
25 capsules and administered.

 Buccal/sublingual administration is a convenient method of administration that provides rapid onset of action of the active component(s) of the composition, and avoids first pass metabolism. Thus, there is no gastric acid or enzymatic degradation, and the absorption of recombinant retroviruses is feasible. There is high bioavailability, and
30 virtually immediate cessation of treatment is possible. Conversely, such administration is

limited to relatively low dosages (typically about 10-15 mg), and there can be no simultaneous eating, drinking or swallowing. Preferred embodiments of the present invention include the buccal/sublingual administration of recombinant retroviruses that contain genes encoding self and/or foreign MHC, or immune modulators, for the treatment
5 of oral cancer; the treatment of Sjogren's syndrome via the buccal/sublingual administration of such recombinant retroviruses that contain IgA or IgE antisense genes; and, the treatment of gingivitis and periodontitis via the buccal/sublingual administration of IgG or cytokine antisense genes.

Rectal administration provides a negligible first pass metabolism effect (there is a
10 good blood/lymph vessel supply, and absorbed materials drain directly into the inferior vena cava), and the method is suitable of children, patients with emesis, and the unconscious. The method avoids gastric acid and enzymatic degradation, and the ionization of a composition will not change because the rectal fluid has no buffer capacity (pH 6.8; charged compositions absorb best). Conversely, there may be slow, poor or erratic absorption,
15 irritation, degradation by bacterial flora, and there is a small absorption surface (about 0.05m^2). Further, lipidophilic and water soluble compounds are preferred for absorption by the rectal mucosa, and absorption enhancers (*e.g.*, salts, EDTA, NSAID) may be necessary. Preferred embodiments of the present invention include the rectal administration of recombinant retroviruses that contain genes encoding colon cancer antigens, self and/or
20 foreign MHC, or immune modulators.

Nasal administration avoids first pass metabolism, and gastric acid and enzymatic degradation, and is convenient. In a preferred embodiment, nasal administration is useful for recombinant retrovirus administration wherein the recombinant retrovirus is capable of expressing a polypeptide with properties as described herein. Conversely, such
25 administration can cause local irritation, and absorption can be dependent upon the state of the nasal mucosa.

Pulmonary administration also avoids first pass metabolism, and gastric acid and enzymatic degradation, and is convenient. Further, pulmonary administration permits localized actions that minimize systemic side effects and the dosage required for
30 effectiveness, and there can be rapid onset of action and self-medication. Conversely, at

times only a small portion of the administered composition reaches the brochioli/alveoli, there can be local irritation, and overdosing is possible. Further, patient cooperation and understanding is preferred, and the propellant for dosing may have toxic effects. Preferred embodiments of the present invention include the pulmonary administration of recombinant retroviruses that express genes encoding IgA or IgE for the treatment of conditions such as asthma, hay fever, allergic alveolitis or fibrosing alveolitis, the CFTR gene for the treatment of cystic fibrosis, and protease and collagenase inhibitors such as α -1-antitrypsin for the treatment of emphysema. Alternatively, many of the same types of polypeptides or peptides listed above for oral administration may be used..

10 Ophthalmic administration provides local action, and permit prolonged action where the administration is via inserts. Further, avoids first pass metabolism, and gastric acid and enzymatic degradation, and permits self-administration via the use of eye-drops or contact lens-like inserts. Conversely, the administration is not always efficient, because the administration induces tearing. Preferred embodiments of the present invention include the ophthalmic administration of recombinant retroviruses that express genes encoding IgA or IgE for the treatment of hay fever conjunctivitis or vernal and atopic conjunctivitis; and ophthalmic administration of recombinant retroviruses that contain genes encoding melanoma specific antigens (such as high molecular weight-melanoma associated antigen), self and/or foreign MHC, or immune modulators.

20 Transdermal administration permits rapid cessation of treatment and prolonged action leading to good compliance. Further, local treatment is possible, and avoids first pass metabolism, and gastric acid and enzymatic degradation. Conversely, such administration may cause local irritation, is particularly susceptible to tolerance development, and is typically not preferred for highly potent compositions. Preferred embodiments of the present invention include the transdermal administration of recombinant retroviruses that express genes encoding IgA or IgE for the treatment of conditions such as atopic dermatitis and other skin allergies; and transdermal administration of recombinant retroviruses encoding genes encoding melanoma specific antigens (such as high molecular weight-melanoma associated antigen), self and/or foreign MHC, or immune modulators.

Vaginal administration provides local treatment and one preferred route for hormonal administration. Further, such administration avoids first pass metabolism, and gastric acid and enzymatic degradation, and is preferred for administration of compositions wherein the recombinant retroviruses express peptides. Preferred embodiments of the present invention include the vaginal administration of recombinant retroviruses that express genes encoding self and/or foreign MHC, or immune modulators. Other preferred embodiments include the vaginal administration of genes encoding the components of sperm such as histone, flagellin, etc., to promote the production of sperm-specific antibodies and thereby prevent pregnancy. This effect may be reversed, and/or pregnancy in some women may be enhanced, by delivering recombinant retroviruses vectors encoding immunoglobulin antisense genes, which genes interfere with the production of sperm-specific antibodies.

Intravesical administration permits local treatment for urogenital problems, avoiding systemic side effects and avoiding first pass metabolism, and gastric acid and enzymatic degradation. Conversely, the method requires urethral catheterization and requires a highly skilled staff. Preferred embodiments of the present invention include intravesical administration of recombinant retrovirus encoding antitumor genes such as a prodrug activation gene such thymidine kinase or various immunomodulatory molecules such as cytokines.

Endoscopic retrograde cystopancreatography (ERCP) (goes through the mouth; does not require piercing of the skin) takes advantage of extended gastroscopy, and permits selective access to the biliary tract and the pancreatic duct. Conversely, the method requires a highly skilled staff, and is unpleasant for the patient.

Many of the routes of administration described herein (*e.g.*, into the CSF, into bone marrow, into joints, intravenous, intra-arterial, intracranial intramuscular, subcutaneous, into various organs, intra-tumor, into the interstitial spaces, intra-peritoneal, intralymphatic, or into a capillary bed) may be accomplished simply by direct administration using a needle, catheter or related device. In particular, within certain embodiments of the invention, one or more dosages may be administered directly in the indicated manner: into the cerebral spinal fluid at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; into bone

marrow at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; into joint(s) at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; intravenously at dosages greater than or equal to 10^8 , 10^9 , 10^{10} or 10^{11} cfu; intra-arterially at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; intra-cranially at dosages greater than or equal to 10^9 , 10^{10} or 10^{11} cfu; intra-muscularly at dosages greater than or equal to 10^{10} or 10^{11} cfu; intra-ocularly at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; pulmonarily at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; nasally at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; sub-lingually at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; rectally at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; orally at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; topically at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; vaginally at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; sub-cutaneously at dosages greater than or equal to 10^9 , 10^{10} or 10^{11} cfu; inter-vesically at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; into an organ such as the lung, liver, spleen, skin, blood or brain at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; intra-tumor at dosages greater than or equal to 10^8 , 10^9 , 10^{10} or 10^{11} cfu; intra-peritoneally at dosages greater than or equal to 10^8 , 10^9 , 10^{10} or 10^{11} cfu; into interstitial spaces at dosages greater than or equal to 10^{10} or 10^{11} cfu; intra-lymphatically at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; into a capillary bed at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu; or intrathecally at dosages greater than or equal to 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} or 10^{11} cfu.

Recombinant retrovirus may be delivered to the target from outside of the body (as an outpatient procedure) or as a surgical procedure, where the vector is administered as part of a procedure with other purposes, or as a procedure designed expressly to administer the vector. Other routes and methods for administration include the non-parenteral routes as well as administration via multiple sites.

As demonstrated in Example 23 herein, viral vectors and particularly retroviral vectors preferentially infect liver cells and spleen cells in order to effect sustained

expression of proteins such as factor VIII and human growth hormone. Furthermore, as described herein expression of proteins in the liver is useful therapeutically, for example, for production of proteins secreted by the liver, and for localized production of therapeutic proteins in the liver which can be used in the treatment of liver diseases.

5 As described above, the gene delivery vehicles of the invention, including recombinant retroviruses are preferentially administered intravenously, generally in a peripheral vein which is readily accessible. However, the gene delivery vehicles of the invention may also be delivered into the hepatic artery or in the portal vein in order to have more effective delivery to the liver. There are many methods available for placing catheters
10 into the human hepatic artery or portal vein which are known to those of skill in the art. For instance, cannulation of the portal vein can be accomplished by a simple dissection of the umbilical vein, thereby eliminating a laparotomy (Storer *et al.*, 1966, *Am J. Surg* 111:56-68, and Weigand *et al.*, 1983, *Rec Res Cancer Res* 86:90-92). For example, the ligamentum teres is cut 4 to 5 cm above the umbilicus and a polyethylene catheter is threaded into the
15 umbilical vein until reaching the junction with the left branch of the portal vein. Following the exertion of firm pressure on the catheter to break through the vessel wall, a brisk return of blood through the catheter indicates the proper entry into the portal system.

As an additional example, an alternative procedure is described by Beart *et al.*, 1990, *Arch Surg* 125:897-901. In this procedure, a 5F catheter is placed through a hole in the
20 abdominal wall which then extends through a mesenteric vein and into the portal vein. The catheter is flushed with saline and fixed to the exterior of the abdominal wall. These and other procedures may be used to deliver the gene delivery vehicles of the invention to the liver.

In addition, the gene delivery vehicles of the invention can be delivered by direct injection into the liver using standard medical procedures.

25 I. Formulation and Administration of Growth Factors

As is described herein, the recombinant viruses of the invention can be administered after induction of cell proliferation by a growth factor, or may be co-administered with a growth factor. The growth factors of the invention are administered by parenteral, topical, oral or by local administration. For example, the growth factors are administered

parenterally, eg. intravenously, subcutaneously, intradermally, or intramuscularly. Preferably, the growth factors are administered intravenously. Administration of the therapeutic agent of the invention can be accomplished by, for example, injection, catheterization, laser-created perfusion channels, cannulization, a particle gun, and a pump.

5 The growth factors of the invention are typically administered with a pharmaceutical carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity. Suitable carriers may be large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids,
10 amino acid copolymers, and inactive virus particles. Such carriers are well known to those of ordinary skill in the art. Pharmaceutically acceptable salts can be used therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. A thorough discussion of pharmaceutically acceptable excipients is
15 available in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. 1991). Pharmaceutically acceptable carriers in therapeutic compositions may contain liquids such as water, saline, glycerol and ethanol. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. Typically, the therapeutic compositions are prepared as injectables, either as
20 liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. Liposomes are included within the definition of a pharmaceutically acceptable carrier. The term "liposomes" refers to, for example, the liposome compositions described in U.S. Patent No. 5,422,120, WO 95/13796, WO 94/23697, WO 91/14445 and EP 524,968 B1. Liposomes may be pharmaceutical
25 carriers for the polypeptides of the invention.

 The growth factors of the invention are administered in therapeutically effective amounts. The term "therapeutically effective amount" as used herein and applied to polypeptide growth factors refers to an amount of a growth factor that is capable of stimulating cell division in a target tissue *in vivo*. Stimulation of cell proliferation in a target
30 tissue means that the number of dividing cells in the target tissue is greater than in the

absence of treatment. The precise effective amount for a subject will depend upon the subject's size and health, the nature and extent of the condition being treated, recommendations of the treating physician, and particular growth factor that is used. The effective amount for a given situation can be determined by routine experimentation and will vary from growth factor to growth factor. For example, for HGF, a dose of 1 ug/kg to 2 mg/kg body weight, and more preferably from 10 ug/kg to 200 ug/kg body weight is used. In the case of KGF, a dose of 100 ug/kg to 5 mg/kg body weight, or more preferably a dose of 1 mg/kg to 50 mg/kg body weight is used. Dose amounts for the other growth factors used in the claimed methods are known to those of skill in the art or can readily be determined experimentally.

Clofibrate, or the other proxisome proliferators, can be administered by IP injection (5-500 mg/kg), or orally (5-500 mg/kg). More preferably the dosages are 10-100 mg/kg. A typical dosing schedule is daily administration for 3-10 days. A tapered dosing can alternatively be employed. Following clofibrate dosing, retroviral vectors can be administered, preferably intravenously, at doses ranging from 10^5 to 10^{11} cfu per injection. Injection schedules of one to three times daily, for one to ten days, will be employed. Repeat administrations of retroviral vector with or without repeat clofibrate or growth factor dosing can be performed.

Collagenase can be administered at a dosage of from .05-5 U/kg, more preferably 0.5-2 U/kg (Liu (994) Hepatology 19:1521). Estrogens with HGF or other growth factors can be administered at a dosage of 0.01-20 ug/g, and more preferably 0.5-2 ug/g. Heparin or other sulfated glycosaminoglycans administered with HGF or other growth factors can be administered at a dose of 100-10,000 U/kg, more preferably 2000-6000 U/kg.

9-cis-retinoic acid is administered at dosages of 10-200 mg/kg, more preferably 60-100 mg/kg. Cyclosporine A is delivered at a dose of 0.1-100 mg/kg, more preferably 10-50 mg/kg. Prostaglandins E1 and/or I1; isoproterenol can be administered at dosages of 0.5-100 ug/kg; more preferably 1-10 ug/kg with or without glucagon at a dose of 0.1-10 mg/kg, more preferably 0.5-2 mg/kg. Phosphatidylethanolamine N-methyltransferase inhibitors such as 3-deazaadenosine or other methylation inhibitors are administered at a dose of 1-50 mg/kg, more preferably 5-15 mg/kg. Liver mitogen nafenopin can be administered at a dose

of 0.1- 1000 mg/kg, more preferably 50-300 mg/kg. Cyproterone acetate can be administered at a dose of (0.1- 600 mg/kg, more preferably 10-100 mg/kg. Triiodothyronine can be administered at at dose of 0.1-10 mg/kg, more preferably 0.5-5.0 mg/kg.

The following examples are offered by way of illustration, and not by way of
5 limitation.

EXAMPLE 1

Preparation of Retroviral Vector Backbones

A. Preparation of Retroviral Backbones KT-1 and KT-3B

The Moloney murine leukemia virus (MoMLV) 5' long terminal repeat (LTR) EcoR
10 I-EcoR I fragment, including *gag* sequences, from the N2 vector (Armentano *et al.*, 1987, *J. Vir.* 61:1647-1650; Eglitis *et al.*, 1985, *Science* 230:1395-1398) is ligated into the plasmid SK⁺ (Stratagene, La Jolla, CA). The resulting construct is designated N2R5. The N2R5 construct is mutated by site-directed *in vitro* mutagenesis to change the ATG start codon to
15 ATT preventing *gag* expression. This mutagenized fragment is 200 base pairs (bp) in length and flanked by Pst I restriction sites. The Pst I-Pst I mutated fragment is purified from the SK⁺ plasmid and inserted into the Pst I site of N2 MoMLV 5' LTR in plasmid pUC31 to
20 replace the non-mutated 200 bp fragment. The plasmid pUC31 is derived from pUC19 (Stratagene, La Jolla, CA) in which additional restriction sites Xho I, Bgl II, BssH II and Nco I are inserted between the EcoR I and Sac I sites of the polylinker. This construct is
designated pUC31/N2R5gM.

A 1.0 kilobase (Kb) MoMLV 3' LTR EcoR I-EcoR I fragment from N2 is cloned into plasmid SK⁺ resulting in a construct designated N2R3⁻. A 1.0 Kb Cla I-Hind III
fragment is purified from this construct.

The Cla I-Cla I dominant selectable marker gene fragment from pAFVXM retroviral
25 vector (Kriegler *et al.*, 1984, *Cell* 38:483; St. Louis *et al.*, 1988, *PNAS* 85:3150-3154), comprising a SV40 early promoter driving expression of the neomycin (neo) phosphotransferase gene, is cloned into the SK⁺ plasmid. This construct is designated SK⁺ SV₂-neo. A 1.3 Kb Cla I-BstB I gene fragment is purified from the SK⁺ SV₂-neo plasmid.

KT-3B or KT-1 vectors are constructed by a three part ligation in which the Xho I-Cla I fragment containing the gene of interest and the 1.0 Kb MoMLV 3' LTR Cla I-Hind III fragment are inserted into the Xho I-Hind III site of pUC31/N2R5gM plasmid. This gives a vector designated as having the KT-1 backbone. The 1.3 Kb Cla I-BstB I *neo* gene fragment from the pAFVXM retroviral vector is then inserted into the Cla I site of this plasmid in the sense orientation to yield a vector designated as having the KT-3B backbone.

EXAMPLE 2

ORAL ADMINISTRATION OF RECOMBINANT RETROVIRUSES EXPRESSING FACTOR VIII

A. Construction of Full-Length and B Domain Deleted Factor VIII cDNA RETROVECTOR™

The following is a description of the construction of several retroviral vectors encoding a full-length factor VIII cDNA. Further discussion is also provided in PCT Patent Publication No. WO 96/21035. Due to the packaging constraints of retroviral vectors and because selection for transduced cells is not a requirement for therapy, a retroviral backbone, *e.g.*, KT-1, lacking a selectable marker gene is employed.

1. Production of Plasmid Vectors Encoding Full-Length Factor VIII

A gene encoding full-length factor VIII can be obtained from a variety of sources. One such source is the plasmid pCIS-F8 (*see* EP 0 260 148), which contains a full-length factor VIII cDNA whose expression is under the control of a CMV major immediate-early (CMV MIE) promoter and enhancer. The factor VIII cDNA contains approximately 80 bp of 5' untranslated sequence from the factor VIII gene and a 3' untranslated region of about 500 bp. In addition, between the CMV promoter and the factor VIII sequence lies a CMV intron sequence, or "cis" element. The cis element, spanning about 280 bp, comprises a splice donor site from the CMV major immediate-early promoter about 140 bp upstream of a splice acceptor from an immunoglobulin gene.

More specifically, a plasmid, designated pJW-2, encoding a retroviral vector for expressing full length factor VIII is constructed using the KT-1 backbone from pKT-1.

Briefly, in order to facilitate directional cloning of the factor VIII cDNA insert into pKT-1, the unique Xho I site is converted to a Not I site by site directed mutagenesis. The resultant plasmid vector is then opened with Not I and Cla I. pCIS-F8 is digested to completion with Cla I and Eag I, for which there are two sites, to release the fragment encoding full-length factor VIII. This fragment is then ligated into the Not I/Cla I restricted vector to generate a plasmid designated pJW-2.

2. Construction of a Truncated Factor VIII Retrovector (ND-5)

A plasmid vector encoding a truncation of about 80% (approximately 370 bp) of the 3' untranslated region of the factor VIII cDNA, designated pND-5, is constructed in a pKT-1 vector as follows: As described for pJW-2, the pKT-1 vector employed has its Xho I restriction site replaced by that for Not I. The factor VIII insert is generated by digesting pCIS-F8 with Cla I and Xba I, the latter enzyme cutting 5' of the factor VIII stop codon. The approximately 7 kb fragment containing all but the 3' coding region of the factor VIII gene is then purified. pCIS-F8 is also digested with Xba I and Pst I to release a 121 bp fragment containing the gene's termination codon. This fragment is also purified and then ligated in a three way ligation with the larger fragment encoding the rest of the factor VIII gene and Cla I/Pst I restricted BLUESCRIPT® KS⁺ plasmid (Stratagene, *supra*) to produce a plasmid designated pND-2.

The unique Sma I site in pND-2 is then changed to a Cla I site by ligating Cla I linkers (New England Biolabs, Beverly, MA) under dilute conditions to the blunt ends created by a Sma I digest. After recircularization and ligation, plasmids containing two Cla I sites are identified and designated pND-3.

The factor VIII sequence in pND-3, bounded by Cla I sites and containing the full length gene with a truncation of much of the 3' untranslated region, is cloned as follows into a plasmid backbone derived from a Not I/Cla I digest of pKT-1 (a pKT-1 derivative by cutting at the Xho I site, blunting with Klenow, and inserting a Not I linker (New England Biolabs)), which yields a 5.2 kb Not I/Cla I fragment. pCIS-F8 is cleaved with Eag I and Eco RV and the resulting fragment of about 4.2 kb, encoding the 5' portion of the full-length factor VIII gene, is isolated. pND-3 is digested with Eco RV and Cla I and a 3.1 kb

fragment is isolated. The two fragments containing portions of the factor VIII gene are then ligated into the Not I/Cla I digested vector backbone to produce a plasmid designated pND-

5.

3. Construction of the B-Domain Deleted Vector

5 The precursor DNA for the B-deleted FVIII is obtained from Miles Laboratory. This expression vector is designated p25D and has the exact backbone as pCISF8 above. The Hpa I site at the 3' of the FVIII8 cDNA in p25D is modified to Cla-I by oligolinkers. An Acc I to Cla I fragment is clipped out from the modified p25D plasmid. This fragment spans the B-domain deletion and includes the entire 3' two-thirds of the cDNA. An Acc I to
10 Cla I fragment is removed from the Retrovector™ JW-2 above, and replaced with the modified B-domain deleted fragment just described. This is designated B-del-1.

B. Assay for Factor VIII Expression

1. Assay of KT-ND5 Vector Expression by Transient Packaging and Transduction of Murine Cells

15 Cell lines, L33, (Dennert, USC Comprehensive Cancer Center, Los Angeles, CA, Patek, *et al.*, 1979, *Int. J. of Cancer* 24:624-628) BC10ME (Patek *et al.*, 1982, *Cell Immuno* 72:113, ATCC# TIB85) L33env, and BCenv (L33env and BCenv express HIV-1 IIIBenv, Warner *et al.*, 1991, *AIDS Res. and Human Retrovirus* 7:645), transduced with the KT-ND5 vector, carrying the amphotropic or VSVG envelope protein are examined for the
20 expression of factor VIII. Non-transduced cells are also analyzed for factor VIII expression and compared with KT-ND5 transduced cells to determine the effect of transduction on protein expression.

Murine cell lines, L33-KT-ND5, L33env-KT-ND5, L33env, L33, BC10ME, BC10ME-KT-ND5, BCenv, and BCenv-KT-ND5, are tested for expression of the KT-ND5
25 molecule. Cells are grown to subconfluent density and the supernatant is removed following centrifugation at 200 xg. The samples are diluted and assayed by the Coamatic® Factor VIII assay (KabiVitrum Diagnostica, Molndal, Sweden).

The assay is performed as follows: 100 μ l of culture media are mixed with 200 μ l of 1x working buffer diluted according to the manufacturer's instructions. Fifty μ l of the mixture are prewarmed at 37°C for 3-4 minutes in wells of a microtiter plate. The 50 μ l of factor reagent from the kit, prewarmed to 37°C are added, and the mixture incubated at 37°C for 4 min., after which 50 μ l of chromogenic substrate (7.7mg S2765 plus 0.2 mg of thrombin inhibitor I2581 in a mannitol carrier, diluted to a total of 6.0 ml with sterile water) are added. After a 10 minute incubation at 37°C, 50 μ l of 20% acetic acid are added to stop the reaction. Absorbance at 405 nm is determined against buffer. A standard curve using dilutions of pooled normal human plasma (1.0 IU factor VIII/ml) is used. Serum levels of factor VIII in non-hemophilic patients are in the range of 200 ng/mL.

When this assay is used for patient samples, 9 volumes of blood are mixed with one volume of 0.1 M sodium citrate, at a neutral pH, and centrifuged at 2,000 x g for 5 - 20 min. at 20 - 25°C to pellet cells. Due to heat lability of factor VIII, plasma samples should be tested within 30 min. of isolation or stored immediately at -70°C, although as much as 20% of factor VIII activity may be lost during freezing and thawing.

2. Assay of KT-ND5 Vector Expression by Transient Packaging and Transduction of Human Cells

Cell lines transduced with KT-ND5 are examined for expression of factor VIII. Non-transduced cells are analyzed to compare with KT-ND5 transduced cells and determine the effect that transduction has on expression.

Two human cell lines, JY and JY-KT-ND5 are tested for expression of KT-ND5. Suspension cells grown to 10^6 cells/ml are removed from culture flasks by pipet and pelleted by centrifugation at 200 xg. The supernatant is removed, diluted, and assayed by the Coamatic[®] Factor VIII assay as described above in Example 2B1.

C. Transient Transfection and Transduction of Packaging Cell Lines HX and DA with the Vector Construct KT-ND5

1. Plasmid DNA Transfection

The packaging cell line, HX (WO92/05266), are seeded at 5.0×10^5 cells on a 10 cm tissue culture dish on day 1 with Dulbecco's Modified Eagle Medium (DMEM) and 10% fetal bovine serum (FBS). On day 2, the media is replaced with 5.0 ml fresh media 4 hours prior to transfection. A standard calcium phosphate-DNA co-precipitation is performed by mixing 40.0 μ l 2.5 M CaCl_2 , 10 μ g plasmid DNA, and deionized H_2O to a total volume of 400 μ l. Four hundred microliters of the DNA- CaCl_2 solution is added dropwise with constant agitation to 400 μ l precipitation buffer (50 mM HEPES-NaOH, pH 7.1; 0.25 M NaCl and 1.5 mM $\text{Na}_2\text{HPO}_4\text{-NaH}_2\text{PO}_4$). This mixture is incubated at room temperature for 10 minutes. The resultant fine precipitate is added to a culture dish of cells. The cells are incubated with the DNA precipitate overnight at 37°C. On day 3, the media is aspirated and fresh media is added. The supernatant is removed on day 4, passed through a 0.45 μ l filter, and stored at -80°C.

Alternatively, 29 2 3 cells (WO 92/05266) (these are 293 cells expressing gag and pol) are transfected with the vector DNA and the plasmid pMLP-VSVG (or other VSVG encoding plasmids) to yield VSVG psuedotyped vector particles that are harvested and stored as described above.

2. Packaging Cell Line Transduction

DA (an amphotropic cell line derived from a D17 cell line ATCC No. 183, WO 92/05266) cells are seeded at 5.0×10^5 cells/10 cm tissue culture dish in 10 ml DMEM and 10% FBS, 4 μ g/ml polybrene (Sigma, St. Louis, MO) on day 1. On day 2, 3.0 ml, 1.0 ml and 0.2 ml of the freshly collected virus-containing HX media is added to the cells. The cells are incubated with the virus overnight at 37°C. On day 3, the media is removed and 1.0 ml DMEM, 10% FBS with 800 μ g/ml G418 is added to the plate. Only cells that have been transduced with the vector and contain the neomycin selectable marker will survive. A G418 resistant pool is generated over a period of a week. The pool of cells is dilution cloned by removing the cells from the plate and counting the cell suspension, diluting the

cells suspension down to 10 cells/ml and adding 0.1 ml to each well (1 cell/well) of a 96 well plate (Corning, Corning, NY). Cells are incubated for 14 days at 37°C, 10% CO₂. Twenty-four clones are selected and expanded up to 24 well plates, 6 well plates then 10 cm plates at which time the clones are assayed for expression and the supernatants are collected and assayed for viral titer.

- 5 The titer of the individual clones is determined by infection of HT1080 cells, (ATCC No. CCL 121). On day 1, 5.0×10^5 HT1080 cells are plated on each well of a 6 well microtiter plate in 3.0 ml DMEM, 10% FBS and 4 µg/ml polybrene. On day 2, the supernatant from each clone is serially diluted 10 fold and used to infect the HT1080 cells in
- 10 1.0 ml aliquots. The media is replaced with fresh DMEM, 10% FBS media, and the cells incubated with the vector overnight at 37°C, 10% CO₂. On day 3, selection of transduced cells is performed by replacing the media with fresh DMEM, 10% FBS media containing 800 µg/ml G418. Cells are incubated at 37°C, 10% CO₂ for 14 days at which time G418 resistant colonies are scored at each dilution to determine the viral titer of each clone as
- 15 colony forming units(cfu)/ml.

Using these procedures, cell lines are derived that produce greater than or equal to 10^6 cfu/ml in culture.

The packaging cell line HX is transduced with vector generated from the DA vector producing cell line in the same manner as described for transduction of the DA cells from

- 20 HX supernatant.

Transduction of the DA or HX cells with vectors lacking a *neo* selectable marker (Example 1) was performed as described above. However, instead of adding G418 to the cells on day 3, the cells are cloned by limiting dilution. Titer is analyzed as described above.

25 3. Generation of Producer Cell Line via One Packaging Cell Line

In some situations it may be desirable to avoid using more than one cell line in the process of generating producer lines. In this case, DA cells are seeded at 5.0×10^5 cells on a 10 cm tissue culture dish on day 1 with DMEM and 10% irradiated (2.5 megarads minimum) FBS. On day 2, the media is replaced with 5.0 ml fresh media 4 hours prior to

transfection. A standard calcium phosphate-DNA coprecipitation is performed by mixing 60 μ l 2.0 M CaCl_2 , 10 μ g MLP-G plasmid, 10 μ g KT-ND5 retroviral vector plasmid, and deionized water to a volume of 400 μ l. Four hundred microliters of the DNA- CaCl_2 solution is added dropwise with constant agitation to 400 μ l 2X precipitation buffer (50 mM HEPES-NaOH, pH 7.1, 0.25 M NaCl and 1.5 mM $\text{Na}_2\text{HPO}_4\text{-NaH}_2\text{PO}_4$). This mixture is incubated at room temperature for 10 minutes. The resultant fine precipitate is added to a culture dish of DA cells plated the previous day. The cells are incubated with the DNA precipitate overnight at 37°C. On day 3, the medium is removed and fresh medium is added. The supernatant containing G-pseudotyped virus is removed on day 4, passed through a 0.45 μ l filter and used to infect the DA packaging cell.

DA cells are seeded at 5.0×10^5 cells on a 10 cm tissue culture dish in 10 ml DMEM and 10% FBS, 4 mg/ml polybrene (Sigma, St. Louis, MO) on day 1. On day 2, 2.0 ml, 1.0 ml or 0.5 ml of the freshly collected and filtered G-pseudotyped virus containing supernatant is added to the cells. The cells are incubated with the virus overnight at 37°C. On day 3 the medium is removed and 10 ml DMEM, 10% irradiated FBS with 800 μ g/ml G418 is added to the plate. Only cells that have been transduced with the vector and contain the *neo* selectable marker will survive. A G418 resistant pool is generated over the period of 1-2 weeks. The pool is tested for expression and then dilution cloned by removing the cells from the plate, counting the cell suspension, diluting the cell suspension down to 10 cells/ml and adding 0.1 ml to each well (1 cell/well) of a 96-well plate. Cells are incubated for 2 weeks at 37°C, 10% CO_2 . Twenty-four clones are selected and expanded up to 24-well plates, then 6-well plates, and finally 10 cm plates, at which time the clones are assayed for expression and the supernatants are collected and assayed for viral titer as described above.

D. Detection of Replication Competent Retroviruses (RCR)

25 1. The Extended S^+L^- Assay

The extended S^+L^- assay determines whether replication competent, infectious virus is present in the supernatant of the cell line of interest. The assay is based on the empirical observation that infectious retroviruses generate foci on the indicator cell line MiCl_1 (ATCC No. CCL 64.1). The MiCl_1 cell line is derived from the Mv1Lu mink cell line (ATCC No.

CCL 64) by transduction with Murine Sarcoma Virus (MSV). It is a non-producer, non-transformed, revertant clone containing a replication defective murine sarcoma provirus, S⁺, but not a replication competent murine leukemia provirus, L⁻. Infection of MiCl₁ cells with replication competent retrovirus "activates" the MSV genome to trigger "transformation" which results in foci formation.

Supernatant is removed from the cell line to be tested for presence of replication competent retrovirus and passed through a 0.45 µ filter to remove any cells. On day 1, Mv1Lu cells are seeded at 1.0 x 10⁵ cells per well (one well per sample to be tested) of a 6 well plate in 2 ml DMEM, 10% FBS and 8 µg/ml polybrene. Mv1Lu cells are plated in the same manner for positive and negative controls on separate 6 well plates. The cells are incubated overnight at 37°C, 10% CO₂. On day 2, 1.0 ml of test supernatant is added to the Mv1Lu cells. The negative control plates are incubated with 1.0 ml of media. The positive control consists of three dilutions (200 focus forming units (ffu), 20 ffu and 2 ffu each in 1.0 ml media) of MA virus (referred to as pAM in Miller *et al.*, *Molec. and Cell Biol.* 5:431, 1985) which is added to the cells in the positive control wells. The cells are incubated overnight. On day 3, the media is aspirated and 3.0 ml of fresh DMEM and 10% FBS is added to the cells. The cells are allowed to grow to confluency and are split 1:10 on day 6 and day 10, amplifying any replication competent retrovirus. On day 13, the media on the Mv1Lu cells is aspirated and 2.0 ml DMEM and 10% FBS is added to the cells. In addition, the MiCl₁ cells are seeded at 1.0 x 10⁵ cells per well in 2.0 ml DMEM, 10% FBS and 8 µg/ml polybrene. On day 14, the supernatant from the Mv1Lu cells is transferred to the corresponding well of the MiCl₁ cells and incubated overnight at 37°C, 10% CO₂. On day 15, the media is aspirated and 3.0 ml of fresh DMEM and 10% FBS is added to the cells. On day 21, the cells are examined for focus formation (appearing as clustered, refractile cells that overgrow the monolayer and remain attached) on the monolayer of cells. The test article is determined to be contaminated with replication competent retrovirus if foci appear on the MiCl₁ cells. Using these procedures, it can be shown that the HBV core producer cell lines are not contaminated with replication competent retroviruses.

2. Cocultivation of Producer Lines and MdH Marker Rescue Assay

As an alternate method to test for the presence of RCR in a vector-producing cell line, producer cells are cocultivated with an equivalent number of *Mus dunni* (NIH NIAID Bethesda, MD) cells. Small scale cocultivations are performed by mixing of 5.0×10^5 *Mus dunni* cells with 5.0×10^5 producer cells and seeding the mixture into 10 cm plates (10 ml standard culture media/plate, 4 μ g/ml polybrene) at day 0. Every 3-4 days the cultures are split at a 1:10 ratio and 5.0×10^5 *Mus dunni* cells are added to each culture plate to effectively dilute out the producer cell line and provide maximum amplification of RCR. On day 14, culture supernatants are harvested, passed through a 0.45 μ cellulose-acetate filter, and tested in the MdH marker rescue assay. Large scale cocultivations are performed by seeding a mixture of 1.0×10^8 *Mus dunni* cells and 1.0×10^8 producer cells into a total of twenty T-150 flasks (30 ml standard culture media/flask, 4 μ g/ml polybrene). Cultures are split at a ratio of 1:10 on days 3, 6, and 13 and at a ratio of 1:20 on day 9. On day 15, the final supernatants are harvested, filtered and a portion of each is tested in the MdH marker rescue assay.

The MdH marker rescue cell line is cloned from a pool of *Mus dunni* cells transduced with LHL, a retroviral vector encoding the hygromycin B resistance gene (Palmer *et al.*, 1987, *PNAS* 84: 1055-1059). The retroviral vector can be rescued from MdH cells upon infection of the cells with RCR. One ml of test sample is added to a well of a 6-well plate containing 10^5 MdH cells in 2 ml standard culture medium (DMEM with 10% FBS, 1% 200 mM L-glutamine, 1% non-essential amino acids) containing 4 μ g/ml polybrene. Media is replaced after 24 hours with standard culture medium without polybrene. Two days later, the entire volume of MdH culture supernatant is passed through a 0.45 μ cellulose-acetate filter and transferred to a well of a 6-well plate containing 5.0×10^4 *Mus dunni* target cells in 2 ml standard culture medium containing polybrene. After 24 hours, supernatants are replaced with standard culture media containing 250 μ g/ml of hygromycin B and subsequently replaced on days 2 and 5 with media containing 200 μ g/ml of hygromycin B. Colonies resistant to hygromycin B appear and are visualized on day 9 post-selection, by staining with 0.2% Coomassie blue.

F. Transduction of Human Cells with KT-ND5 Vector Construct

On day one, HT1080 cells are set up at 2×10^4 cells per well in six well tissue culture plates containing 2 mls standard growth media (DME + 10% FBS). On day two, ND-5 FVIII retroviral vector particles from a confluent vector producing cell line are harvested as a HX-ND-5 clone. They are filtered through .45 μ m syringe filters prior to testing the supernatants. (Alternatively the filtered media supernatants may be frozen at 80 in aliquots for later use.) Polybrene is added to each well such that the final concentration is 8 μ g per ml. Thirty minutes later, either diluted or undiluted retroviral vector supernatant is added to duplicate wells. Typical volumes and dilutions are 0.5 ml per well and four or more 1:3 serial dilutions in growth media. As a control, two wells are transduced with the same volume of growth media only. On day three, the wells are refed with 2mls of fresh media and the cells allowed to reach confluence, which may typically be about day four or five. On this day, the cells are again refed with one ml per well fresh growth media. Twenty four hours later the media is harvested and filtered as above.

15 G. Expression of Transduced Vector for FVIII

The expression of vector transduced human cells for FVIII is detected by the Coamatic^R assay as described above in Example 2B1. Activity is assayed relative to supernatant from the control wells by counting the cells per well from the two control wells and normalizing FVIII expression data per 1×10^6 cells per 24 hours.

20 H. Administration of Vector Construct

1. Animal Administration Protocol

The intestinal epithelium is an attractive site for gene delivery due to its rapidly proliferating tissue mass and the known location of stem cells in the crypts of Lieberkuhn. The deep location of the stem cells in the crypts and the protective role of the mucus gel layer, makes the retrovirus relatively inaccessible to the tissue cells. However, the accessibility of the retroviral vector to these stem cells can be improved in animal models

by the *in vivo* mucus removal method of Sandberg *et al.*, 1994, *Human Gene Therapy* 5:3232-329.

Male Sprague-Dawley rats obtained from Charles River Breeding Laboratories (Portage, MD.) are anesthetized and the cecum is identified upon opening the peritoneal cavity. A 3 cm ileal segment is isolated from the last Peyer's patch in the terminal ileum and ligated at each end. A plastic catheter attached to a syringe is inserted into the segment and two milliliters of the mucolytic agents dithiothreitol and N-acetyl-cysteine is instilled under mild pressure for two minutes, then removed. This procedure is repeated once again before filling the segment with 0.2 to 2.0 ml of retroviral vector particles at 10^6 to 10^{10} cfu/ml. The ligatures are removed 1 to 4 hours later and the abdominal cavity is sutured. Control animals are instilled with formulation buffer only.

Blood is collected from the tail vein and assayed for factor VIII production by a sandwich ELISA specific for human factor VIII (according to the modified procedure of Zatloukal, K., *et al.*, *PNAS* 91:5148-5152, 1994). The ELISA is based on two Diagnostica) ESH 4 (25 μ g/ml in 1.0 M NaHCO_3 /0.5 M NaCl, pH 9.0) is coupled to the ELISA plates overnight at 4°C, washed with 0.1% Tween 20 in PBS, and blocked with 1% BSA in PBS. The samples are applied in 0.05 M Tris-HCl/1 M NaCl/2% BSA, pH 7.5, over 4 hr at room temperature, the plates are washed, and ESH 8 (2.5 μ g/ml in 0.05 M Tris-HCl/1 M NaCl/2% BSA, pH 7.5,) which has been biotinylated with *N*-hydroxysuccinimidobiotin (Pierce, Rockford, IL.) is added for 2 hr at room temperature. The color reaction is performed with peroxidase-conjugated streptavidin (Boehringer Mannheim, Indianapolis, IN.) and o-phenylenediamine dihydrochloride as substrate. The human factor VIII:c standard (from the National Institute for Biological Standards and Control, Hertfordshire, U.K.) and normal rat plasma are used as references.

2. Human Administration Protocol

Lyophilized recombinant retrovirus containing the gene for Factor VIII expression is formulated into an enteric coated tablet or gel capsule according to known methods in the art. These are described in the following patents: US 4,853,230, EP 225,189, AU 9,224,296, AU 9,230,801, and WO 92144,52.

The capsule is administered orally to be targeted to the jejunum. At 1 to 4 days following oral administration of the recombinant retrovirus, expression of Factor VIII is measured in the plasma and blood by the Coamatic[®] Factor VIII assay as described in Example 2B1.

EXAMPLE 3

INTRAVESICAL ADMINISTRATION OF RECOMBINANT RETROVIRUSES EXPRESSING TK

A. Construction of TK Vector Constructs

1. Construction of plasmids containing vector LTR sequences

All of the following retroviral vectors are based on the N2 vector (Keller et al., *Nature* 318:149-154, 1985). Briefly, 5' and 3' Eco RI LTR fragments (2.8 and 1.0 Kb, respectively) (Armentano, *J. Vir.* 61:1647, 1987; Eglitis, *Science* 230:1395, 1985) are initially subcloned into the Eco RI site of plasmids SK⁺ (Stratagene, San Diego, CA) and pUC31. pUC31 is a modification of pUC19 (Stratagene, San Diego, CA) carrying additional restriction sites (Xho I, Bgl II, BssH II, and Nco I) between the Eco RI and Sac I sites of the polylinker. Plasmid N2R3+/- is thereby created from ligation of the SK⁺ plasmid with the 1.0 Kb 3' LTR fragment. The plasmids p31N2R5+/- and p31N2R3+/- are constructed from the ligation of pUC31 with the 2.8 Kb 5' LTR and packaging signal (Y) or the 1.0 Kb 3' LTR fragment, respectively. In each case N2 refers to the vector source, R refers to the fact that the fragment is an Eco RI fragment, 5 and 3 refer to 5' or 3' LTRs, and + or - refers to the orientation of the insert (see Figures 1-6 for examples of LTR subclones).

In one case, a 1.2 Kb Cla I/Eco RI 5' LTR and W fragment from N2 is subcloned into the same sites of an SK⁺ vector. This vector is designated pN2CR5. In another case, the 5' LTR containing a 6 bp deletion of the splice donor sequence (Yee et al., Cold Spring Harbor, Quantitative Biology, 51:1021, 1986) is subcloned as a 1.8 Kb Eco RI fragment into pUC31. This vector is designated p31N25D[+], Figure 6.

2. Construction of plasmids containing HSVTK

The coding region and transcriptional termination signals of HSV-1 thymidine kinase gene (HSVTK) are isolated as a 1.8 Kb Bgl II/Pvu II fragment from plasmid 322TK

(3.5 kb Bam HI fragment of HSV-1 (McKnight *et al.*) cloned into Bam HI of pBR322 (ATCC No. 31344)) and cloned into Bgl II/Sma I-digested pUC31. This construct is designated pUCTK. For constructs which require deletion of the terminator signals, pUCTK is digested with Sma I and Bam HI and the 0.3 Kb fragment containing the (A)_n signal is removed. The remaining coding sequences and sticky-end Bam HI overhang are reconstituted with a double-stranded oligonucleotide made from the following oligomers:

5' GAG AGA TGG GGG AGG CTA ACT GAG 3' (SEQUENCE ID. NO. 1)

5' GAT CCT CAG TTA GCC TCC CCC ATC TCT C 3' (SEQUENCE ID. NO. 2)

The resulting construct is designated pTKD A, Figure 7.

10 For diagnostic purposes, the oligonucleotides are designed to destroy the Sma I site while maintaining the Ava I site without changing the translated protein.

The plasmid pPrTKDA (Figure 8), which contains the HSVTK promoter and coding sequence (lacking an (A)_n signal), is constructed as follows.

1. pTKD A is linearized with Bgl II treated with alkaline phosphatase, and gel
15 purified.
2. A 0.8 Kb fragment containing the HSVTK transcriptional promoter is isolated as a Bam HI/Bgl II fragment from p322TK.
3. Products from (1) and (2) are ligated, transformed into bacteria, and positive clones are screened for the proper orientation of the promoter region. A resultant clone is
20 designated pPrTKDA (Figure 8).

3. Construction of retroviral provectors expressing HSVTK from a constitutive promoter

The retroviral provectors pTK-1 and pTK-3 are constructed essentially as described
25 below.

1. The 5 Kb Xho I/Hind III 5' LTR and plasmid sequences are isolated from p31N2R5(+) (Figure 1).
2. HSVTK coding sequences lacking transcriptional termination sequences are isolated as a 1.2 Kb Xho I/Bam HI fragment from pTKDA (Figure 2).

3. 3' LTR sequences are isolated as a 1.0 Kb Bam HI/Hind III fragment from pN2R3(-) (Figure 2).
4. The fragments from steps 1-3 are mixed, ligated, transformed into bacteria, and individual clones identified by restriction enzyme analysis. The construct is designated TK-1 (Figure 9).
5. pTK-3 is constructed by linearizing TK-1 with Bam HI, filling in the 5' overhang and blunt-end ligating a 5'-filled Cla I/Cla I fragment containing the bacterial lac UV5 promoter, SV40 early promoter, plus Tn5 neo^r gene obtained from pAFVXM retroviral vector (Krieger *et al.*, 1984, *Cell* 39:483; St. Louis *et al.*, 1988, *PNAS* 85:3150).
- 10 Kanamycin-resistant clones are isolated and individual clones are screened for the proper orientation by restriction enzyme analysis (Figure 9).

These constructs were used to generate infectious recombinant vector particles in conjunction with a packaging cell line, such as DA as described above.

15 B. Determination of the Effect of Ganciclovir on Mouse Colon Carcinoma Cells With or Without TK-3 Vector

An experiment was performed to determine whether or not ganciclovir had an effect on CT26 cells (colon tumor 26, Brattain, Baylor College of Medicine, Houston, TX) that were transduced with DA/TK-3. CT26 cells are transduced with G-pseudotyped TK-3 vector. Twenty-four hours after adding the viral supernatant, the CT26 cells are placed under G-418 selection (450 lg/ml). After 10 days incubation, a G-418 selected pool is obtained and designated CT26TK Neo. CT26 TK Neo cells were seeded into six 10 cm² plates at a density of 2.5 X 10⁶ per plate. As controls, each of two other cell types, CT26 and CT26 beta-gal (this cell line was transduced with a retroviral vector carrying the reporter gene β -galactosidase from *E. coli.*), were also seeded into six 10 cm² plates as controls. Five plates of each cell type were treated twice per day for four consecutive days with medium containing ganciclovir concentrations of 100 μ g/ml, 50 μ g/ml, 25 μ g/ml, 12.5 μ g/ml and 6.25 μ g/ml. One plate of each cell type was left untreated. Afterwards, the cells were removed from each dish using trypsin-EDTA, resuspended in DMEM with 10% FBS and counted. The data in Figure 10 shows that even the lowest dose of ganciclovir had a

dramatic cytotoxic effect on the CT26 TKneo cells. This dose of ganciclovir (6.25 $\mu\text{g/ml}$) or even the next higher dose (12.5 $\mu\text{g/ml}$) did not have an effect on either the CT26 or CT26 beta-gal cells. However, beginning at a ganciclovir dose of 25 $\mu\text{g/ml}$, a dose-dependent decrease in cell growth could be seen, although CT26 TK Neo cells were always more sensitive to the drug.

C. Determination of a Ganciclovir Dose For the Treatment of Mice Injected with CT26 TK Neo Cells

In order to test whether *in vivo* transduction of a murine tumor could be used to treat the disease, an experiment was performed to determine the optimal concentration of ganciclovir necessary to eliminate a tumor that was transduced and selected *in vitro* to assure 100% transduction. Twelve groups of 3 mice each are injected with 2×10^5 CT26TK Neo cells. Six groups of mice are injected with these cells intraperitoneally (I.P.) and six groups of mice are injected subcutaneously (S.C.). Two other groups of 3 mice each are injected with 2×10^5 unmodified CT26 cells (as a control) either I.P. or S.C..

Ten days after the injection of the CT26 or CT26TK Neo cells into these groups of mice, several concentrations of ganciclovir treatment are initiated. Each dose regimen consists of 2 daily AM and PM I.P. injections of ganciclovir. The experiment is summarized in Table A below.

TABLE A

Group	Innoculum	Injection Route	Concentration of Ganciclovir (Mg/Kg)
1	CT26 I.P.	0	
2	CT26 TKneo	I.P.	0
3	CT26 TKneo	I.P.	15.63
4	CT26 TKneo	I.P.	31.25
5	CT26 TKneo	I.P.	62.5
6	CT26 TKneo	I.P.	125.0
7	CT26 TKneo	I.P.	250.0
8	CT26 TKneo	I.P.	500.0
9	CT26 Subq.	0	
10	CT26 TKneo	Subq.	0
11	CT26 TKneo	Subq.	15.63
12	CT26 TKneo	Subq.	31.25
13	CT26 TKneo	Subq.	62.5
14	CT26 TKneo	Subq.	125.0
15	CT26 TKneo	Subq.	250.0
16	CT26 TKneo	Subq.	500.0

After 5 days, all of the mice in the 125 mg/Kg, 250 mg/Kg and 500 mg/Kg treated groups were dead due to the toxic effects of ganciclovir. Mice in the 15.63 mg/Kg, 31.25 mg/Kg and 62.5 mg/Kg treated groups were treated for an additional 7 days and were able to tolerate the treatment. Tumor measurements were made for 23 days (Figure 11). CT26TK Neo grew only slightly slower than unmodified CT26 in the absence of ganciclovir. Complete tumor regression was seen in the groups of mice treated with the 62.5 mg/Kg regimen. Partial tumor regression was seen in the 31.25 mg/Kg treated groups. Little or no effect was seen in the 15.63 mg/Kg treated groups as compared to the untreated control groups. Even though there was some toxicity observed in the 62.5 mg/Kg

groups, it was not life threatening and reversible upon the discontinuation of the treatments so this concentration was used for future studies (Figure 11). After 24 days, the I.P. injected animals were sacrificed and evaluated. The optimal concentration for anti-tumor effect was similar whether the tumor was grown I.P. or S.C.

5 D. Comparison of Cytotoxicity on CT26 and CT26TK Neo In Vivo Tumor Growth

In order to determine whether ganciclovir has an effect on the growth of unmodified CT26 tumor cells *in vivo*, 2 groups of 7 mice are injected S.C. with 2×10^5 unmodified CT26 cells and 2 groups of 7 mice are injected S.C. with 2×10^5 CT26TK Neo cells. Seven days after tumor implantation, one group of CT26 injected mice and one group of
10 CT26TK Neo injected mice are placed on a twice daily (AM and PM) regimen of I.P. ganciclovir at 62.5 mg/Kg. These mice are treated for 12 days or until the CT26TK Neo injected animals have no detectable tumor burden. Tumor growth is monitored over a three week period. Mice injected with CT26 and treated with ganciclovir had tumors that were somewhat smaller than untreated mice injected with CT26, indicating a small HSVTK-
15 independent inhibition of tumor growth (Figure 12). However, a dramatic decrease in tumor burden was observed if, and only if, CT26 TKneo containing mice were treated with ganciclovir (Figure 12).

E. Determination of the Effect of Ganciclovir on AY-27 Rat Carcinoma Cells With or Without the TK-3 Vector

20 An experiment is performed to determine whether or not ganciclovir has an effect on AY-27 cells that are transduced with DA/TK-3. AY-27 cells are rat carcinoma cells which have been induced by N-[4-(5-nitro-2-furyl)-2-thiazolyl] formamide (Selman *et al.*, 1986, *J. Urol* 136:141, 1986). The AY-27 cells are transduced with G-pseudotyped TK-3 vector. Twenty-four hours after adding the viral supernatant, the AY-27 cells are placed under G-
25 418 selection (450 μ g/ml). After 10 days incubation, a G-418 selected pool is obtained and designated AY-27TK Neo. AY-27TK Neo cells are seeded into six 10 cm² plates at a density of 2.5×10^6 per plate. As controls, each of two other cell types, AY-27 and AY-27beta-gal (this cell line is transduced with a virus carrying the reporter gene β -

galactosidase from *E. coli.*), are also seeded into six 10 cm² plates as controls. Five plates of each cell type are treated twice per day for four consecutive days with medium containing ganciclovir concentrations of 100 µg/ml, 50 µg/ml, 25 µg/ml, 12.5 µg/ml and 6.25 µg/ml. One plate of each cell type is left untreated. Afterwards, the cells are removed from each dish using trypsin-EDTA, resuspended in DMEM with 10% FBS and counted. The data generated can be used to determine the concentration which has the most cytotoxic effect on the AY-27, AY-27 beta-gal, or AY-27 TK Neo cells.

F. Determination of a Ganciclovir Dose For the Treatment of Rats Injected with AY-27 TK Neo Cells

10 In order to test whether *in vivo* transduction of a murine tumor could be used to treat the disease, an experiment was performed to determine the optimal concentration of ganciclovir necessary to eliminate a tumor that was transduced and selected *in vitro* to assure 100% transduction. Twelve groups of 3 rats each are injected with 2 X 10⁵ AY-27TK Neo cells. Six groups of rats are injected with these cells intravesically. Six other groups of 3 rats each are injected intravesically with 2 X 10⁵ unmodified AY-27 cells (as a control).

15 Ten days after the injection of the AY-27 or AY-27TK Neo cells into these groups of rats, several concentrations of ganciclovir treatment are initiated. Each dose regimen consists of 2 daily AM and PM I.P. injections of ganciclovir. The experiment is summarized in Table B below.

20

TABLE B

Group	Innoculum	Injection Route	Concentration of Ganciclovir (Mg/Kg)
1	AY-27		0
2	AY-27 TKneo	intravesically	0
3	AY-27 TKneo	intravesically	15.63
4	AY-27 TKneo	intravesically	31.25
5	AY-27 TKneo	intravesically	62.5
6	AY-27 TKneo	intravesically	125.0
7	AY-27 TKneo	intravesically	250.0
8	AY-27 TKneo	intravesically	500.0

Rats are treated for 12 days, eliminating those that die due to the higher concentrations of ganciclovir. Tumor measurements are made for 20 days, assessing tumor regression in order to determine an optimal ganciclovir concentration.

G. Administration Protocol

5 1. Rat Administration Protocol

AY-27 rat bladder carcinoma cells are transplanted into the bladders of 20 male Fischer 344 rats (Charles River Breeding Laboratories, Portage, MD). One to fourteen days following transplantation, tumor-bearing rats weighing between 200 to 300 grams are anesthetized, their bladders are surgically exteriorized and evacuated of urine with a 27-
10 gauge needle. Viral vector particles are instilled into the bladders of 5 rats at 10^6 to 10^{10} in cfu 200 to 2,000 μ l of formulation buffer. The addition of 4 to 8 μ g/ml of polybrene increases the efficiency of transduction. In order to prevent leakage, the cystotomy is repaired with 7-zero nonabsorbable suture. The virus is allowed to incubate in the presence of the tumor cells for 0.5 to 1.0 hour by keeping the animal anesthetized and thereby
15 preventing voiding. Ten control rats receive 500 μ l of formulation buffer only.

Alternatively, 200 to 2,000 μ l of vector can be instilled directly into the bladder by catheterization through the urethra following urine evacuation and rinsing with saline.

At 24 to 72 hours after vector treatment, the rats are placed on twice daily (AM and PM) injections of I.P. ganciclovir at the previously determined optimum dose (e.g. 62.5 mg/Kg body weight) for 4 to 12 days. Finally, the rats receive a single daily dose of ganciclovir until the end of the experiment (1 to 10 weeks). Whole bladders are removed and tumor growth is measured.

2. Human Administration Protocol

A urinary (Foley) catheter is inserted through the urethra into the bladder and secured in place. The bladder is evacuated of urine and washed with 100 to 500 mls of sterile saline. Recombinant retroviruses containing the gene for thymidine kinase expression are instilled through the catheter into the bladder at 10^5 to 10^{11} cfu in 10 to 500 ml of formulation buffer preferably containing 4 to 8 μ g/ml of polybrene, or other enhancing excipients. The viral particles are allowed to incubate for 0.25 to 12 hours prior to removal of the catheter. After 1 to 7 days, ganciclovir is administered at 1 to 5mg/Kg I.V. (at a constant rate over 1 hour) every 12 hours for 2 to 21 days. The vector can be readministered multiple times (2 to 20), followed by ganciclovir administration. Due to the frequency of granulocytopenia and thrombocytopenia in patients receiving ganciclovir, it is recommended that neutrophil and platelet counts be performed every two days during the dosing of the drug. Tumor regression is monitored by x-ray and/or biopsy and the treatment repeated if required.

EXAMPLE 4

Pulmonary Administration of Recombinant Retroviruses Expressing Factor VIII

A. Construction of Full-Length and B Domain Deleted Factor VIII cDNA Retrovector™

The construction of the full-length and B domain deleted Factor VIII retrovectors are described in Example 2A.

B. Aerosolization of Recombinant Retroviruses Expressing Factor VIII

The KT-ND5 viral supernatant in formulation buffer, with 4 to 8 µg/ml of polybrene or other transduction enhancing excipient, is nebulized using a DeVilbiss #15 Atomizer (DeVilbiss Health Care Division, Somerset, PA) designed to produce 0.3 to 0.5 µm particles (Rousculp, 1992, *Human Gene Therapy* 3:471-477). The aerosol produced by this nebulizer uses compressed air in a laminar flow hood. The mist is directed into a polypropylene tube, and the condensed vapor, as well as the control viral supernatant, is resterilized by 0.22 µm filtration. Retroviral particles pass through this filter without any significant loss of functional activity.

C. Administration of Vector Construct

1. Rat Administration Protocol

Rats are anesthetized and the trachea is exposed by anterior midline incision. Recombinant retroviral particles expressing the factor VIII gene product are diluted in 300 µl of formulation buffer, with or without 4 to 8 µg/ml of polybrene or other transduction enhancing excipient, at 10^5 to 10^{10} cfu/ml and instilled into the trachea through a small gauge needle. Control animals are administered 300µl of formulation buffer only. The incision is sutured and the rats are allowed to recover. Two to fourteen days following viral instillation, blood is drawn from the tail vein and examined for factor VIII production as described in Example 2Hi.

2. Human Administration Protocol

The recombinant retrovirus is administered using a DeVilbiss #15 Atomizer (as described above) for 2 to 60 minutes. Two to fourteen days following administration of the recombinant retroviruses, expression of Factor VIII is measured in the blood and plasma by the Coamatic^R Factor VIII assay (as described in Example 2B1).

EXAMPLE 5

Transdermal Administration of Recombinant Retroviruses

A. Construction of TK-3 Vector Construct

Construction and verification of the TK-3 vector construct and recombinant
5 retroviruses are described in Examples 3A through 3F.

B. Administration of Vector Construct

1. Animal Administration Protocol

Cottontail rabbit papillomavirus (CRPV) provides an animal model for the highly
10 oncogenic human papillomavirus (HPV). Papillomas can be induced in the cottontail rabbit
and virus infection leads to three different outcomes in the rabbit (Lin, Y. et al., *J Virol*
67:382, 1993). First, papillomas appear and persist for the lifetime of the rabbit; second,
papillomas spontaneously regress 2 to 3 months after infection; and third, papillomas
progress to carcinomas after 8 to 15 months.

15 In this experiment, twelve cottontail rabbits (E. Johnson, Rago, KA) are
injected with the B strain of CRPV (Stevens, J. et al., *J Virol* 30:891, 1979) by intradermal
injection as described by Stevens, J., et al., (*J Virol* 30:891, 1979). Four to six months after
infection, when papillomas form, the animals are divided into two groups. In the first
group, papillomas of 6 animals are injected with 25 to 100 μ l of formulation buffer, with or
20 without 4 to 8 μ g/ml of polybrene or other transduction enhancing excipient, at 10^5 to 10^{10}
cfu/ml through a small gauge needle. In the second group, control animals are administered
25 to 100 μ l of formulation buffer only. At 24 to 72 hours after vector treatment, the rabbits
are placed on twice daily (AM and PM) injections of I.P. ganciclovir at the previously
determined optimum dose (e.g. 62.5 mg/Kg body weight) for 4 to 12 days. Finally, the
25 rabbits receive a single daily dose of ganciclovir until the end of the experiment (1 to 10
weeks). Papilloma regression is visually monitored for 2 to 14 days.

2. Human Administration Protocol

The clinical cutaneous lesions that result from the human papillomavirus (HPV) include common warts, filiform warts, plantar warts, and anogenital warts (reviewed in Cobb *et al.*, 1990, *J. Am. Acad. Derm.* 22:547). In this experiment, patients are divided into two groups. In the first group, 100 to 500 μ l of recombinant retrovirus particles at a concentration of 10^9 cfu/ml in a formulated ointment, preferably containing 4 to 8 μ g/ml of polybrene or other enhancing excipients, are applied to the warts using a transdermal delivery system (TDS).

Transdermal delivery systems (TDS) are capable of delivering a drug through intact skin so that it reaches the systemic circulation in sufficient quantity to be therapeutically beneficial. TDS provides a variety of advantages including elimination of gastrointestinal absorption problems and hepatic first pass effect, reduction of dosage and dose intervals, and improved patient compliance. The major components of TDS are a controlled release device composed of polymers, the drug to be administered, excipients, and enhancers, and a fastening system to fix the device to the skin. A number of polymers have been described which include gelatin, gum arabic, paraffin waxes, and cellulose acetate phthalate (Sogibayasi, K., *et al.*, *J. Controlled Release*, 29:177-185, 1994).

The second group receives 100 to 500 μ l of vector particle in formulation buffer only. The areas are covered and the viral particles are allowed to incubate for 0.25 to 12 hours prior to removal of the TDS. After 1 to 7 days, ganciclovir is administered at 1 to 5mg/Kg I.V., preferably at a constant rate over 1 hour) every 12 hours for 2 to 21 days. The vector can be readministered multiple times (2 to 20), followed by ganciclovir administration. Due to the frequency of granulocytopenia and thrombocytopenia in patients receiving ganciclovir, it is recommended that neutrophil and platelet counts be performed every two days during the dosing of the drug. The regression of the wart is visually monitored for 2 to 14 days.

EXAMPLE 6

Ocular Administration of Recombinant Retroviruses for E3/19K

A. Cloning of E3/19K Gene into KT-3B

5 1. Isolation and Purification of Adenovirus

The isolation and purification of adenovirus is described by Green *et al.* (*Methods in Enzymology* 58: 425, 1979). Specifically, five liters of Hela cells ($3-6.0 \times 10^5$ cells/ml) are infected with 100-500 plaque forming units (pfu) per ml of adenovirus type 2 (Ad2) virions (ATCC No. VR-846). After incubation at 37°C for 30-40 hours, the cells are placed on ice,
10 harvested by centrifugation at 230 xg for 20 minutes at 4°C, and resuspended in Tris-HCl buffer (pH 8.1). The pellets are mechanically disrupted by sonication and homogenized in trichlorotrifluoroethane prior to centrifugation at 1,000 xg for 10 min. The upper aqueous layer is removed and layered over 10 mls of CsCl (1.43 g/cm^3) and centrifuged in a SW27 rotor for 1 hour at 20,000 rpm. The opalescent viral band is removed and adjusted to 1.34 g/cm^3
15 g/cm^3 with CsCl and further centrifuged in a Ti 50 rotor for 16-20 hours at 30,000 rpm. The visible viral band in the middle of the gradient is removed and stored at 4°C until purification of adenoviral DNA.

2. Isolation and Purification of Adenovirus DNA

20 The adenovirus band is incubated with protease for 1 hour at 37°C to digest proteins. After centrifugation at 7,800 xg for 10 minutes at 4°C, the particles are solubilized in 5% SDS at room temperature for 30 minutes before being extracted with equal volumes of phenol. The upper aqueous phase is removed, re-extracted with phenol, extracted three times with ether, and dialyzed in Tris buffer for 24 hours. The viral Ad2 DNA is
25 precipitated in ethanol, washed in ethanol, and resuspended in Tris-EDTA buffer (pH 8.1). Approximately 0.5 mg of viral Ad2 DNA is isolated from virus produced in 1.0 L of cells.

3. Isolation of E3/19K Gene

The viral Ad2 DNA is digested with EcoR I and separated by electrophoresis on a 1% agarose gel. The resulting 2.7 Kb Ad2 EcoR I D fragments, located in the Ad2 coordinate region 75.9 to 83.4, containing the E3/19K gene (Herisse *et al.*, *Nucleic Acids Research* 8:2173, 1980, Cladaras *et al.*, *Virology* 140:28, 1985) are eluted by electrophoresis, phenol extracted, ethanol precipitated, and dissolved in Tris-EDTA (pH 8.1).

4. Cloning of E3/19K Gene into KT-3B

The E3/19K gene is cloned into the EcoR I site of PUC1813. PUC1813 is prepared as essentially described by Kay *et al.* (*Nucleic Acids Research* 15:2778, 1987) and Gray *et al.* (*PNAS* 80:5842, 1983). The E3/19K is retrieved by EcoR I digestion and the isolated fragment is cloned into the EcoR I site of phosphatase-treated pSP73 plasmid. This construct is designated SP-E3/19K. The orientation of the SP-E3/19K cDNA is verified by using appropriate restriction enzyme digestion and DNA sequencing. In the sense orientation, the 5' end of the cDNA is adjacent to the Xho I site of the pSP73 polylinker and the 3' end adjacent to the Cla I site. The Xho I-Cla I fragment containing the E3/19K cDNA in either sense or antisense orientation is retrieved from the SP-E3/19K construct and cloned into the Xho I-Cla I site of the KT-3BB retroviral. This construct is designated KT-3B/E3/19K.

B. Cloning of PCR Amplified E3/19K Gene into KT-3B

1. PCR Amplification of E3/19K Gene

The Ad2 DNA E3/19K gene, including the amino terminal signal sequence, followed by the intraluminal domain and carboxy terminal cytoplasmic tail which allow the E3/19K protein to embed itself in the endoplasmic reticulum (ER), is located between viral nucleotides 28,812 and 29,288. Isolation of the Ad2 E3/19K gene from the viral genomic DNA is accomplished by PCR amplification, with the primer pair shown below:

The forward primer corresponds to the Ad2 nucleotide sequences 28,812 to 28,835.
(Seq ID No. 3) 5'-TATATCTCCAGATGAGGTACATGATTTTAGGCTTG-3'

The reverse primer corresponds to the Ad2 nucleotide sequences 29,241 to 29,213.
5 (Seq ID No. 4) 5'-TATATATCGATTCAAGGCATTTTCTTTTCATCAATAAAAC-3'

In addition to the Ad2 complementary sequences, both primers contain a five nucleotide "buffer sequence" at their 5' ends for efficient enzyme digestion of the PCT amplicon products. This sequence in the forward primer is followed by the Xho I
10 recognition site and by the Cla I recognition site in the reverse primer. Thus, in the 5' to 3' direction, the E3/19K gene is flanked by Xho I and Cla I recognition sites. Amplification of the E3/19K gene from Ad2 DNA is accomplished with the following PCR cycle protocol:

Temperature°C	Time (min)	No. Cycles
94	2.0	1
94	0.5	
55	0.17	5
72	3.5	
94	0.5	30
70	3.5	
72	10.0	10

2. Ligation of PCR Amplified E3/19K Gene into KT-3B

15 The E3/19K gene from the SP-E3/19K construct, approximately 780 bp in length, is removed and isolated by 1% agarose/TBE gel electrophoresis. The Xho I-Cla I E3/19K fragment is then ligated into the KT-3B retroviral backbone. This construct is designated KT-3B/E3/19K. It is amplified by transforming *E. coli*, DH5 alpha bacterial strain (Bethesda Research Labs, Gaithersburg, Maryland) with the KT-3B/E3/19K construct.
20 Specifically, the bacteria is transformed with 1-100 ng of ligation reaction mixture DNA.

The transformed bacterial cells are plated on LB plates containing ampicillin. The plates are incubated overnight at 37°C, bacterial colonies are selected and DNA prepared from them. The DNA is digested with Xho I and Cla I. The expected endonuclease restriction cleavage fragment sizes for plasmids containing the E3/19K gene are 780 and 1,300 bp.

5

C. Transduction of Packaging Cell Line DA with the Recombinant Retroviral Vector KT-3B/E3/19K

1. Plasmid DNA Transfection

293 2-3 cells (a cell line derived from 293 cells ATCC No. CRL 1573, (WO 92/05266) 5.0×10^5 cells are seeded at approximately 50% confluence on a 6 cm tissue culture dish. The following day, the media is replaced with 4 ml fresh media 4 hours prior to transfection. A standard calcium phosphate-DNA coprecipitation is performed by mixing 10.0 µg of KT-3B/E3/19K plasmid and 10.0 µg MLP G plasmid with a 2M CaCl_2 solution, adding a 1x HEPES buffered saline solution, pH 6.9, and incubating for 15 minutes at room temperature. The calcium phosphate-DNA coprecipitate is transferred to the 293 2-3 cells, which are then incubated overnight at 37°C, 5% CO_2 . The following morning, the cells are rinsed three times in 1x PBS, pH 7.0. Fresh media is added to the cells, followed by overnight incubation at 37°C, 10% CO_2 . The following day, the media is collected off the cells and passed through a 0.45 µ filter. This supernatant is used to transduce packaging and tumor cell lines. Transient vector supernatant for other vectors are generated in a similar fashion.

15
20

2. Packaging Cell Line Transduction

Packaging cell line transduction is performed as described in Example 2C.

25

3. Detection Of Replication Competent Retroviruses

Detection of replication competent retroviruses is performed as described in

Example 2D

D. Transduction of Cell Lines with the Recombinant Retroviral Vector KT-3B/E3/19K

The following adherent human and murine cell lines are seeded at 5×10^5 cells/10 cm dish with 4 μ g/ml polybrene: HT 1080, Hela, and BC10ME. The following day, 1.0 ml of filtered supernatant from the DA E3/19K pool is added to each of the cell culture plates.

- 5 The following day, 800 μ g/ml G418 is added to the media of all cell cultures. The cultures are maintained until selection is complete and sufficient cell numbers are generated to test for gene expression. The transduced cell lines are designated HT 1080-E3/19K, Hela-E3/19K and BC10ME-E3/19K, respectively.

- EBV transformed cell lines (BLCL), and other suspension cell lines, are
10 transduced by co-cultivation with irradiated producer cell line, such as DA-E3/19K. Specifically, irradiated (10,000 rads) producer line cells are plated at 5.0×10^5 cells/6 cm dish in growth media containing 4 μ g/ml polybrene. After the cells have been allowed to attach for 2-24 hours, 10^6 suspension cells are added. After 2-3 days, the suspension cells are removed, pelleted by centrifugation, resuspended in growth media containing 1mg/ml
15 G418, and seeded in 10 wells of a round bottom 96 well plate. The cultures were expanded to 24 well plates, then to T-25 flasks.

E. Expression of E3/19K in the Recombinant Retroviral Vector Construct KT3B-E3/19K

20 1. Western Blot Analysis

- Radio-immuno precipitation assay (RIPA) lysates are made from selected cultures for analysis of E3/19K expression. RIPA lysates are prepared from confluent plates of cells. Specifically, the media is first aspirated off the cells. Depending upon the size of the culture plate containing the cells, a volume of 100 to 500 ml ice cold RIPA lysis buffer (10 mM
25 Tris, pH 7.4; 1% Nonidet P40; 0.1% SDS; 150 mM NaCl) is added to the cells. Cells are removed from plates using a micropipet and the mixture is transferred to a microfuge tube. The tube is centrifuged for 5 minutes to precipitate cellular debris and the supernatant is transferred to another tube. The supernatants are electrophoresed on a 10% SDS-polyacrylamide gel and the protein bands are transferred to an Immobilon membrane in
30 CAPS buffer (10 mM CAPS (Aldrich, Milwaukee, WI) pH 11.0; 10% methanol) at 10 to 60

volts for 2 to 18 hours. The membrane is transferred from the CAPS buffer to 5% Blotto (5% nonfat dry milk; 50 mM Tris, pH 7.4; 150 mM NaCl; 0.02% Na azide, and 0.05% Tween 20) and probed with a mouse monoclonal antibody to E3/19K (Severinsson *et al.*, 1985, *J. Cell. Biol.* 101:540-547). Antibody binding to the membrane is detected by the use of ¹²⁵I-Protein A.

2. FACS Analysis of KT3B-E3/19k-Vector Transduced Cells to Demonstrate Decreased Levels of Class I Expression Compared to Non-Transduced Cells.

Cell lines transduced with the KT3b-E3/19K-vector are examined for MHC class I molecule expression by FACS analysis. Non-transduced cells are also analyzed for MHC class I molecule expression and compared with E3/19K transduced cells to determine the effect of transduction on MHC class I molecule expression.

Murine cell lines, BC10ME, BC10ME-E3/19K, P815 (ATCC No. TIB 64), and P815-E3/19K, are tested for expression of the H-2D^d molecule on the cell surface. Cells grown to subconfluent density are removed from culture dishes by treatment with Versene and washed two times with cold (4°C) PBS plus 1% BSA and 0.02% Na-azide (wash buffer) by centrifugation at 200g. Two million cells are placed in microfuge tubes and pelleted in a microfuge at 200g before removing the supernatant. Cell pellets are resuspended with the H-2D^d-specific Mab 34-2-12s (50 ml of a 1:100 dilution of purified antibody, ATCC No. HB 87) and incubated for 30 min at 4°C with occasional mixing. Antibody labeled cells are washed two times with 1 ml of wash buffer (4°C) prior to removing the supernatant. Cells are resuspended with a biotinylated goat anti-mouse kappa light chain Mab (50 ml, of a 1:100 dilution of purified antibody) (Amersham, Arlington Height, IL) and incubated for 30 min at 4°C. Cells are washed, resuspended with 50ml of avidin conjugated FITC (Pierce, Rockford, IL), and incubated for 30 min at 4°C. The cells are washed once more, resuspended in 1 ml of wash buffer, and held on ice prior to analysis on a FACStar Analyzer (Becton Dickinson, Los Angeles, CA). The mean fluorescence intensity of transduced cells is compared with that of non-transduced cells to determine the effect E3/19K protein has on surface MHC class I molecule expression.

F. Administration of Vector Construct

1. Rat Administration Protocol

Rats are anesthetized and one eye is instilled with 5 to 100 μ l of recombinant retroviral particles at a concentration of 10^5 to 10^{10} cfu/ml in formulation buffer, with or without 4 to 8 μ g/ml of polybrene or other transduction enhancing excipient,. Five to one hundred μ l of solution containing formulation buffer only is added to the other eye to be used as a control. The solution is allowed to incubate for 1 hour before each eye is rinsed 3 times with 100 μ l of saline. Two to seven days following the treatment, the rat is sacrificed, the cornea is removed, and homogenized in 2 ml ice cold RIPA lysis buffer. Expression of E3/19K is detected by Western blot analysis as described in Example 6E1.

2. Human Administration Protocol

Although the rate of corneal transplant rejection is relatively low, the current therapy for those with rejection requires continuous treatment of steroid compounds. This eventually leads to cataract formation, requiring surgery. Therefore, the introduction of a recombinant retrovirus expressing the E3/19K would prevent the need for such a steroid regimen. Ten to five hundred μ l of recombinant retroviruses at a concentration of 10^5 to 10^{10} cfu/ml in formulation buffer, with or without 4 to 8 μ g/ml of polybrene or other transduction enhancing excipient in formulation buffer, are administered to the eye of a patient lying in a prone position. The solution is allowed to incubate for 15 to 30 minutes before being washed with saline.

Alternatively, the cornea may be incubated for 1 hour in 1 ml of retroviral vector particles at a concentration of 10^5 to 10^{10} cfu/ml in formulation buffer, with or without 4 to 8 μ g/ml of polybrene or other transduction enhancing excipient, just prior to surgical attachment. In either of the above cases, the progress of the transplant is monitored by visually observing tissue viability.

EXAMPLE 7

Intranasal Administration of Recombinant Retroviruses Expressing Factor VIII

A. Construction of Full-Length and B Domain Deleted Factor VIII cDNA
Retrovector™

The construction of the full-length and B domain deleted Factor VIII retrovectors are described in Example 2A

5 B. Administration of Vector Construct

1. Rat Administration Protocol

The nasal route has been shown to be effective for the administration of a number of molecules due to the extensive network of capillaries located under the nasal mucosa. This facilitates effective systemic absorption and when the drug is administered with absorption
10 promoters, absorption occurs rapidly with high bioavailability (review in Gizurarson *et al.*, 1990, *Acta Pharm* 2:105).

One group of six Fischer-344 rats are used for nasal administration of the retroviral vector particles for Factor VIII. One to fifty μ l of retroviral vector particles at 10^9 cfu/ml in formulation buffer, with or without 4 to 8 μ g/ml of polybrene or other transduction
15 enhancing excipients are applied with a pipette inserted about 3 to 5 mm into each nostril. Another group is administered formulation buffer without vector in the same manner. Blood samples are collected from the jugular or tail vein 1 to 14 days later and assayed for factor VIII production as described in Example 2Hi.

2. Human Administration Protocol

20 Several types of drug delivery devices for the nasal cavity exists (reviewed in Chien *et al.*, 1987, *Crit Rev Therap Drug Carr Sys* 4:67). These systems include nasal spray, nose drops, saturated cotton pledget, aerosol spray, and insufflator. The metered-dose nebulizer can deliver a predetermined volume of the formulation to the nasal cavity.

Two groups of patients are used in this study. One group of patients receives 100 to
25 500 μ l of retroviral vector particles at 10^9 cfu/ml in formulation buffer, with or without 4 to 8 μ g/ml of polybrene or other transduction enhancing excipients, applied to each nostril via nasal spray or nasal drops. Another group receives formulation buffer only applied in the

same manner. Blood samples are collected 1 to 14 days later and assayed for factor VIII production as described in Example 2B1.

EXAMPLE 8

5 PREPARATION OF RECOMBINANT RETROVIRUS FOR DELIVERY OF HUMAN GROWTH HORMONE

A. Preparation of hGH containing vectors

Vector pDHF828 containing the full-length human growth hormone gene is constructed essentially as follows. Briefly, plasmid pDHF811, was constructed by
10 removing the XhoI- ClaI fragment of the KT-1 retroviral vector described above, and inserting the following oligonucleotide linkers by ligation of the cohesive ends:

Linker sequences:

(SEQUENCE ID# 5) 5' TCGAGGATCC GCCCGGGCGG CCGCATCGAT GTCGACG
15 3'
(SEQUENCE ID# 6) 5' CGCGTCGA CATCGATGCG GCCGCCCGGG CGGATCC 3'

In particular, the linkers were annealed at 65°C for 20 minutes, 42°C for 20 minutes, 37°C for 20 minutes, and room temperature for 2 hours. The concentrations of both
20 oligonucleotides was 18mM and the salt concentration was 100 mM NaCl. After annealing, 50ml of 1.8 mM annealed linker was digested with ClaI overnight to generate ClaI ends. For ligation, 3nM of KT-1 XhoI - ClaI fragment was mixed with 90nM of linker, and the resultant mixture incubated at 15°C for 3 hours. The ligated DNA sample was transformed into DH-5α competent cells, followed by screening of transformants.

25 Plasmid chGH 800 containing the full length cDNA of the hGH gene (Martial *et al.*, 1979, *Science* 205:602) was digested with Hind III, blunt-ended with the Klenow fragment enzyme, and cloned into the SrfI site of pDHF811. The resultant plasmid was designated pDHF828.

B. Preparation of hGH expressing recombinant retrovirus

The pDHF828 plasmid was then introduced into the HX packaging cell, using standard procedures and assayed using the HGH Chemiluminescence Kit (HGH 100T) (Nichols Institute, San Juan Capistrano, CA.), according to a preferred modification of the kit protocol. On day 1, the kit components were warmed to room temperature and gently mixed by inversion before opening any vials. Test samples were centrifuged for 5' at top speed in a microfuge before using them in order to remove fibrin and other debris. All samples were measured in quadruplicate, including the standards. The incubations are performed in 12 x17 polypropylene tubes that have been stored in the dark. One hundred
10 fifty ul of sample or standard were aliquoted into each tube and ul of antibody is added and the samples were mixed gently. One bead was added to each well using the forceps provided in the kit. The tubes were capped, covered with foil, and shaken on an orbital shaker for 24 hr at room temperature. Standards contain 530 pg/ml (STD D), and serial dilutions were made in zero standard of Std D of 250, 100, 50, 25, 10, 5, and 2.5 pg/ml.

15 After 24 hours, the tubes were uncapped and 0.5 ml of wash buffer were added. These wash solution was added with enough force to make the bead bounce up off the bottom of the tube. The samples were washed three times with 2.0 ml nanopure water, and aspirated completely each time. The luminometer determinations were done in 12x75 polycarbonate (clear plastic) tubes stored in the dark. The luminometer was pretested with
20 performance control standards.

Using this assay, HX/HGH retrovector producing cell lines were generated with titers of 4.8×10^6 cfu/ml. Introduction of the plasmid into DX packaging cells resulted in production of clonal producer cells with a titer of 1.6×10^7 cfu/ml.

EXAMPLE 9

25 Preservation of a Recombinant Retrovirus

A. Lactose Formulation of a Recombinant Retrovirus

Crude recombinant retrovirus is obtained from a Celligan bioreactor (New Brunswick, New Brunswick, NJ) containing DA cells transformed with the recombinant

retrovirus bound to the beads of the bioreactor matrix. The cells release the recombinant retrovirus into the growth media that is passed over the cells in a continuous flow process. The media exiting the bioreactor is collected and passed initially through a 0.8 micron filter then through a 0.65 micron filter to clarify the crude recombinant retrovirus. The filtrate is concentrated utilizing a cross flow concentrating system (Filtron, Boston, MA).

Approximately 50 Units of DNase (Intergen, New York, NY) per ml of concentrate is added to digest exogenous DNA. The digest is diafiltrated using the same cross flow system to 150 mM NaCl, 25 mM tromethamine, pH 7.2. The diafiltrate is loaded onto a Sephadex S-500 gel column (Pharmacia, Piscataway, NJ), equilibrated in 50 mM NaCl, 25 mM tromethamine, pH 7.4. The purified recombinant retrovirus is eluted from the Sephadex S-500 gel column in 50 mM NaCl, 25 mM tromethamine, pH 7.4.

The formulation buffer containing lactose was prepared at a 2X concentrated stock solution. The formulation buffer contains 25 mM tromethamine, 70 mM NaCl, 2 mg/ml arginine, 10 mg/ml HSA, and 100 mg/ml lactose in a final volume of 100 mls at a pH 7.4.

The purified recombinant retrovirus is formulated by adding one part 2X lactose formulation buffer to one part S-500 purified recombinant retrovirus. The formulated recombinant retrovirus can be stored at -70°C to -80°C or dried.

The formulated retrovirus is lyophilized in an Edwards Refrigerated Chamber (3 Shelf RC3S unit) attached to a Supermodulyo 12K freeze dryer (Edwards High Vacuum, Tonawanda, NY). When the freeze drying cycle is completed, the vials are stoppered under a vacuum following a slight nitrogen gas bleeding. Upon removal, vials are crimped with aluminum seals.

In the given lactose study, formulated liquid product was stored at both -80°C and at -20°C cycling freezer. In Figure 13 viral infectivity of these samples were compared to the viral infectivity of lyophilized samples. The lyophilized samples were stored at -20°C, refrigerator temperature and room temperature. Activity of the samples upon reconstitution are determined by titer assay.

The lyophilized recombinant retrovirus is reconstituted with 1.0 ml water. The infectivity of the reconstituted recombinant retrovirus is determined by a titer activity assay. The assay is conducted on HT 1080 fibroblasts or 3T3 mouse fibroblast cell line (ATCC

No. CCL 163). Specifically, 1×10^5 cells are plated onto 6 cm plates and incubated overnight at 37°C, 10% CO₂. Ten microliters of a dilution series of reconstituted recombinant retroviruses are added to the cells in the presence of 4 mg/mL polybrene (Sigma, St. Louis, MO) and incubated overnight at 37°C, 10% CO₂. Following incubation, cells are selected for neomycin resistance in G418 containing media and incubated for 5 days at 37°C, 10% CO₂. Following initial selection, the cells are re-fed with fresh media containing G418 and incubated for 5-6 days. After final selection, the cells are stained with Commassie blue for colony detection. The titer of the sample is determined from the number of colonies, the dilution, and the volume used.

Figure 13 demonstrates that storage in lyophilized form at -20°C to refrigerator temperatures retains similar viral activity as a recombinant retrovirus stored in liquid at -80 to -20°C permitting less stringent temperature control during storage.

B. Mannitol Formulation of a Recombinant Retrovirus

The recombinant retrovirus utilized in this example was purified as described in Example 9A.

The formulation buffer containing mannitol was prepared as a 2X concentrated stock solution. The formulation buffer contains 25 mM tromethamine, 35 mM NaCl, 2 mg/ml arginine, 10 mg/ml HSA and 80 mg/ml mannitol at a final volume of 100 mls at a pH 7.4.

The purified recombinant retrovirus is formulated by adding one part mannitol formulation buffer to one part S-500 purified recombinant retrovirus. The formulated recombinant retrovirus can be stored at this stage at -70°C to -80°C or dried.

The formulated retrovirus is dried in an Edwards Refrigerated Chamber (3 Shelf RC3S unit) attached to a Supermodulyo 12K freeze dryer. When the freeze drying cycle is completed, the vials are stoppered under a vacuum following nitrogen gas bleeding to 700 mbar. Upon removal, vials are crimped with aluminum seals.

In the given mannitol study, formulated liquid product was stored at both -80°C and at -20°C in cycling freezers. The viral infectivity of these samples were compared to the viral infectivity of lyophilized samples, Figure 14. The lyophilized samples were stored at -

20°C, refrigerator temperature and room temperature. Activity of the samples upon reconstitution are determined using the titer assay described in Example 9A.

Figure 14 demonstrates that storage in lyophilized form at -20°C to refrigerator temperature retains significant viral activity as compared to recombinant retrovirus stored in liquid at -80°C or -20°C, permitting less stringent temperature control during storage.

C. Trehalose Formulation of a Recombinant Retrovirus

The recombinant retrovirus utilized in this example was purified as described in Example 9A.

10 The formulation buffer containing trehalose was prepared as a 2X concentrated stock solution. The formulation buffer contains 25 mM tromethamine, 70 mM NaCl, 2.0 mg/ml arginine, 10.0 mg/ml HSA and 100 mg/ml trehalose at a final volume of 100 mls at a pH 7.2.

15 The purified recombinant retrovirus is formulated by adding one part trehalose formulation buffer to one part S-500 purified recombinant retrovirus. The formulated recombinant retrovirus can be stored at this stage at -70°C to -80°C or dried.

The formulated retrovirus is dried in an Edwards Refrigerated Chamber (3 Shelf RC3S unit) attached to a Supermodulyo 12K freeze dryer. When the freeze drying cycle is completed, the vials are stoppered under a vacuum following nitrogen gas bleeding to 20 700 mbar. Upon removal, vials are crimped with aluminum seals.

In the given trehalose study, formulated liquid product was stored at both -80°C and at -20°C in cycling freezers. The viral infectivity of these samples was compared to the viral infectivity of lyophilized samples, Figure 15. The lyophilized samples were stored at -20°C, refrigerator temperature and room temperature. Activity of the samples upon 25 reconstitution are determined using the titer assay as described in Example 9A.

Figure 15 demonstrates that storage in lyophilized form at -20°C to refrigerator temperature retains similar viral activity as compared to recombinant retrovirus stored in liquid at -80°C to -20°C permitting less stringent temperature control during storage.

Viral infectivity of liquid formulated recombinant retrovirus samples stored at -80°C 30 was compared to viral infectivity of lyophilized formulated recombinant retrovirus stored at

-20°C. Initially, a bulk of recombinant retrovirus was received and formulated in four different ways as shown below. The formulated recombinant retrovirus was then frozen in bulk for 1.5 months subsequent to being quick thawed and freeze dried. Positive controls were stored at -80°C for comparison with lyophilized samples which were stored at -20°C after freeze-drying. The formulations are listed below:

Formulation	Buffer				Human Serum
	Sugar	Concentration	Salt	Arginine	Albumin
	Concentration (mg/ml)	(mM tromethamine)	Concentration (mM NaCl)	Concentration (mg/ml)	Concentration (mg/ml)
Mannitol	40	25	25	1	5
Lactose	40	25	75	1	5
Sucrose	50	25	60	1	5
Trehalose	50	25	60	1	5

In the graphs of Figure 16, the y-axis on each of the 4 graphs (A, B, C, D) represent the normalized titer. At an initial time point after lyophilization, $t = 0$, a titer value was established for both the -80°C liquid sample and the -20°C lyophilized sample. At each time point of the stability study, the titer obtained was divided by the zero time point titer value and the % of original entered onto the graph.

The data demonstrates that post-lyophilization activity is maintained in the lyophilized sample (stored at -20°C) relative to the liquid sample (stored at -80°C). The formulated lyophilized recombinant retrovirus was stored in a -20°C freezer (a frost-free cycling freezer). Comparison to the formulated liquid recombinant retrovirus stored at -80°C indicates the lyophilized form permits less stringent control of storage conditions.

EXAMPLE 10

Analysis of Crude and Purified Recombinant Retrovirus

- Crude and purified solutions of recombinant retrovirus particles may be separated on gradient polyacrylamide gels utilizing, for example, the PHASTGEL system (Pharmacia Biotech). Briefly, samples are placed on 4-15% polyacrylamide gels without pretreatment and electrophoresed for 35 minutes at 250V. The gels are then removed and stained with coomassie blue in order to detect virus and other protein components. The gels are then scanned by laser densitometry in order to determine the content of virus and other components.
- 10 Virus bands may be identified by their relative molecular weight and by reverse transcriptase activity (RT). The purpose of this assay is to quantify the activity of reverse transcriptase (RT), an enzyme exclusively associated with all retroviruses. The relative amount of retrovirus in a sample can be determined by measuring the activity of this enzyme in a given preparation.
- 15 Briefly, moloney murine leukemia virus reverse transcriptase (Pharmacia, Newark, NJ) is diluted to a concentration of 1 µg/ml by addition of 1x Tris/EDTA buffer solution containing 10 mM Tris-HCl and 1mM EDTA, pH 8.0. One hundred microliters of this solution is added 6.84 ml of sterile dH₂O, 500 µl of 1M Tris HCl pH 8.0, 10 µl of 0.1M MnCl₂, 200 µl of 1M dithiothreitol, 50 µl of 10% Nonidet P40 (NP40), 2 µl of 100 µM dNTP (Pharmacia, Newark, NJ, dNTP Ultrapure KitTM), and 300 µl Methyl - ³H
- 20 Thymidine 5' - Triphosphate (30-50 Ci/mmol). This mixture is incubated for 1 hour at 37°C in a water bath. Following incubation the sample is placed on ice. Approximately 1.0 ml of 2N HCl is added to the cooled sample. The precipitated radiolabeled DNA fragments are vacuum filtered onto glass fiber filters using a Millipore sampling manifold (Millipore, Philadelphia, PA). The filters are washed, dried, placed in scintillation cocktail, and
- 25 counted in a Beckman LS5000TD scintillation counter (Beckman, Dallas, TX).

EXAMPLE 11

Analysis of Complement Resistance in Various Packaging Cell Lines

A. Packaging Cell Line Preparation and Transduction

Four different packaging cell lines were used to package pCB β -gal into infectious
5 virions. Two cell lines were derived from D17 dog cells ("D") (ATCC CCL 183), and two
were derived from the human embryonic kidney cell line 293 ("2") (ATCC CRL 1573). In
the case of both dog and human packaging cells, one cell line expressed the amphotropic
envelope from the 4070A virus (*i.e.*, DA and 2A, respectively) and the other expressed the
xenotropic envelope from the NZB9-1 virus (*i.e.*, DX and 2X, respectively). All packaging
10 cells were grown in DMEM media (Irvine Scientific, Santa Ana, CA) supplemented with
10% heat inactivated fetal bovine serum (Hyclone, Salt Lake City, UT).

Each of the four packaging cell lines was transduced with a G-pseudotyped pCB β -
gal (*see* Burns et al., 1993, *Proc. Nat'l. Acad. Sci. USA* 90:8033-8037, for a pseudotyping
procedure). After G418 selection, the selected cell pools were dilution cloned (except for
15 the amphotropic vector producing human cell line which was maintained as a non-clonal
pool). Vector producing cell lines producing the highest titer on HT1080 cells (ATCC No.
CCL 121) were selected. Titers were determined approximately 2 days after transduction
by G418 colony forming units or by X-gal staining of the monolayers. (*See Current
Protocols in Molecular Biology*, Ed. Ausubel et al., for more details on the titering assays.)

20 Retrovirus containing supernatants were collected from each of the confluent
monolayers of the individual amphotropic envelope (DA and 2A) or xenotropic envelope
(DX or 2X) vector producing cell lines and filtered through a 0.45 μ m filter prior to
aliquoting and storage at -70°C.

B. Serum Inactivation Assay

25 Serum inactivation titer assays were performed as follows: Serum was drawn from at
least two different human volunteers, chimpanzees, baboons, and macaque monkeys.
Approximately 20-70 mL of blood was collected from each donor into serum separating
tubes (Becton Dickinson, Los Angeles, CA). Blood was allowed to clot for 20-30 minutes
at room temperature, after which time the samples were centrifuged at 2000 x g for 10

minutes at 4°C. Serum was frozen in approximately 1.1 mL aliquots and stored at -80°C. Vials from each batch were tested for total classical complement activity (Quantiplate, Kallestad Labs, Inc., Chaska, MN), and only batches with "normal" levels of complement activity were used. 1 mL aliquots of fresh, 100% serum were used for each inactivation
5 assay. Complement inactivated controls of each serum were prepared by heat inactivation for 30 minutes at 56°C. Undiluted and diluted preparations of supernatants containing recombinant retroviral particles (BCFU greater than or equal to 10^3 per mL sera) were mixed 1:10 with medium control, sera, or heat inactivated sera. These mixtures were then incubated at 37°C for 30 minutes. Treated vector particles were then titered by a standard
10 blue colony forming unit assay (*Current Protocols in Molecular Biology, supra*).

The results of this experiment are presented below in Table 1 as BCFU/mL.

Table 1**Packaging Cell Line**

Serum source	DA	DX	2A	2X
HUMAN				
control	19,000	110,000	5,700	6,700
100% sera	400	0	6,100	7,600
H.I. sera	18,000	100,000	10,500	8,800
CHIMP				
control	16,200	31,600	27,800	22,400
100% sera	900	1,700	21,500	18,300
H.I. sera	28,300	47,000	42,700	48,500
BABOON				
control	16,000	13,000	8,000	7,200
100% sera	0	0	1,800	1,000
H.I. sera	24,000	9,600	15,000	9,400
MACAQUE				
control	170,000	13,000	2,500	2,300
100% sera	0	0	0	10
H.I. sera	270,000	12,000	3,000	2,200

H.I. = heat inactivated

control = DMEM + 10% H.I. fetal bovine serum + Sodium pyruvate

Note: zero's indicate that no blue colonies were detected in 0.1 or 0.2 ml undiluted volumes, therefore indicating that titers were conservatively less than 20-40 BCFU/ml.

Briefly, these results demonstrate that recombinant retroviruses which are made in human packaging cell lines exhibit no detectable sensitivity to inactivation by a heat labile component of human serum, presumably complement, in *in vitro* assays. In contrast, they show partial sensitivity to inactivation by chimp, and then increasing sensitivity to baboon and macaque serum, in order of increasing phylogenetic distance from man. In addition, these *in vitro* results demonstrate that recombinant retroviruses produced in D17 derived packaging cell lines exhibit near total sensitivity to inactivation by a heat labile component, presumably complement, of human, chimp, baboon and macaque serum.

Other experiments which were conducted employing another recombinant retroviral genome packaged in either of two of the above-described cell lines (dog cell line DA, and human cell line 2A), as well as a different human cell line (HX, a HT1080-derived packaging cell line; confirm that the conclusions drawn from the above results are not vector dependent. Moreover, human producer cell lines derived from 293 and HT1080 generated equally complement resistant vector.

Additional experimental data which was generated using recombinant retroviral particles made in human HT1080-derived packaging cells expressing a different murine envelope tropism, poly, similarly showed insensitivity to heat labile human serum components, further confirming the conclusions above.

EXAMPLE 12

Recombinant Retrovirus Production from Hollow Fiber Cultures

A. Culture Initiation

To initiate a hollow fiber culture, first condition the hollow fiber bioreactor (HFB) for 48 hours prior to seeding by simulating a run condition with 100-200 mL of complete growth media at 37°C. The growth media should be what ever the cell line has been adapted to. All liquids in the HFB when originally shipped should be aspirated and replaced with the complete growth media. When seeding the bioreactor, the cells should not have been split more than 48 hours earlier and should be in log growth phase at the time of

harvest for the seeding of the HFB. The cells are harvested by trypsinization and pelleted by centrifugation. The cell pellet is then resuspended in 4 mL of 25% pre-conditioned media and delivered to the extra-capillary space by syringe using the side syringe ports found on the HFB. After seeding the HFB, allow the cells to adhere for 20 to 30 minutes before starting the circulation pump. During this time replace the media used to condition the HFB with 100-200 ml using 25% pre-conditioned media. The circulation feed pump is initiated with the starting flow rate set at 25 mL/min. (setting 5 with 2 long pump pins). After 1 hour from the time of switching the pump on, a one mL sample of media is collected in order to record the initial levels of lactate and ammonia. On a daily schedule, 1 ml samples are collected every 24 hours to assay for the daily production of lactate and ammonia. The initial 100-200 ml of old is exchanged media with fresh media when lactate levels begin to reach 2.0 g/l (or the equivalent to 22 mM/L) The same volume of media is replaced until the culture approaches daily levels of 20 mmol/L. When daily levels of lactate reach 20 mmol/L increase your reservoir bottle size to a 500 mL bottle containing 500 mL of fresh media. The flow feed rate is then increased to 50 mL/min. when the culture begins to produce 2.2 mmol/day of lactate. When daily 500 mL volumes reach 20 mmol/L of lactate, the original Cellco supplied reservoir feeding cap is exchanged for the larger reservoir cap (Unisyn-vender part #240820) adapted for the Cellco system with the addition of tubing and male luer lock fittings. This reservoir cap will accommodate the large 2 Liter Corning bottles. (To avoid the exchange of the reservoir caps during a culture run, initiate the culture run with the larger reservoir cap which can also support smaller bottle sizes.) When daily lactate readings are assayed and recorded, one can calculate the daily levels of lactate production of the culture in order to determine when the culture reaches maximum cell density when the rate of lactate decreases and levels off.

25 B. Seeding density for the 2X- β -gal

In order to establish specific seeding requirements, two hollow fiber runs are established. One run is seeded with a low number of cells and one with a high number of cells. Progress of the culture is tracked by analyzing the daily glucose consumption and

lactate production levels. Figure 20 is a representative graph of data generated over a two week period of the vector producer cell line 2X- β -GAL₁₇₋₁₄.

In this experiment, one HFB was seeded with 1.3×10^7 cells (to represent the low seed culture) the other seeded with 1.6×10^8 cells. In this experiment, the cell line (2x- β -GAL₁₇₋₁₄) was able to initiate a good hollow fiber run under low and high seeding conditions. Being able to incubate the HFB with fewer cells, is only convenient for reducing the effort required for generating the number of cells required to start a culture. However a low seed start also extends the time it takes to reach optimal cell densities which usually yield the highest titers. In this experiment, the cell line used, adapted very well to hollow fiber cultures which eventually required daily media changes of 500 ml per day.

C. Cell Culture Health and Maximum Cell Densities:

In the original Cellco design, it was observed that the original media reservoir cap was not suited to fit larger bottles other than the standard 100 and 500 ml media bottles. This is a problem when aggressive growth cultures require greater than 500 ml daily exchanges of media. Daily multiple changes of media increases the likelihood of culture contamination by increasing the daily handling time of the system. If one does not opt to perform multiple daily exchanges of media then one exposes the culture to daily toxic levels of waste products which can affect the cell expansion of the system along with the length at which the culture run will survive. Figure 20-B demonstrates lactate concentrations of a culture which required daily 500 mL exchanges of fresh media after reaching day 7 of culture. Figure 20-D is one indication of the health of the culture by tracking the amount of lactate being produced on a daily basis. The graphs indicate that the culture was no longer allowed to expand based on the plateauing of daily production of lactate. Another indicator of the health of the cell culture was the drop in peak titer production which also correlated with the daily exposure of high levels of lactate (See Figure 21). These findings would indicate that optimal titers can be correlated with the maximum cell densities and the relative health of the culture.

D. Optimal Titer Concentrations, Frequency of Harvests and Total Harvest Amounts

β -gal titers for the above experiment were determined from frozen samples and were
titered on 293 cells assayed 48 or 72 hours after transduction. The transduced cells were
stained for β GAL activity and individual cells counted on a hemocytometer giving a titer
5 based on the number of blue cells /mL (BCT/mL). As shown in Figure 21, optimum titers
were obtained on day 7 of the high seed culture at 1.8×10^8 BCT/mL from a 72 hour blue
cell titer on 293 cells. The duplicate culture initially seeded with a 10 fold lower seeding
density, peaked at 5.2×10^7 BCT/mL from a 48 hour blue cell titer. Previous flat stock
cultures of 2X b-GAL₁₇₋₁₄ cultures have been titered using 48 hour blue cell titers on
10 HT1080 cells and have been calculated to be 5×10^6 BCT/mL. If one uses the values
obtained from 48 hour blue cell titers, the increase in titer by using hollow fiber systems is
ten fold higher than crude supernatants obtained from tissue culture dishes or flasks. These
maximum titers were reached prior to hitting the daily 20 mmol/L toxic levels lactate which
appeared to reduce the titer produced the following week. Crude supernatants can be
15 harvested every 9 hours with out any loss of titer (See Figure 2). It is predicted that 3
harvests per day can be achieved with minimum loss of titers. In addition, continuous
hollow fiber cultures can be maintained for several weeks. When titers were compared
between the low and the high seed culture, there was little differences by day 11 between
the two seed cultures averaging 4×10^7 BCT/mL.

20

EXAMPLE 13

Two-Phase Purification of Recombinant Retroviruses

A. Concentration of DA/ND-7 recombinant particles

Five milliliters of formulated DX/ND-7 recombinant retroviral particles at a titer of
 3.5×10^8 cfu/ml (total of 1.75×10^9 cfu) is diluted in 1400 ml of media (DMEM containing
25 5% Fetal Bovine Serum). Three hundred milliliters of two-phase partitioning components
(PEG-8000 (autoclaved), dextran-sulfate, and NaCl) are added to a final concentration of
6.5% PEG, 0.4% dextran-sulphate, and 0.3 M NaCl. The resultant solution is placed into a

two-liter separatory funnel, and left in a cold room for 24 hours (including two mixing steps approximately 6 to 16 hours apart).

Following the 24 hour period, the bottom layer (approximately 20 mL) is carefully eluted, and the interphase (approximately 1 mL) is collected in a 15 mL conical FALCON tube. The interphase containing vector is diluted to 10 mL by addition of PBS, and incubated at 37°C in order to bring the solution to room temperature and destabilize the micelles.

To one-half of the diluted interphase, KCl is added to a final concentration 0.4 M, and mixed well. The tube is then placed on ice for ten minutes, and spun for 2 minutes at 2,000 rpm in a bench-top centrifuge. The supernatant is removed and filtered through a 0.45 um syringe filter.

The other half of the interphase containing vector is separated by S-500 Sephadex chromatography in 1X PBS.

The results of these concentration processes, as determined in a BCFU assay, are shown below in Table 1:

TABLE 1

<u>PHASE</u>	<u>CONCENTRATION</u>
Crude	1.1×10^9 bcfu
Separation: Top phase	1.4×10^8 bcfu
Separation: Interphase	$7(+/-3) \times 10^8$ bcfu
Separation: Bottom phase	2×10^6 bcfu
Final step: KCl separation	$*6(+/-3) \times 10^8$ bcfu
Final step: S-500 separation	$*1.8(+/-0.3) \times 10^8$ bcfu

* Note that since the sample was split into two halves, that these numbers were doubled in order to represent the level of purification that would be expected if the entire 1 mL interphase was separated as indicated.

In summary, 1.4 liters of crude research grade supernatant containing recombinant retroviral particles may be reduced to a 10 ml volume, with approximately 50% (+/-20%) being recovered when KCl separation is utilized as the final step. When S-500 chromatography is utilized as the final step, only about 10% of the initial recombinant retroviral particles are recovered in a 14 ml.

In order to complete concentration of the retroviral vector particles, the vector-containing solution may be further subjected to concentration utilizing an MY-membrane Amicon filter, thereby reducing the volume from 10 to 14 mL, down to less than 1 mL.

10

EXAMPLE 14

Production of Vector From DX/ND7 β gal Clone 87 Utilizing a Cell Factory

DX/ND7 β gal clone 87, an expression vector, was grown in cell factories. Cells were grown in DMEM supplemented with Fetal Bovine Serum in roller bottles until enough cells to seed 20 10-layer cell factories (NUNC) at a 1:3 dilution were obtained. Each 10-layer cell factory is seeded with approximately 0.8 liters of cell medium.

Cells were seeded into the cell factory by pouring media containing cells into the factory so that the suspensions evenly fill the 10 layers. The factory is then carefully tilted away from the port side to prevent the suspension from redistribution in the common tube. Finally, the cell factory is rotated into its final upright position. A hepaent filter is attached to each port. The factory was then placed in a CO₂ incubator.

In three days, and for each of the next three days, supernatant containing vector was harvested. The cell factory is placed in a tissue culture hood. One filter is removed and sterile transfer tubing is connected to the open port. The factory is lifted so that supernatant drains into the tubing. Approximately 2 liters of supernatant is harvested from each factory. Fresh DMEM/FBS is used to replenish the lost medium. The transfer tubing is removed and the factory replaced in the incubator. From 20 cell factories, approximately 90 liters of crude vector containing supernatant were obtained.

Verification of the vector was performed by transduction of HT1080 cells. These cells were harvested 2 days later and stained for β -gal protein. The titer of the supernatant was determined to be 2×10^7 /ml.

EXAMPLE 15

Concentration of Recombinant Retroviruses by Low-Speed Centrifugation

A. Retrovector Supernatant Preparation

5 Producer cell lines DA/βgal and HX/DN-7 were cultured in a culture flask and a roller bottle, respectively, containing Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% fetal bovine serum plus 1 mM L-Glutamine, Sodium pyruvate, non-essential amino acids and antibiotics. Viral supernatant was harvested from the flask and roller bottle, and were filtered through a 0.45 μm syringe filter. The filtered supernatants were stored either at 4°C (HX/ND7), or frozen at -70°C (DAβ-gal).

10 B. Virus Concentration

Viral supernatant was aliquoted into 50 ml sterile OAKRIDGE screw cap tubes, and placed into an SS34 rotor for use in a Sorvall centrifuge. The tubes were spun for 1 hour at 16,000 rpm (25,000g-force) at 4°C. Upon completion of the spin, the tubes were removed, the supernatant decanted and a small opaque pellet resuspended in the DMEM media
15 described above.

C. Virus Titration

Concentrated virus was titered on HT1080 cells plated 24 hours earlier at a cell density of 2×10^5 cells per well in a six well plate + 4 μg/ml polybrene. Briefly, virus preps
20 were diluted from 1/10 to 1/10,000 and 50 μl of each dilution was used to infect one well from the six well plate. Plates were incubated overnight at 37°C. Forty-eight hours later, cells were fixed and stained with X-gal. The results are set forth below in Table 1.

Table 1. Virus Concentration through Low Speed Centrifugation

Parameter description	Experiment number				
	1	2		3	
Virus source	DA β -gal	DA β -gal	HX/ND7	DA β -gal	HX/ND7
Titer of normal harvest	4.4×10^6	2.1×10^6	3.2×10^5	5×10^6	5×10^5
Titer of virus concentrate	6×10^8	7.4×10^7	3.2×10^7	2.9×10^8	3.9×10^7
Starting volume	80 ml	.39 ml	39 ml	118ml	40ml
Final concentrate volume	.5 ml	.36 ml	.36 ml	.78ml	.28ml
Fold virus concentration	136X	34X	100X	58X	78X
Virus recovery	87%	30%	91%	50%	99%

As is evident from Table 1, virus recovery ranged from 30% to 99%, with the best recovery
 5 being obtained from human producer cells (HX/ND7; recovery ranged from 91% to 99%).

EXAMPLE 16

Concentration of Recombinant Retroviruses by Ultrafiltration

S-500 purified supernatant containing the β -gal expressing recombinant retrovirus
 DX/CB- β gal and partially concentrated supernatant containing the same virus were each
 10 filtered through a 0.45 μ m filter, and loaded into a CENTRIPREP-100 filter (product #4308,
 Amicon, MA). The supernatants were kept at a temperature of 4 °C throughout this
 procedure, including during centrifugation. The CENTRIPREP filters were spun three
 times each for 45 to 60 minutes at 500 x G. Between each spin the filtrate was decanted.
 The retentate was thus sequentially reduced, such that the initial 15 ml (or 10 ml) volume
 15 was reduced to approximately 0.6 mL per unit.

The resultant titer was determined by assaying HT1080 target cells set up at a
 concentration of 1×10^5 cells per well 24 hours prior to transduction of the viral sample.
 Cells were transduced in the presence of 8 μ g/ml polybrene and 2 mL growth media

(DMEM plus 10% FBS) per well. As shown in Table 1 below, approximately one hundred percent of the virus was recovered utilizing this procedure (note that titers are in BCFU/ml).

MTable 1

	<i>Pre-centrifuge titer/volume.</i>	<i>Final titer/volume</i>
S-500	$4 \times 10^7/15 \text{ ml}$	$1.3 \times 10^7/0.6 \text{ ml}$
<i>part. conc.</i>	$3 \times 10^7/10 \text{ ml}$	$1 \times 10^{10}/0.6 \text{ ml}$

5

EXAMPLE 17Preparation of Recombinant Retrovirus in a BioreactorA. Freezing protocol

Producer cells are frozen in DMEM media containing 10% to 20% FBS, and 5 to 10 15% DMSO, at a concentration of 1×10^7 cells/ml/vial. Cells are frozen in a controlled rate freezer (Series PC, Controlled Rate Freezing System, Custom Biogenic Systems, Warren MI) at a rate of from 1 to 10 °C per minute. Frozen cells are stored in liquid nitrogen.

B. Bioreactor protocol

Cells are thawed from frozen vials at 37°C, washed once with media to remove 15 DMSO, and expanded into 850 cm² "FALCON" roller bottles (Corning, Corning, N.Y.) Expanded cell culture is used to inoculate a "CELLIGEN PLUS" bioreactor (5 liter working volume; New Brunswick, Edison, N.J.). The cells are grown on microcarriers (*i.e.*, Cytodex 1 or Cytodex 2; Pharmacia, Piscataway, N.J.) at a concentration of 3 to 15 g/L microcarrier. Initial inoculation densities are from 4 to 9 cells/bead at half to full volume for 2 to 24 20 hours. The media constituents for virus production are DMEM-high glucose (Irvine Scientific, Santa Ana, CA.) basal media supplemented with FBS (10 to 20%), Glutamine (8 to 15mM), glucose (4.5 to 6.5 g/L), Nonessential amino acids (1X), RPMI 1640 amino acids (0.2 to 9.6X), 10 mM HEPES, RPMI 1640 Vitamins (0.2 to 5X).

During culture, pH (6.9 to 7.6) and dissolved oxygen ("DO" 5 to 90%) are controlled 25 by the use of a four gas system which includes air, oxygen, nitrogen, and carbon dioxide.

After several days of batch growth the culture is then continuously perfused with fresh media with concurrent continuous harvesting in an escalating perfusion rate of 0.5 to 2.5 volumes/day. Cell retention is the result of differential sedimentation of cell covered beads in a decanting column.

During operation the bioreactor is monitored for viable cells, titer, glucose, lactate, ammonia levels, and lack of contamination. Viable cells and titer range from 1×10^5 cells/ml to 1×10^7 cells/ml. Glucose ranges from 6 to 0.25 g/L, Lactate from 1 to 25 mM, and Ammonia ranges from 0.5 to 30 mM. Cells are incubated in the bioreactor for 5 to 25 days.

EXAMPLE 18

Long Term Expression of Human Factor VIII in Rabbits Following

Intravenous Expression of Vector

Recombinant retroviral vector expressing B domain deleted factor VIII was constructed, packaged, and expressed as described in Example 2 herein. Large scale production was carried out in Cell Factories as described for DX/ND7 in Example 14. The retroviral vector was then lactose formulated as described in Example 9 herein. High titer formulated retroviral vector was injected into 500 g juvenile rabbits according to the following schedule: 1.5E8 vector particles were injected into the ear vein at 2.5 hr intervals, three times per day for three days (Day 0, 1, and 2). On days 3, 4, 5, 6, 7, 56, and every 7 days subsequently, citrated plasma was obtained by venipuncture of the ear vein and analyzed for human factor VIII antigen as follows.

Day 1: Coat Immulon IV plates (Dynatech, Chantilly, VA, USA) . Plate 1 was coated with 200 ul anti-factor VIII light chain monoclonal ESH8 (American Diagnostica, Greenwich, CT, USA) at 3 ug/ml in coating buffer (0.1M bicarbonate, pH 9.2) at 4 degrees overnight. Plate 2 and 3 were coated with 300 ul Sigma fractionated ascites anti-beta galactosidase (Sigma #G4644, Sigma Chemical Company, St. Louis, MO, USA) at 8 ug/ml. These are the "absorption" plates.

Day 2: Plates were washed once with buffer DB2 (1000 ml = 7.8 g Tris HCl/ 58.4 g NaCl/20 g BSA/1 ml Tween 20, pH 7.5)., then blocked with 300 ul DB2, covered with
30 parafilm, and incubated at room temperature for at least 90 min. on an orbital shaker at 80

rpm. Plates were then washed in DB2. 200 ul samples in duplicate were diluted in DB2 (plasma samples were diluted at least 1:3), centrifuged for 5' at room temperature at 14,000 rpm in an Eppendorf microfuge, and added to plate 1. Plate 1 was covered with parafilm and incubated two hours at room temperature shaking at 80 rpm.

5 00 ul of polyclonal anti-FVIII:C at 20 ug/ml in buffer DB2 (Enzyme Research Labs, South Bend, Indiana, USA) was placed on absorption plate 2, covered with parafilm, and incubated at room temperature shaking at 80 rpm. Likewise, 300 ul donkey anti-sheep IgG conjugated to alkaline phosphatase (Sigma Chemical Company, St Louis Mo., USA) was absorbed on absorption plate 3 at 1:30,000 in DB2, covered and incubated at room
10 temperature shaking at 80 rpm.

After two hours at room temperature, plate 1 was washed and 200 ul of absorbed sheep anti-VIII:C was transferred from plate 2 to plate 1. Plate 1 was covered with parafilm and incubated at room temperature for 1-2 hrs, shaking at 80 rpm.

After 2 hr, plate 1 was washed and 200 ul absorbed donkey anti-sheep conjugate was
15 transferred from plate 3 to plate 1. Plate 1 was covered and incubated 1 hr at room temperature shaking at 80 rpm.

Plate 1 was washed, and freshly prepared Attophos substrate (JBL Scientific, Huntingdon, England, UK) prepared as recommended by the manufacturer was added at 200 ul per well. the plate was read at 2 min intervals (30 cycles) on Cytofluor II fluorometer,
20 excitation 450, emission 580 filters; mix=1; gain = 65.

Readings were compared to a standard curve using duplicate dilutions of NHP (pooled normal plasma, George King, Overland Park, Kansas, USA) as a factor VIII source diluted in 1:3 rabbit plasma at 3-fold dilutions starting at 1:10 NHP.

Data from Day 0 (prebleeds) to day 261 from 7 experimental and two control
25 rabbits are shown in Figure 22. Levels remained low during the first 7 days, with no significant differences being seen between the experimental and control groups ($p > 0.25$). On Day 56 and subsequent days, two rabbits displayed 30-50 ng/ml human factor VIII antigen in their plasmas. These levels, with slight fluctuations, were maintained at least until Day 261 (8 months later), the last time point tested. The experimental group as a whole
30 over the entire time course was significantly different from the controls ($p < 0.03$).

Therefore, rabbits injected via ear vein with high titer formulated retroviral vectors displayed high level systemic expression of factor VIII following a lag phase of 7-56 days.

Figure 23 shows the data in this experiment extended through Day 429. The two high-expressing rabbits described above maintain their expression levels through day 429.

- 5 Medium-level expressing rabbits do as well, while the control rabbits remain negative. Thus, expression is observed at least from Day 56-429, indicating >373 days of expression from a single course of vector administration.

Example 19

High Level Expression of Human Growth Hormone With a Lag Phase in Rabbits Following Intravenous Injection of Retroviral Vector

Human growth hormone-expressing retroviral vectors were produced as described in Example 8 herein. Large scale production in Cell Factories was performed as described for DX/ND7 in Example 14 herein. The retroviral vector was lactose formulated as described in Example 9 herein. The titer was $8E8$. Vector was diluted 5-fold in formulation buffer, and 1 ml injections were given to 500 g juvenile rabbits 3 time per day for three days as described in Example 18 for factor VIII vector. Serum was collected on days 0 (prebleed), 7, 14, and every 7 days thereafter. Serum was analyzed for human growth hormone content using a commercial ELISA kit (Boehringer Mannheim Biochemicals, Indianapolis, Indiana, USA) according to the directions of the manufacturer. The results are shown in Figure 24. A modest degree of expression (100 pg/ml) was observed in rabbit #861 at Day 7. Several other rabbits displayed smaller initial expression. Expression in all rabbits declined, to zero except for rabbit #861, by Day 14. Unexpectedly, expression in several rabbits (#861, 863, 858, 866) began to rise between Day 21 and Day 36. Two rabbits (#863 and 861) displayed peak expression levels at Day 36 of 200-350 pg/ml. Expression in all rabbits subsequently declined to zero at Day 56. A periodic small rise and decline was seen thereafter, reaching a low steady state of approximately 50 pg/ml in one rabbit (#863) after Day 84. Therefore, rabbits injected via ear vein with high titer formulated retroviral vector displayed a high level systemic expression of human growth hormone with a lag phase of 20-36 days.

Example 20

High Level Expression of Human Growth Hormone with a Lag Phase in Mice Following Intravenous Injection of Retroviral Vector

Human growth hormone-expressing retroviral vectors were produced as described in Example 8, herein. Large scale production in Cell Factories was performed as described for DX/ND7 in Example 14, herein. The retroviral vector was then lactose formulated as described in Example 9, herein. The titer was $8E8$. Three week old mice were given a single injection of 200 ul vector via the tail vein. Human growth hormone was analyzed in mouse

serum using a commercial ELISA kit (Boehringer Mannheim Biochemicals) according to the instructions of the manufacturer. The results are shown in Figure 25. Due to the low serum volumes that could be obtained from young mice relative to the volumes required for ELISA analysis, pooled test mice were tested on Day 14, and therefore the results shown on this date represent the average value for the indicated animals. All three pools (two test, on control) yielded values of < 50 pg/ml on Day 14. By Day 28, animals had grown sufficiently that individual animals could be tested. Animals 2 and 3 displayed hGH levels of approximately 300 pg/ml on Day 28; animal 9 showed a level; of 200 pg/ml. All other mice were <100 pg/ml. The two controls, Animals #11 and 12, were 0 throughout the test period. By day 48, all 10 test animals displayed increasing levels of hGH, and all were clearly above 100 pg/ml, with animals #1, 2, 3, 4, 5, and 10 showing levels of 300-600 pg/ml. Therefore, mice injected a single time with high-titer formulated retroviral vector expressed human growth hormone at high levels for at least 20 days following a lag phase of 14-48 days.

15

Example 21

High Level Expression of Human Growth Hormone in Adult Mice

Human growth hormone-expressing retroviral vectors were produced as described in Example 8 herein. Large scale production was carried out in Cell Factories as described for DX/ND7 in Example 14 herein. The retroviral vectors were then lactose formulated as described in Example 9, herein. The titer was 8E8. Adult mice were given a single injection of 100 µl vector via the tail vein. One group of mice also received 600 µg cyclosporin A once daily i.m. starting 1 day prior to injection of vector and continuing through Day 13. Human growth hormone was analyzed in mouse serum using a commercial ELISA kit (Boehringer Mannheim Biochemicals) according to the instructions of the manufacturer. The results are shown in Figure 26. "Positive control" refers to mice that received no cyclosporin. High level (400-1200 pg/ml) human growth hormone was observed on Day 3, declining to zero by Day 14. Negative controls (formulation buffer-injected) expressed no human growth hormone. The injection of cyclosporin A using the above schedule neither augmented nor diminished the growth hormone response. Therefore, intravenous injection

of high titer formulated retroviral vectors induces high level expression of growth hormone in adult animals with no lag phase.

Example 22

5 Use of Peroxisome Proliferators to Induce Liver Mitosis and Retroviral Transduction In Vivo

Balb/C mice were treated IP (500 μ l) or via gavage (200 μ l) with the peroxisome proliferator WY 14643 (Chem Syn, Lenexa, Kansas, USA) (15 mg/ml) once per day for 4 days. On day 4, 200 μ l beta galactosidase retrovector (1E9 cfu/ml, DA-Bgal) was injected via tail vein. On day 7, the mice were sacrificed, tissues were fixed in 2% formaldehyde/PBS for 24 hr, and blocked into slices of 2-5 mm. Sections were rinsed in PBS and stained in fresh XGAL solution (5 mM potassium ferricyanide/5 mM potassium ferrocyanide/ 2 mM MgCl₂/ 0.5mg/ml XGAL in DMF (Gold Scientific)/1X PBS). Tissue was embedded in paraffin (-50 degrees), sectioned at 5 microns (PML, San Diego), and counterstained with Hematoxylin/Eosin. Sections were compared from both periportal and peripheral regions from normal liver (XGAL stain only), control liver (no WY14643, with vector), WY control liver (WY14643 only, no vector), liver from group I (IP WY14643 plus vector), liver from group II (gavage WY14643 plus vector), and spleen from Group I. Numerous blue cells were observed in both treatment groups, both periportal and on the liver periphery. Very rare blue cells were observed in the control liver (without mitogen). No blue cells were seen in the normal liver (no mitogen or vector) or the WY liver (mitogen, no vector). The results indicate that the mitogen, administered either IP or by gavage, promoted liver transduction by beta-galactosidase-expressing vector, while no background staining was induced by mitogen in the absence of vector.

Example 23

Biolocalization of Intravenously Administered Retroviral Vector Following Long-Term Expression of Gene of Interest

Rabbits which had received intravenous administration retroviral vector resulting in
5 > 373 days of expression of human factor VIII (See Example 18, herein) were sacrificed on
day 511. Brain, spleen, and liver samples were collected using a fresh set of sterilized
instruments for each organ in each animal. DNA was isolated using lysis buffer (8M urea,
2% SDS, 0.35M NaCl, 10 mM Tris, pH 8.0) followed by phenol/chloroform extraction.
DNA was quantitated by fluorometry and diluted to 500 ng/25 ul.. Integrity of each genomic
10 DNA samples was verified by 0.6% agarose gel electrophoresis. PCR was performed on
each sample (0.5 uM primer LTR 492S (CCC TGT GCC TTA TTT GAA CTA ACC) (Seq
ID No. 12), 0.55 uM primer LTR 1072AS (CCC ACC ACA ACC ACA TAT CCC TCC)
(Seq ID No.13), 1X PCR Buffer, 2.5 mM MgCl₂, 200 uM dNTPs, 2.5 U Taq in a total
volume of 50 ul (Perkin-Elmer)). The reactions are incubated 10 min at 94o, then subjected
15 to 32 cycles of 94o x 30 sec, 68o x 30 sec, 72o x 30 sec. The amplicon consisted of a 580 bp
region spanning the LTR/Psi region of the vector backbone (see Example 1). Each sample
was subjected to four replicate reactions. Two other replicate reactions were spiked with 5
copies of positive control DNA (DA6A3 HBV-IT VCL genomic DNA; see Townsend *et*
al., 1997, *J. Virol.* 71:3365, to control for interference with PCR amplification. Ten
20 microliters of each amplification reaction were dot-blotted onto a nylon membrane (Zeta
Probe). The membrane was prehybridized in buffer (6X SSPE, 30% formamide, 5X
Denhardt's Solution, 0.5% SDS, 1.5 mg/ml herring sperm DNA) at 42o for 60 min, the
hybridized at 42o for 4 hr in the same buffer to a 32P-end labeled probe (LTR-P - CCA
GTC CTC CGA TTG ACT G [Seq ID No. 14]) corresponding to a sequence internal to the
25 LTR-Psi amplicon. The blot was washed (at room temperature in 2x SSPE/1% SDS, then 3x
10 min at 42o in 2x SSPE/1% SDS) and exposed to film. The film was inspected for
number of positive blots per total number of reactions. Tissue samples from Rabbit #82,
which expressed high levels of human factor VIII protein, and from Rabbit #95, a control
animal that had received no vector, are shown in Figure 27. The spiked reactions from all

tissues were positive, indicating that negatives are not due to PCR inhibiting substances from the tested tissues. Liver samples from 10 locations, a spleen sample, and samples from two brains locations were all negative in the control rabbit. The brain sample was negative in the expressing rabbit, while spleen and all 10 liver samples were positive in the
5 expressing rabbit. The results suggest that liver and spleen have been transduced in the vector-treated rabbit, that marked cells are still present after 511 days, and that these are candidate organs for the source of the circulating factor VIII.

Juvenile mice were treated with DA/beta-gal vector as in Example 20 herein. Expressing mice were sacrificed at Day 59 or 105. Brain, liver, kidney, lung, and spleen
10 were analyzed for vector sequences by quantitative PCR. Quantitative PCR is performed as described above, except that 1 μ l of alpha-32P-dCTP is included in the PCR buffer, and a standard curve is prepared from the same positive control DNA in 3-fold dilutions from 1000 copies/sample down to 1.4 copies/sample, maintaining a constant concentration of 500 ng DNA/reaction by adding human PBMC DNA to each dilution. Five μ l of each reaction
15 is blotted onto Whatman DE81 Ion Exchange Chromatography paper, dried at 42°C x 5 min, washed 3x (40 ml 5M NaCl, 21 ml 1M monobasic sodium phosphate, 29 ml 1 M dibasic sodium phosphate, qs to 1 liter), and quantitated relative to the standard curve on the PhosphorImager (Molecular Dynamics). Liver was positive in all four animals tested, spleen in 3/4, and kidney in 1/4. Lung and brain were negative in all four, and ovary was negative
20 in the two animals tested (See Figure 28). Tissue lysates were also prepared from several animals. One third to 1/2 of each tissue was placed in homogenization buffer (Dianon PCNA ELISA kit) and stored on ice, until all of the samples were collected. Samples were liquefied in dounce homogenizers. The lysates were transferred to 1.5 ml microfuge tubes, and centrifuged at 14,000 rpm, 4°C, for 15 minutes. The aqueous layer between the lipid
25 and cell debris layers was aspirated by syringe and stored at -80°C. Samples were thawed at 37°C, diluted 1:3 and 1:10, and tested in duplicate for hGH by ELISA (BMB kit). Protein in each sample was determined in a Bradford protein assay, and results are expressed relative to total protein levels. Growth hormone antigen was found in liver in 4/4 animals tested, in spleen in 2/4, and kidney in 1/4. Brain was negative in 2/2 animals tested (See
30 Figure 29). These data are consistent with the proposal that liver and spleen are the major

sites of transduction and synthesis of genes of interest when retroviral vectors are introduced intravenously.

Example 24

Long Term Expression of Human Factor VIII in Normal Dogs Following Intravenous

Expression of Vector

5 Recombinant retroviral vector expressing B domain deleted factor VIII was constructed, packaged, and expressed as described in Example 2 herein. Large scale production was carried out in Cell Factories as described for DX/ND7 in Example 14 herein. The retroviral vector was then lactose formulated as described in Example 9 herein.
10 High titer formulated retroviral vector was injected into 8 week old normal juvenile dogs according to the following schedule: 4E8 vector particles were injected into the cephalic vein at 2.5 hr intervals, three times per day for three days (Day 0, 1, and 2). On days 4, 7, 10, 13, 20 and every 7 days subsequently, citrated plasma was obtained by venipuncture of the cephalic vein and analyzed for human factor VIII antigen as in Example 18, except that
15 the diluent was 1:3 citrated dog plasma (Harlan).

 Data from Day 0 (prebleeds) to day 279 from 2 experimental and one control dog are shown in Figure 30. Levels remained undetectable through Day 7, with no significant differences being seen between the experimental and control groups ($p > 0.25$). On Day 7 and 10, low levels (3-10 ng/ml) of factor VIII were observed in all three dogs. On day 13
20 and subsequent days until day 90, Control Dog 39 and Experimental Dog 47 ("nonresponder") displayed 0-8 ng/ml human factor VIII antigen in their plasmas. In contrast, Experimental Dog 55 ("responder") displayed levels of 40-110 ng/ml factor VIII. Levels of factor VIII in Dog 55 peaked around Day 160, gradually declining until Day 279, the last date tested. On day 90, 91, and 92, the nonresponder dog #47 was boosted with
25 dosage schedule recapitulating that used on day 0, 1, and 2. On Day 95, dog 47 displayed 95 ng/ml, comparable to the 77 ng/ml found in the unboosted responder dog 55 on that date. Unlike Dog 55, boosted Dog 47 did not maintain these levels, which declined to barely detectable levels by Day 125.

Example 25

Transduction of Splenocytes and Possible Stem Cell Transduction by Intravenous Administration of Retroviral Vector

5 Data in Example 24 herein show transduction of spleen and liver in long-term expressing animals treated with retroviral vector intravenously. To test for expression of hGH by splenocytes, an adoptive transfer experiment was carried out. Fragments of spleens (1/4 to 1/3 spleen) were collected from BALB/c mice which had received 1×10^7 cfu DA-827 51 days earlier via tail vein injection as described in Example 20 herein. Spleens were 10 processed individually. Single-cell suspensions were prepared in sterile Hank's balanced salt solution (HBSS; Irvine Scientific) by passing tissue fragments through a 70 mm nylon mesh strainer (Becton Dickinson #2350). After pelleting, cells from each spleen were resuspended in 0.8 ml HBSS; 0.3 ml of this suspension was injected intravenously into lethally irradiated (900 R) recipient BALB/c mice. An estimated $6 - 7 \times 10^6$ cells were 15 transferred to each recipient. Serum levels of hGH were monitored every two weeks in recipients as in Example 20. levels of hGH expression in recipients roughly corresponded to the levels that had been present in the original donors (See Figure 31), suggesting that these levels correlated with transduced cell number. Peak expression was observed 56 days following spleen cell transfer, with a decrease by day 70. The increase of expression with 20 time as the animals reconstituted indicates that stem cells have been transduced.

Example 26

Expression of Human Factor VIII in Hemophiliac Dogs Following Intravenous Expression of Vector

25 Two factor VIII-deficient hemophiliac dogs (8 weeks old) were obtained from the closed colony at U. North Carolina, Chapel Hill. Two normal littermates were also obtained and used without further treatment to provide baseline values for assays. The two test dogs were injected with retroviral vector as in Example 24, herein, except that the injection dates were Days 1, 2, and 3. Blood samples were obtained on Days 4, 7, 10, 14, and every 7 days subsequently. Citrated samples were used for measurements of whole blood clotting time

(See Figure 32). On day -4, samples from the hemophiliac dogs were incoagulable (WBCT > 60 min), whereas the normal littermates had WBCT of 6-8 minutes. The WBCT of the normal littermates were maintained throughout the experiment. On day 4-10, the WBCT of the injected hemophiliac dogs dropped to 12-16 minutes. This represents expression of pg levels of factor VIII (Dr. Tim Nichols, UNC-Chapel Hill, personal communication). On day 14-28, the hemophiliac dog WBCT was again incoagulable (>50 min). Starting at day 35, the clotting time was again reduced, reaching 17 minutes for one dog and 16 minutes for the second, indicating a second peak of factor VIII clotting activity. This data demonstrates the successful expression of biologically active factor VIII in dogs by intravenous administration of a retroviral vector expressing a B-domain deleted factor VIII protein.

Example 27

Construction of pBA-5a, pBA-5b, and pBA-5c, PBA-9b and pBA-8bL1 Retroviral Vector Backbones

This example describes several modifications of the retroviral vector pKT1 resulting in decreased sequence homology to the retroviral gag/pol and envelope expression constructs. pKT-1 is described in Example 1 herein and is shown schematically in Figure 33. (The construction of the pKT-1 starting material used in the present example is further described in detail in PCT WO 95/30763 and in co-owned U.S. Serial No. 08/721,327, both of which are hereby incorporated by reference in their entirety. In addition, two stop codons were introduced in the DNA sequence of the packaging signal sequence in order to increase the safety of these vectors. All modifications are summarized in Fig. 36 and the resulting retroviral backbones are called pBA-5a, pBA-5b, and pBA-5c. Further details on the construction of pBA-5a, pBA-5b, and pBA-5c are provided in co-owned U.S. Serial No. 08/721,327 and co-owned application, attorney docket 1147.004, filed May 5, 1997, entitled "Crossless Retroviral Vectors" and which is also hereby incorporated by reference.

A. Substitution of Nonsense Codons in the Extended Packaging Sequence (Ψ^+)

This example describes modification of the extended packaging signal (Ψ^+) by PCR on the template KT-1 using primers that introduce two stop codons in the extended

packaging signal sequence. In particular, the template pKT-1 contains the modification ATT at the normal ATG start site of *gag* (position 621-623 of SEQ ID NO: 15). Here the start site was further modified to the stop codon, TAA, and an additional stop codon TGA was added to replace the codon TTA at position 645-647 of SEQ ID NO: 15.

5 Briefly, two sets of PCR reactions were carried out on pKT1, each introducing one stop codon. The primers for the PCR were designed such that the two PCR products had overlapping regions and a splice-overlap extension PCR (SOE-PCR) was carried out with the two PCR products in order to combine the two introduced stop codons on one strand. The first set of oligonucleotides introducing the change from ATT to TAA were 5'-GGG-
10 AGT-GGT-AAC-AGT-CTG-GCC-TTA-ATT-CTC-AG (SEQ ID NO: 16) and 5'-CGG-TCG-ACC-TCG-AGA-ATT-AAT-TC (SEQ ID NO: 17) and the second set of oligonucleotides introducing the change from TTA to TGA were 5'-CTG-GGA-GAC-GTC-CCA-GGG-ACT-TC (SEQ ID NO: 18) and 5'-GGC-CAG-ACT-GTT-ACC-ACT-CCC-TGA-AGT-TTG-AC (SEQ ID NO: 19). The flanking primers of the final 708 base pair
15 PCR product introduced the *AatII* and the *XhoI* sites, at the 5' and 3', respectively.

The ends of the 708 base pair product were blunted and phosphorylated and the product introduced into the *SmaI* and *EcoRV* digested vector pBluescript SK- (Stratagene, San Diego, Calif.). The resulting plasmid was designated pBA-2, and is shown diagrammatically in Figure 34.

20

B. Removal of Retroviral Sequences Upstream and Downstream from the 3' LTR and Upstream and within the 5' LTR

Retroviral envelope sequence was removed upstream of the 3' LTR between the *Clal* site and the TAG stop codon of the envelope coding sequence. The DNA sequence was
25 modified by PCR such that the TAG stop codon was replaced by a *Clal* site and the 97 nucleotides upstream from this new *Clal* site to the original *Clal* site were deleted, as well as the 212 base pairs of retroviral sequence downstream of the 3' LTR.

Briefly, the following two oligonucleotides were used for the PCR: 5'-CATCGATAAA ATAAAAGATT TTATTTAGTC (SEQ ID NO: 20) and 5'-
30 CAAATGAAAG ACCCCCGCTG AC (SEQ ID NO: 21) and the template was pKT1. The

PCR product was cloned into pPCRII (Invitrogen, San Diego, Calif.) using the TA cloning kit (Invitrogen, San Diego, Calif.) and called pBA-1.

Subsequently, pBA-2 (described in section A above) was digested with *XbaI* and *AatII* which deleted a part of the multiple cloning site and into this linearized vector the 780 base pair fragment from *NheI* to *AatII* from pKT1 was cloned, resulting in the plasmid pBA-3. This plasmid pBA-3 combined the shortened 5' LTR with the above described packaging region including the two introduced stop codons.

Subsequently, pBA-1 was digested with *ClaI* and *ApaI* resulting in a 640 base pair fragment that was cloned into the *ClaI* and *ApaI* digested pBA-3 resulting in the plasmid pBA-4. This plasmid combines the above described 5' LTR and the packaging signal with the 3' LTR.

Subsequently, pBA-4 was digested with *ApaI* and *EcoRI*, ends blunted and religated in order to remove extraneous 3' polylinker sites, resulting in plasmid pBA-5a.

Subsequently, pBA-5a was cut with *NotI* (blunted) and *EcoRI* and introduced into *SmaI* and *EcoRI* digested pUC18 (GIBCO/BRL, Gaithersburg, MD) resulting in pBA-5b. This construct moved the retroviral vector from a pBluescript into an alternate pUC18 vector.

pBA-5c is constructed as identical to pUC-18 derived pBA-5b except for the *XhoI/ClaI* multicloning site was introduced into pUC-19.

This example further describes several modifications of the retroviral vector pBA-5b which result in a vector with multiple unique restriction enzyme sites for convenient cloning. In order to prepare the pBA-9b vector, the herpes simplex virus thymidine kinase (HSVTK) gene was retrieved by digesting pBH-1 with *XhoI* and *EcoRI* resulting in a 1.2kb fragment. (pBH-1 was prepared as described in WO 91/02805 entitled "Recombinant Retroviruses Delivering Vector Constructs to Target Cells" which is hereby incorporated by reference.) The neomycin gene driven by the SV40 promoter was retrieved by digesting pKT-1 with *EcoRI* and *BstBI* resulting in a 1.3 kb fragment. Both fragments were cloned into a *XhoI* and *ClaI* digested pBA-5b resulting in the retroviral vector pMO-TK.

The TK gene from the retroviral vector pMO-TK was isolated as a *XhoI-ClaI* fragment and inserted into the *XhoI-ClaI* digested pBA-5b, resulting in the plasmid pBA-

5bTK. In order to delete the restriction enzyme sites HindIII, SphI, PstI, SalI and HincII upstream of the 5' LTR, pBA-5bTK was digested with HindIII and HincII, overhanging ends removed using T4 polymerase and blunt ends ligated using the T4 DNA ligase, resulting in plasmid pTJBA-5bTK with 16 bases (TGC ATG CCT GCA GGT C) removed from upstream of the 5'LTR. The plasmid pTJBA-5bTK has a BamHI upstream of the 5'LTR. It is desirable to remove this BamHI site since it is a common site used for cloning. In order to destroy the BamHI site upstream of the 5'LTR, the BamHI-containing TK gene in pTJBA-5bTK was replaced by the IL-2 gene via XhoI-ClaI digest, resulting in plasmid pTJBA-5bIL-2. The plasmid pTJBA-5bIL-2 was digested with BamHI, the ends filled in with the Klenow fragment and the ends religated, resulting in pTJBA-5bIL-2 (BamHI del.).

In order to produce the plasmid pBA-9b, the IL-2 gene from pTJBA-5bIL-2 (BamHI del.) is deleted via XhoI-ClaI digest and replaced with a linker that introduces a multiple cloning site (MCS) and codes for the restriction enzyme sites 5' -XhoI-ApaI-BglII-NotI-NruI-SalI-HindIII-BamHI-ClaI- 3'. The sequence of the two primers that is used to produce the linker are as follows: 5' - TCG AGG GGC CCA GAT CTG CGG CCG CTC GCG AGT CGA CAA GCT TGG ATC CAT - 3' (Seq ID No. 26) as the primer for the + strand and 5' -CGA TGG ATC CAA GCT TGT CGA CTC GCG AGC GGC CGC AGA TCT GGG CCC C - 3' (Seq ID No. 27) as the primer for the - strand.

This example describes several modifications of the retroviral vector pBA-5b which result in a vector coding for the human placental alkaline phosphatase gene (PLAP), driven by the SV40 promoter.

The plasmid pBA-8bL1 was constructed in a three-way ligation using the following three fragments: The NdeI-ClaI fragment from pBA-5b (described above) containing the 3'LTR and the pUC18 backbone, the ClaI-HindIII fragment from pCI-PLAP coding for PLAP and the HindIII-NdeI fragment from pBA-6bL1 containing the 5'LTR and the SV40 promoter. Plasmid pBA-6bL1 is based on pBA-6b (described above) where the HIVenv/rev coding region was deleted via XhoI-ClaI digest and replaced with the L1 linker coding for several restriction enzyme sites including XhoI at the 5'end and ClaI at the 3'end.

Example 28

Preparation Of pBA-5a, pBA-5b, and pBA-5c Retroviral Vectors Expressing
B-Domain Deleted Factor VIII

A B-domain deleted factor VIII cDNA fragment thus constructed was obtained by XhoI/NotI digestion, as described below. A retroviral vector (pMBF8) expressing B-
5 domain-deleted factor VIII is constructed from the expression plasmid pSVF8-200 which is prepared as described in Truett, 1985, *DNA* 4:333) and in U.S. Patent No. 5,045,455, and which was deposited with the ATCC on July 17, 1985, and which has been assigned ATCC number 40190.

A DNA fragment encoding the β -domain deleted Factor VIII molecule was obtained
10 from pSVF8-302, which has a nine base pair deletion in the 5' non-coding region after the poly G tail. Plasmid SVF8-302 was constructed in a similar manner as pSVF8-200, which is described in detail in Truett and in U.S. Patent No. 5,045,455. Construction of pSVF8-302 is also described in U.S. Patent No. 5,595,886.

The procedure outlined below describes the construction of retroviral vectors
15 expressing a β -domain deleted Factor VIII protein obtained from pSVF8-302. However, alternatively, the same procedure can be used to construct such retroviral vectors from pSVF8-200.

The full-length cDNA sequence of the factor III is shown in Seq ID No. 44 and the full-length amino acid sequence is shown in Seq ID No. 45. The cDNA sequence of the B-
20 domain deleted SQN deletion is shown in Seq ID No. 46 and the SQN deletion amino acid sequence is shown in Seq. ID No. 47.

Fragment 1, encompassing nt 5500-6248 of pSVF8-200 (see Figure 8 of Truett), was obtained by VENT-PCR using factor VIII primers encoding a PFIM1 site at the 3' end and the 5' SQN sequence plus a HindIII site at the 5' end. The 5' primer encompasses the region
25 2446-2460 of the 5' SQN and the region 5144-5167 just downstream of the 3' SQN sequence. Thus, this fragment spans the sequence between the two SQN sites within the B domain (positions 2461 and 5142). The particular primer sequences used were:

1. 5' GAAGCTTCTCCCAGAACCCACCAGTCTTGAAACGCCATC (Seq ID No. 22)
2. 5' GTACCAGCTTTTGGTCTCATCAAAG (Seq ID No. 23).

Fragment 1 was blunt-end cloned into vector SK- that had been cut with *Sma*I and dephosphorylated, forming pSK-Pfl. Fragment 2, encompassing nt 1190 and 2448 was isolated following *Hind*III digestion and cloned into the *Hind*III site of SK-Pfl to form SK-Pfl-*Hind*. The orientation of the insert was determined using *Acc*I and *Pst*I digests. pSVF8-200 was digested with *Hpa*I and religated to remove two small *Hpa*I fragments 3' to the factor VIII cDNA insert, forming pF8-300-del-Hpa. The remaining *Hpa*I site was converted to a *Not*I site using *Not*I phosphorylated linkers, forming F8-300-Hpa/*Not*. Fragment 3, encompassing nt 5885-7604, was isolated following PflMI and *Not*I digestion and cloned into SK-Pfl-*Hind* following PflMI and *Not*I digestion of the latter to form pF8:213.

Fragment 4 (encompassing nt 104-133 to 1204) was obtained following VENT-PCR of PSV7dF8-300 with primers containing 5' *Xho*I and 3' *Acc*I sites respectively. The 5' primer encompasses nt 104-133 and the 3' primer encompasses nt 1200-1224. pF8:213 was digested sequentially with *Xho*I followed by *Acc*I and ligated to Fragment 4 which was digested with *Xho*I and *Acc*I, pF8:4213. The primer sequences for the 5' UT and *Xho*I primers were:

3. 5' CTCCTCGAGCTAAAGATATTTAGAGAAGAATTAAC (Seq ID No. 24)
4. 5' TTCCTCTGGACAGCTGTCTACTTTG (Seq ID No. 25)

The crossless backbones pBA-9b, pBA-5a, pBA-5b and pBA-5c were modified by linearizing with *Cla*I, blunt ending and religating in the presence of *Not*I phosphorylated linkers. The modified cDNA fragment was cloned into the *Xho*I/*Not*I linearized vectors. Similarly, the modified cDNA described above is cloned into the backbone pMBA backbone described above which has been digested with *Xho*I and *Not*I

Example 29

Construction of Recombinant Adeno-Associated Virus (rAAV) Vectors that Express the Heavy and Light Chains of Human Factor VIII

We constructed two rAAV vectors, one expressed the light chain, and the other expressed the heavy chain. Both chains contain the Factor VIII leader sequence and a variable amount of the B domain.

To clone the heavy chain, a region of the Factor VIII gene from 174 bp 5' of the ATG to amino acid 745 was amplified by PCR. This fragment includes the entire heavy chain and first five amino acids of the B domain. The oligos used were: forward, 5'-CACCGTCGTCGACTTATGCT-3' (Seq ID No. 28), and reverse, 5'-GACCGTCGACTCAATTCTGGGAGAAGCTTCTTGG-3' (Seq ID No. 29). The plasmid used as a template in the PCR reaction was pCMVKmHSTB, a B deleted factor VIII expression construct. The amplified fragment was digested with Sal I, and cloned into a CMV expression vector, pCMVKmLINK digested with Sal I and Xho I. This plasmid was called pCMVKm90H. pCMVKmLINK is an expression vector containing the CMV promoter/intron, a polylinker for cloning genes of interest, and a bovine growth hormone polyadenylation signal. To make a rAAV vector expressing the heavy chain, pCMVKm90H was digested with Sal I and Bam HI, the Bam HI site was filled in with T4 DNA polymerase, and this fragment was cloned into the rAAV vector pKm201CMV-CI digested with Sal I and Eco RV. pKm201CMV-CI contains the inverted terminal repeats of AAV, the CMV promoter, the chimeric intron from pCI (Promega, Madison, WI), and the bovine growth hormone polyadenylation signal. The final AAV vector was called pKm201-90H.

To clone the light chain, we first amplified the factor VIII 5' untranslated and leader region using the following primers: forward, 5'-CACCGTCGTCGACTTATGCT-3' (Seq ID No. 30), and reverse, 5'-CAACGCTCGAGAAGCAGAATCGCAAAGGC-3' (Seq ID No. 31). Again, pCMVKmHSTB was used as a template in the PCR reaction. The amplified region includes sequences from 174 bp upstream of the ATG to amino acid 19 of factor VIII. This fragment was digested with Xho I and Sal I and cloned into pCMVKmLINK digested with Xho I and Sal I. This plasmid was called pCMVKmF8L (for factor VIII leader). To amplify the light chain, the following primers were used: forward, 5'-TCGGCTCGAGGCATCAACGGGAAATAACTCGT-3' (Seq ID No. 32), and reverse, 5'-CCGACTCGAGTCAGTAGAGGTCCTGTGCCTC-3' (Seq ID No. 33). Again, pCMVKmHSTB served as the template for the PCR. The amplified fragment included sequences from amino acid 1645 of factor VIII to the STOP codon after amino acid 2332. This included the last four amino acids of the B domain and the complete light chain. This

fragment was digested with Xho I and cloned into the Xho I site of pCMVKmF8L. This resulted in a light chain construct containing the factor VIII leader which was called pCMVKm80L. pCMVKm80L was digested with Sal I and Bam HI to remove the light chain construct, and this fragment was cloned into pKm201CMV-CI digested with Sal I and Bam HI to generate pKm201-80L. See Figures 35 and 36 for diagrams pKm201-80L and pKm201-90H, respectively.

Example 30

Co-Infection of Cells with rAAV Vectors Expressing the Heavy and Light Chains of Factor VIII Results in the Production of Biologically Active Factor VIII

10 The heavy and light chain constructs, pKm201-80L and pKm201-90H, were packaged following standard procedures for the production of rAAV (Zhou *et al.*, 1994, *J. Exp. Med.* 179:1867-1875). rAAV was purified as described (Wang *et al.*, 1995, *Proc. Natl. Acad. Sci. USA* 92:12416-12420) and used to infect 293 cells plated in 6-well plates. Supernatants of infected cells were collected at 48 h after infection and assayed for

15 biologically active factor VIII by Coamatic Factor VIII (KabiVitrum, Stockholm) following manufacturer's instructions. Normal human plasma (George King Bio-Medical, Inc., Overland Park, Kansas) was used to generate a standard curve. Cells were infected at a multiplicity of infection (MOI) of 6000 rAAV particles per cell. The experiment was done both with and without the addition of etoposide (0.3 M) to the medium. Etoposide has been

20 shown to increase the transduction efficiency of rAAV vectors (Russell *et al.*, 1995, *Proc. Natl. Acad. Sci. USA* 92:51719-51723). As shown in Table 5 below, co-infection of rAAV-80L and rAAV-90H resulted in the production of biologically active factor VIII. The amount of factor VIII was increased in the presence of etoposide.

Table 5: Co-infection of rAAV-80L and rAAV-90H results in the production of biologically active factor VIII.

Virus	Factor VIII (ng/ml)	
	- etoposide	+ etoposide
80L	0	0
90H	0	0
80L + 90H	1.4	6.1

5

Example 31

Preparation of Retroviral Vectors with Liver-Specific Promoters

(a) Preparation of Interferon -alpha Sequences Utilizing PCR

Construction of the Retroviral Vector Encoding Interferon- α 2a, α 2b, α 2c, α 54, and α 76

DNA clones of interferon- α 2a, α 2b, α 2c α 54, and α 76 are prepared utilizing PCR.

10 The interferon- α 2a DNA is obtained from a cDNA library as described by Goeddel *et al*, 1980, *Nature* 287:411-416. The interferon- α 2b DNA is obtained from a cDNA library described by Streuli *et al.*, 1980, *Science* 209:1343-1347, and the interferon- α 2c DNA is obtained as described by Saveliev *et al.*, 1986, *Antibiot. Med. Biotechnol.* 3:592-596. DNA clones of interferon- α 54 and α 76 are prepared utilizing PCR. Both the interferon- α 54 DNA
15 and the interferon- α 76 DNA are obtained as described in U.S. Patent No. 4,975,276 and U.S. Patent No. 5,098,703.

A reaction mixture is then prepared according to procedures specified by New England Biolabs (Beverly, MA). More specifically, a reaction mixture is prepared containing 1 ug purified plasmid, 10 ul of 10X ThermoPol reaction buffer, 2 ul 2.5 mM of
20 each dATP, dCTP, dGTP and dTTP, 0.5 ul of 2 units/100 ul Vent polymerase, 1 - 3 ul of 100mM MgSO₄, and 0.5-1.0 ug of the primers specified below (Mattila *et al.*, 1991, *Nucleic Acids Research* 19:4967-4973, Eckert, K.A. and Kunkel, T. A., PCR Methods and Applications 1, 17-24, 1991).

a. Interferon- α 2a and α 2b

For interferon- α 2a and α 2b, the coding region is identical except for AAA codon at position 23 for interferon- α 2a and AGA codon at position 23 for interferon- α 2b. The sense
5 primer for both interferon- α 2a and α 2b is from the 5' region of the coding sequence, 6 bp upstream from the ATG start codon until 25 bp downstream from the start codon. The 5' end of the primer contains the Xho I restriction site.

(Seq ID No. 36) 5'-3': CGCG CCG CTC GAG TCT ACA ATG GCC TTG ACC TTT GCT TTA CTG G

For interferon- α 2a and α 2b, the second primer corresponds to the anti-sense nucleotide
10 sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Cla I restriction site.

(Seq ID No. 37) 5'-3': GCG CCC ATC GAT TCA TTC CTT ACT TCT TAA ACT TTC TTG CAA G

b. Interferon- α 2c

For interferon- α 2c, the sense primer is from the 5' region of the coding sequence, 7 bp
15 upstream from the ATG start codon until 28 bp downstream from the start codon. The 5' end of the primer contains the Xho I restriction site.

(Seq ID No. 38) 5'-3': CGCG CCG CTC GAG CAT CCA ATG GCC CTG TCC TTT TCT TTA CTT ATG G

For interferon- α 2c, the second primer corresponds to the anti-sense nucleotide sequence,
20 extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Cla I restriction site.

(Seq ID No. 39) 5'-3': CC ATC GAT TCA ATC CTT CCT CCT TAA TCT TTT TTG CAA G

c. Interferon- α 54

The sense primer is from the 5' region of the coding sequence, 6 bp upstream from the ATG
25 start codon until 24 bp downstream from the start codon. The 5' end of the primer contains the Xho I restriction site.

(Seq ID No. 53) 5'-3': CGCG CCG CTC GAG TCT ACA ATG GCT TTG CCT TTT GCT TTA CTG

The second primer corresponds to the anti-sense nucleotide sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Cla I restriction site.

(Seq ID No. 54) 5'-3': GCG CCC ATC GAT TTA TTC CTT CCT CCT TAA CCT TTC TTG CAA G

5 d. Interferon- α 76

For interferon-76, the sense primer is from the 5' region of the coding sequence, 7 bp upstream from the ATG start codon until 28 bp downstream from the start codon. The 5' end of the primer contains the Xho I restriction site.

(Seq ID No. 55) 5'-3': CGCG CCG CTC GAG CAT CCC AAT GGC CCT GTC CTT TTC TTT ACT GAT

10 GG

The second primer corresponds to the anti-sense nucleotide sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Cla I restriction site.

(Seq ID No. 56) 5'-3': CC ATC GAT TCA ATC CTT CCT CCT TAA TCT TTT TTG CAA G

15 The reaction mixture is brought up to 100 μ l with DI H₂O, and each tube is placed into a thermocycler (Gene Amp PCR System 9600, Perkin-Elmer, Cetus, Calif.). The PCR program regulates the temperature of the reaction vessel first at 94°C for 10 minutes to hot start reactions. Vent polymerase is added next and the cycle begins with 94°C for 2 minutes, next at 56°C for 30 - 60 seconds, and 72°C for 30 - 60 seconds. This cycle is
20 repeated 35 times. After the 35th cycle, product ends are sealed by the reactions at 72°C for 3 - 5 minutes. Reactions are held at 4°C. If necessary, PCR amplification can be optimized using touchdown or stepdown PCR protocol (White, B. A., PCR Cloning Protocols 67, 39-45, 1997).

The PCR product from the reaction yields the interferon- α 2a, α 2b and α 2c gene.

25 Following the PCR reaction, the solution is transferred to a fresh 1.5 ml microfuge tube. Fifty microliters of 3 M sodium acetate is added to this solution followed by 500 μ l of chloroform:isoamyl alcohol (24:1). The mixture is vortexed and then centrifuged at 14,000

rpm for 5 minutes. The aqueous phase is transferred to a fresh microfuge tube and 1.0 ml 100% EtOH is added. This solution is incubated at -20°C for 4.5 hours, and then centrifuged at 10,000 rpm for 20 minutes. The supernatant is decanted, and the pellet rinsed with 500 ul of 70% EtOH. The pellet is dried by centrifugation at 10,000 rpm under
5 vacuum and then resuspended in 10 ul deionized H₂O. One microliter of the PCR product is analyzed by electrophoresis in a 1.0% agarose gel.

These PCR products, approximately 570 bp in length, are digested with Xho I and Cla I restriction endonucleases, electrophoresed through a 1.0% agarose gel and the DNA is purified from the gel slice by GeneClean II (Bio 101, Vista, California). These Xho I-Cla I
10 PCR products are subcloned into the respective sites of pBluescript KSII+ (Stratagene, La Jolla, California). These constructs are designated KSII+ Xho-Cla IFN- α 2a, α 2b, α 2c, α 54 and α 76 respectively, and are verified by DNA sequencing.

(b) Insertion of interferon-alpha sequences into retroviral vectors

1. Retroviral Vector Backbone - pBA6bL1

15 The pBA6bL1 retroviral vector backbone is described above. Two fragments are purified for the construction of the interferon α 2a, α 2b, α 2c, α 54 and α 76 pBA6bL1 retroviral vector. First, pBA6bL1 is digested with Xho I and Cla I and the 6.2 kb fragment containing the retroviral vector backbone is isolated. Second, from KSII+ Xho-Cla IFN- α 2a, α 2b, α 2c, α 54 and α 76, the 570 Xho I-Cla I bp fragment containing the interferon- α 2a, α 2b,
20 α 2c, α 54 and α 76 gene is isolated and inserted into the Xho I-Cla I sites of pBA6bL1. The vector constructs are designated pBA6b-IFN α 2a, pBA6b-IFN α 2b, pBA6b-IFN α 2c, pBA6b-IFN α 54 and pBA6b-IFN α 76, respectively.

25 2. Retroviral Vector Backbone - pLXSN

The retroviral vector backbone, pLXSN is available by Clontech (Palo, Alto, Calif.). Two fragments are purified for the construction of the LXSN retroviral vector encoding interferon- α 2a, α 2b, α 2c, α 54 and α 76. First, pLXSN is digested with Xho I and Bam HI and the 5.9 kb fragment containing the retroviral vector backbone is isolated. Second, from

KSII+ Xho-Cla IFN- α 2a, α 2b, α 2c, α 54 and α 76, the 570 bp Xho-Bam HI fragment containing either the interferon- α 2a, α 2b, α 2c, α 54 or α 76 gene is isolated and inserted into the respective sites of pLXSN. This vector construct is designated pLXSN- IFN α 2a, pLXSN-IFN α 2b, pLXSN-IFN α 2c, pLXSN- IFN α 54 and pLXSN-IFN α 76.

5

3. Lentiviral Backbone

An HIV retroviral vector backbone is made, similar to V653RSN as described by Parolin *et al.* (*J. Virol.* 68:3888-3895, 1994). From KSII+ Xho-Cla IFN- α 2a, α 2b, α 2c, α 54 and α 76, the 570 bp Xho I-Xba I fragment containing the interferon- α 2a, α 2b, α 2c, α 54 and α 76 is isolated and inserted into the Xho I-Xba I sites of a CMV expression vector, pCI (Promega, Madison, WI) and is designated pCI -IFN α 2a, IFN α 2b, IFN α 2c, IFN α 54 and IFN α 76, respectively. From pCI- IFN α 2a, IFN α 2b, IFN α 2c, IFN α 54 and IFN α 76, the Bgl II-Bam HI fragment is isolated and inserted into the Bam HI sites of V653RSN where the SL3NEO sequences have been removed. To determine the orientation of the interferon α insert, vector constructs are sequenced. These vector constructs are designated V653-IFN α 2a, V653-IFN α 2b, V653-IFN α 2c, V653-IFN α 54 and V653-IFN α 76, respectively.

15

4. Retroviral Vector Backbone - pBA8bL1

The pBA8bL1 retroviral vector backbone has been described before in Example 27.

20

a. Interferon- α 2a and α 2b

In a separate set of PCR reactions, the sense primer for both interferon- α 2a and α 2b is from the 5' region of the coding sequence, 6 bp upstream from the ATG start codon until 25 bp downstream from the start codon. The 5' end of the primer contains the Bam HI restriction site.

25 (Seq ID No. 40) 5'-3': CC GGA TCC TCT ACA ATG GCC TTG ACC TTT GCT TTA CTG G

For interferon- α 2a and α 2b, the second primer corresponds to the anti-sense nucleotide sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Not I restriction site.

30 (Seq ID No. 41) 5'-3': CGCG CCG GCG GCC GC TCA TTC CTT ACT TCT TAA ACT TTC TTG CAA G

b. Interferon- α 2c

For interferon- α 2c, the sense primer is from the 5' region of the coding sequence, 7 bp upstream from the ATG start codon until 28 bp downstream from the start codon. The 5' end of the primer contains the Bam HI restriction site.

5 (Seq ID No. 42) 5'-3': CC GGA TCC CAT CCA ATG GCC CTG TCC TTT TCT TTA CTT ATG G

For interferon- α 2c, the second primer corresponds to the anti-sense nucleotide sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Not I restriction site.

(Seq ID No. 43) 5'-3': CGCG CCG GCG GCC GC TCA ATC CTT CCT CCT TAA TCT TTT TTG CAA G

10 **c. Interferon- α 54**

In a separate set of PCR reactions, the sense primer for both interferon- α 54 is from the 5' region of the coding sequence, 6 bp upstream from the ATG start codon until 25 bp downstream from the start codon. The 5' end of the primer contains the Bam HI restriction site.

15 (Seq ID No. 57) 5'-3': CC GGA TCC TCT ACA ATG GCT TTG CCT TTT GCT TTA CTG

For interferon- α 54, the second primer corresponds to the anti-sense nucleotide sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Not I restriction site.

(Seq ID No. 58) 5'-3': CGCG CCG GCG GCC GC TTA TTC CTT CCT CCT TAA CCT TTC TTG CAA G

20 **d. Interferon- α 76**

For interferon- α 76, the sense primer is from the 5' region of the coding sequence, 7 bp upstream from the ATG start codon until 28 bp downstream from the start codon. The 5' end of the primer contains the Bam HI restriction site.

(Seq ID No. 59) 5'-3': CC GGA TCC CAT CCC AAT GGC CCT GTC CTT TTC TTT ACT GAT GG

For interferon- α 76, the second primer corresponds to the anti-sense nucleotide sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Not I restriction site.

(Seq ID No. 60) 5'-3': CGCG CCG GCG GCC GC TCA ATC CTT CCT CCT TAA TCT TTT TTG CAA G

5 These PCR products, approximately 570 bp in length, are digested with Bam HI and Not I restriction endonucleases, electrophoresed through a 1.0% agarose gel and the DNA is purified from the gel slice by GeneClean II (Bio 101, Vista, California). This Bam HI-Not I PCR product is subcloned into the respective sites of pBluescript KSII+ (Stratagene, La Jolla, California). These constructs, designated KSII+ Bam-Not IFN- α 2a, α 2b, α 2c, α 54
10 and α 76 are verified by DNA sequencing.

Two fragments are purified for the construction of the interferon α 2a, α 2b, α 2c, α 54 and α 76 pBA8bL1 retroviral vector. First, pBA8bL1 is digested with Not I and Srf I and the 6.1 kb fragment containing the retroviral vector backbone is isolated. Second, from KSII+ Bam-Not IFN α 2a, α 2b, α 2c, α 54 and α 76, the 570 bp Sma I-Not I fragments
15 containing the interferon- α 2a, α 2b, α 2c, α 54 or α 76 are isolated and inserted into the Srf I-Not I sites of pBA8bL1. These vector constructs are designated pBA8b-IFN- α 2a, pBA8b-IFN- α 2b, pBA8b- α 2c, pBA8b-IFN- α 54 and pBA8b-IFN α 76.

5. Retroviral Vector Backbone - pBA6bL1/TK

The herpes simplex virus (HSV) thymidine kinase (TK) cDNA is excised from
20 pTJBA-5bTK retroviral vector by Xho I/Cla I double digestion as described before in Example 27B and inserted into the Sal I/Cla I sites of pSP72 plasmid (Promega, Madison, WI) and designated pSP72-TK. The human interleukin-2 (IL-2) cDNA is excised from KT3-IL-2 (described in PCT Patent Publication Nos. WO 94/21792 and WO 96/21015) and inserted into the Xho I/Cla I sites of pBluescript SK- and designated pBSSK-IL2. The Hind
25 III/Sma I double digest of pSP72-TK excises TK cDNA and this fragment is inserted into the Hind III/Sma I sites of pBSSK-IL2. This construct has the following: Xho I-IL2 cDNA-Cla I-Hind III-TK cDNA-Sma I sequence and is designated pBSSK-IL2/TK.

From KSII+ Xho-Cla IFN- α 2a, α 2b, α 2c, α 54 or α 76, the 570 Xho I-Cla I bp fragment containing the interferon- α 2a, α 2b, α 2c, α 54 or α 76 gene is isolated. From pBSSK-IL2/TK, the IL2 fragment is excised by Xho I/Cla I double digestion, and replaced with Xho/Cla I fragments containing the interferon- α 2a, α 2b, α 2c, α 54 or α 76 genes. The vector constructs are designated pBSSK-IFN α 2a /TK, pBSSK-IFN α 2b /TK, pBSSK-IFN α 2c /TK, pBSSK-IFN α 54/TK and pBSSK-IFN α 76/TK, respectively. The Xho I/Not I fragments from pBSSK-IFN α 2a /TK, pBSSK-IFN α 2b /TK, pBSSK-IFN α 2c /TK, pBSSK-IFN α 54 /TK and pBSSK-IFN α 76 /TK are excised and inserted into the Xho I-Not I sites of pBA6bL1. These di-cistronic retroviral vectors encoding IFN α 2a, α 2b, α 2c α 54 and α 76 are designated pBA6bL1 IFN α 2a/TK, pBA6bL1 IFN α 2b/TK, pBA6bL1 IFN α 2c/TK, pBA6bL1 IFN α 54/TK and pBA6bL1 IFN α 76/TK.

6. Retroviral Vector Backbone with a Liver-specific promoter

The ApoE enhancer-alpha 1-antitrypsin promoter yields high levels of expression from liver cells (Okuyama *et al.*, 1996, *Hum Gene Ther* 7:637-645). The retroviral vector encoding the liver-specific promoter is constructed as follows.

a. ApoE enhancer cassette

The apolipoprotein E (Apo E) enhancer sequence has been identified as a 154 bp fragment by Shachter *et al.*, 1993, *J. Lipid Res* 34:1699-1707. The ApoE enhancer cassette was generated by sets of four oligonucleotides that span the entire sequence. For the first reaction, four oligonucleotides were synthesized that were phosphorylated at the 5' end. The first oligonucleotide is the sense strand and contains Hind III restriction site at the 5' end.

(Seq ID No. 61) 5'-3': AG CTT GCT GTT TGT GTG CTG CCT CTG AAG TCC ACA CTG AAC AAA
CTT CAG CCT ACT CAT GTC CCT AAA ATG GGC AAA CAT TGC AAG CAG C

The second oligonucleotide is the sense strand, spanning the second half of the Apo E enhancer and contains the Hind III restriction site at the 3' end.

(Seq ID No. 62) 5'-3': AAA CAG CAA ACA CAC AGC CCT CCC TGC CTG CTG ACC TTG GAG CTG
GGG CAG AGG TCA GAG ACC TCT CTG A

The third oligonucleotide is the antisense strand of the second half of the Apo E enhancer and contains the Hind III site at the 5' end.

5

(Seq ID No. 63) 5'-3': AG CTT CAG AGA GGT CTC TGA CCT CTG CCC CAG CTC CAA GGT CAG
CAG GCA GGG AGG GCT GTG TGT TTG CTG TTT GCT GCT TG

The fourth oligonucleotide is the antisense strand of the first half of the Apo E enhancer and contains the Hind III site at the 3' end.

10

(Seq ID No. 64) 5'-3': CAA TGT TTG CCC ATT TTA GGG ACA TGA GTA GGC TGA AGT TTG TTC
AGT GTG GAC TTC AGA GGC AGC ACA CAA ACA GC A

This set of four oligonucleotides of approximately 80 bases in length are annealed,
and ligated into the dephosphorylated Hind III sites of pBluescript KSII+ (Stratagene, La
Jolla, California) following standard molecular cloning protocols. Recombinants are
selected and sequenced. Correct monomer Apo E enhancer elements are excised by Hind
III, electrophoresed through a polyacrylamide gel and the DNA is purified following
procedures described by Asubel et al., *Current Protocols in Molecular Biology*, Greene
Publishing and Wiley-Interscience, New York (1987). The monomer enhancer element is
ligated into the dephosphorylated Hind III sites of pBluescript KSII+ at an insert:vector
ratio of 10:1 in order to generate recombinants containing dimers. Recombinant plasmids
were assessed by DNA sequencing to select clones which carried both enhancer fragments
in the same orientation with the 5' end closer to the Eco RV site and the 3' end closer to the
Cla I site within the multi-cloning site of pBluescript. This construct is designated KSII+
Hind III ApoE(x2)

25

For the second reaction, another set of four oligonucleotides were synthesized that
were phosphorylated at the 5' end. The first oligonucleotide is the sense strand and contains
Eco RI restriction site at the 5' end.

30

(Seq ID No. 65) 5'-3': AAT TC GCT GTT TGT GTG CTG CCT CTG AAG TCC ACA CTG AAC AAA
CTT CAG CCT ACT CAT GTC CCT AAA ATG GGC AAA CAT TGC AAG CAG C

The second oligonucleotide is the sense strand, spanning the second half of the Apo E enhancer and contains the Eco RI restriction site at the 3' end.

5

(Seq ID No. 66) 5'-3': AAA CAG CAA ACA CAC AGC CCT CCC TGC CTG CTG ACC TTG GAG CTG
GGG CAG AGG TCA GAG ACC TCT CTG G

The third oligonucleotide is the antisense strand of the second half of the Apo E enhancer and contains the Eco RI site at the 5' end.

10

(Seq ID No. 67) 5'-3': AA TTC CAG AGA GGT CTC TGA CCT CTG CCC CAG CTC CAA GGT CAG
CAG GCA GGG AGG GCT GTG TGT TTG CTG TTT GCT GCT TG

The fourth oligonucleotide is the antisense strand of the first half of the Apo E enhancer and contains the Eco RI site at the 3' end.

15

(Seq ID No. 68) 5'-3': CAA TGT TTG CCC ATT TTA GGG ACA TGA GTA GGC TGA AGT TTG TTC
AGT GTG GAC TTC AGA GGC AGC ACA CAA ACA GC G

In a second reaction, this set of four oligonucleotides of approximately 80 bases in length are annealed, and ligated into the dephosphorylated Eco RI sites of pBluescript

20 KSII+ (Stratagene, La Jolla, California). Recombinants are selected and sequenced.

Correct monomer Apo E enhancer elements are excised by Eco RI, electrophoresed through a polyacrylamide gel and the DNA is purified following standard procedures. The

monomer enhancer element is ligated into the dephosphorylated Eco RI sites of pBluescript KSII+ at an insert:vector ratio of 10:1 in order to generate a second set of recombinants

25 containing dimers. Recombinant plasmids were assessed by DNA sequencing to select clones which carried both enhancer fragments in the same orientation with the 5' end closer to the Pst I site and the 3' end closer to the Eco RV site. This construct is designated KSII+ Eco RI ApoE(x2).

The 300 base pair Pst I/Eco RV fragment from KSII+ Eco RI ApoE(x2) is inserted into the Pst I and Eco RV sites within the multi-cloning sites of KSII+ Hind III ApoE(x2). This construct is designated KSII+ ApoE(x4), and contains 4 copies of the 154 base pair Apo E enhancer.

5 b. Human alpha 1-antitrypsin promoter

The human alpha 1-antitrypsin (hAAT) promoter has been identified as a 400 base pair region by Shen *et al.*, 1989, *DNA* 8:101-108. The hAAT promoter DNA is obtained from a genomic library as was done by Long *et al.*, 1984, *Biochem* 23:4828-4837, or can be obtained from ATCC (ATCC No. 61596).

- 10 A PCR reaction is conducted in order to isolate the hAAT promoter. The 5' end of the primer contains the Sma I restriction site.

(Seq ID No. 69) 5'-3': CGCG CCG CCC GGG GTA GAT CTT GCT ACC AGT GG

- 15 The second primer corresponding to the anti-sense nucleotide sequence contains the Not I restriction site.

(Seq ID No. 70) 5'-3': GCG CCC GCG GCC GC CAC TGT CCC AGG TCA GTG GTG GTG CC

- 20 This PCR product, approximately 400 bp in length, is digested with Sma I and Not I and is cloned into the Sma I and Not I sites within the multi-cloning sites of KSII+ ApoE(x4). This construct is confirmed by sequencing and is designated KSII+ ApoE/hAAT.

c. Construction of the retroviral vector pBA6 ApoE/hAAT-IFN α 2a and pBA6 ApoE/hAAT-IFN α 2b

1. Interferon- α 2a and α 2b

The sense primer for both interferon- α 2a and α 2b is from the 5' region of the coding sequence, 6 bp upstream from the ATG start codon until 25 bp downstream from the start codon. The 5' end of the primer contains the Not I restriction site.

(Seq ID No. 71) 5'-3': CGCG CCG GCGG CCGC TCT ACA ATG GCC TTG ACC TTT GCT TTA CTG G

10 For interferon- α 2a and α 2b, the second primer corresponds to the anti-sense nucleotide sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Cla I restriction site.

(Seq ID No. 72) 5'-3': GCG CCC ATC GAT TCA TTC CTT ACT TCT TAA ACT TTC TTG CAA G

15 The product of this PCR reaction is digested with Not I and Cla I and purified. KSII+ ApoE/hAAT is digested with Not I and Cla I and the 1 kb fragment containing the Apo E enhancer and the hAAT promoter is isolated. From pBA6bL1, the 6.1 kb Xho/Cla I retroviral vector backbone is isolated. In a three part ligation (a) the Xho I/Not I fragment containing the Apo E enhancer and the hAAT promoter, (b) the Not I/ Cla I fragment encoding the IFN α 2a or α 2b gene and (c) the Xho I/Cla I fragment encoding the retroviral vector backbone are joined to generate pBA6 ApoE/hAAT-IFN α 2a or α 2b.

25 7. Construction of the Marker-less retroviral vector pBA9 ApoE/hAAT-IFN α 2a and

pBA9 ApoE/hAAT-IFN α 2b

a. Interferon- α 2a and α 2b

The sense primer for both interferon- α 2a and α 2b is from the 5' region of the coding sequence, 6 bp upstream from the ATG start codon until 25 bp downstream from the start codon. The 5' end of the primer contains the Not I restriction site.

30

(Seq ID No. 34) 5'-3': CGCG CCG GCGG CCGC TCT ACA ATG GCC TTG ACC TTT GCT TTA CTG G

For interferon- α 2a and α 2b, the second primer corresponds to the anti-sense nucleotide sequence, extending 31 bp into the coding sequence from the 3' end of the interferon gene. The 5' end of the primer contains the Cla I restriction site.

5

(Seq ID No. 35) 5'-3': GCG CCC ATC GAT TCA TTC CTT ACT TCT TAA ACT TTC TTG CAA G

The product of this PCR reaction is digested with Not I and Cla I and purified. KSII+ ApoE/hAAT is digested with Not I and Cla I and the 1 kb fragment containing the Apo E enhancer and the hAAT promoter is isolated. From pBA9b, the 4.8 kb Xho/Cla I retroviral vector backbone is isolated. In a three part ligation (a) the Xho I/Not I fragment containing the Apo E enhancer and the hAAT promoter, (b) the Not I/ Cla I fragment encoding the IFN α 2a or α 2b gene and (c) the Xho I/Cla I fragment encoding the pBA9b retroviral vector backbone are joined to generate pBA9 ApoE/hAAT-IFN α 2a or α 2b.

15

8. Construction of the retroviral vector pBA6 TK/ApoE/hAAT-IFN α 2a and pBA6 TK/ApoE/hAAT-IFN α 2b

The thymidine kinase (TK) gene is excised from pTJ5b-TK with Xho/Cla I and inserted into the Xho I/Cla I sites of KSII+ ApoE/hAAT. This construct is designated KSII+ TK/ApoE/hAAT.

20

PCR products containing either the IFN α 2a or IFN α 2b genes described in the example above is digested with Not I and Cla I and purified. The Xho/Not fragment containing the TK/ApoE/hAAT sequence is excised from KSII+ TK/ApoE/hAAT. From pBA6bL1, the 6.1 kb Xho/Cla I retroviral vector backbone is isolated.

25

In a three part ligation (a) the Xho I/Not I fragment containing the TK gene and the Apo E enhancer with the hAAT promoter, (b) the Not I/ Cla I fragment encoding the IFN α 2a or α 2b gene and (c) the Xho I/Cla I fragment encoding the retroviral vector backbone are joined to generate pBA6 TK/ApoE/hAAT-IFN α 2a or pBA6 TK/ApoE/hAAT-IFN α 2b.

9. Construction of the Marker-less retroviral vector pBA9 TK/ApoE/hAAT-IFN
 α 2a
and pBA9 TK/ApoE/hAAT-IFN α 2b

5 The thymidine kinase (TK) gene is excised from pTJ5b-TK with Xho/Cla I and inserted into the Xho I/Cla I sites of KSII+ ApoE/hAAT. This construct is designated KSII+ TK/ApoE/hAAT.

 PCR products containing either the IFN α 2a or IFN α 2b genes described in the example above is digested with Not I and Cla I and purified. The Xho/Not fragment
10 containing the TK/ApoE/hAAT sequence is excised from KSII+ TK/ApoE/hAAT. From pBA9b, the 4.8 kb Xho/Cla I pBA9 retroviral vector backbone is isolated.

 In a three part ligation (a) the Xho I/Not I fragment containing the TK gene and the Apo E enhancer with the hAAT promoter, (b) the Not I/ Cla I fragment encoding the IFN
15 α 2a or α 2b gene and (c) the Xho I/Cla I fragment encoding the pBA9 retroviral vector backbone are joined to generate pBA9 TK/ApoE/hAAT-IFN α 2a or pBA9 TK/ApoE/hAAT-IFN α 2b.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Chiron Corporation

(ii) TITLE OF INVENTION: Methods for Administration of
Recombinant Gene Delivery Vehicles for Treatment of
Hemophilia

(iii) NUMBER OF SEQUENCES: 81

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Chiron Corporation
(B) STREET: 4560 Horton Street
(C) CITY: Emeryville
(D) STATE: California
(E) COUNTRY: U.S.A.
(F) ZIP: 94608

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: Unassigned
(B) FILING DATE: Even date herewith
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Kruse, Norman J.
(B) REGISTRATION NUMBER: 35,235
(C) REFERENCE/DOCKET NUMBER: 1155.200

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (510) 923-3520
(B) TELEFAX: (510) 655-3542

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAGAGATGGG GGAGGCTAAC TGAG

24

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GATCCTCAGT TAGCCTCCCC CATCTCTC

28

- (2) INFORMATION FOR SEQ ID NO:3:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:
TATATCTCCA GATGAGGTAC ATGATTTTAG GCTTG 35
- (2) INFORMATION FOR SEQ ID NO:4:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 40 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:
TATATATCGA TTCAAGGCAT TTTCTTTTCA TCAATAAAAC 40
- (2) INFORMATION FOR SEQ ID NO:5:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 37 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
TCGAGGATCC GCCC GGCGG CCGCATCGAT GTCGACG 37
- (2) INFORMATION FOR SEQ ID NO:6:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
CGCGTCGACA TCGATGCGGC CGCCGGGCG GATCC 35

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 77 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AGTGAATTCTG AGCTCGGTAC CCGGGGATCC TCTAGAGTCG ACCTGCAGGC ATGCAAGCTT
GGCGTAATCA TGGTCAT

60

77

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Ala Arg Glu Met Gly Glu Ala Asn
1 5

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CCCAGAGAT GGGGGAGGCT AACTGAG

27

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGGCTCTCTA CCCCTCCGA TTGACACCTA G

31

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Thr Ile Met Thr Met
1 5

(2) INFORMATION FOR SEQ ID NO:12:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 24 base pairs	
(B) TYPE: nucleic acid	
(C) STRANDEDNESS: single	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: DNA (genomic)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:	
CCCTGTGCCT TATTGAACT AACC	24
(2) INFORMATION FOR SEQ ID NO:13:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 24 base pairs	
(B) TYPE: nucleic acid	
(C) STRANDEDNESS: single	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: DNA (genomic)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:	
CCCACCACAA CCACATATCC CTCC	24
(2) INFORMATION FOR SEQ ID NO:14:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 19 base pairs	
(B) TYPE: nucleic acid	
(C) STRANDEDNESS: single	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: DNA (genomic)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:	
CCAGTCCTCC GATTGACTG	19
(2) INFORMATION FOR SEQ ID NO:15:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 8332 base pairs	
(B) TYPE: nucleic acid	
(C) STRANDEDNESS: single	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: DNA (genomic)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:	
GCGCCAGTCC TCCGATTGAC TGAGTCGCCC GGGTACCCGT GTATCCAATA AACCCCTCTG	60
CAGTTGCATC CGACTTGTGG TCTCGCTGTT CTTGGGAGG GTCTCCTCTG AGTGATTGAC	120
TACCCGTCAG CGGGGTCTT TCATTTGGG GCTCGTCCG GATCGGGAGA CCCCTGCCCA	180
GGGACCACCG ACCCACCACC GGGAGGTAAG CTGGCCAGCA ACTTATCTGT GTCTGTCCGA	240
TTGTCTAGTG TCTATGACTG ATTTTATGCG CCTGCGTCGG TACTAGTTAG CTAAC TAGCT	300
CTGTATCTGG CGGACCCGTG GTGGAAGTGA CGAGTTCGGA ACACCCGGCC GCAACCCTGG	360
GAGACGTCCC AGGGACTTCG GGGGCCGTTT TTGTGGCCCG ACCTGAGTCC AAAAATCCCG	420
ATCGTTTTTG ACTCTTTGGT GCACCCCTT TAGAGGAGG ATATGTGGT CTGGTAGGAG	480
ACGAGAACCT AAAACAGTTC CCGCCTCCGT CTGAATTTT GCTTTCGGT TGGGACCGAA	540
GCCGCGCCGC GCGTCTTGTC TGCTGCAGCA TCGTCTGTG TTGTCTCTGT CTGACTGTGT	600
TTCTGTATTT GTCTGAGAAT ATGGGCCAGA CTGTTACCAC TCCCTTAAGT TTGACCTTAG	660
GTCAGTGAA AGATGTCGAG CGGATCGCTC ACAACCAGTC GGTAGATGTC AAGAAGAGAC	720

GTTGGGTTAC	CTTCTGCTCT	GCAGAATGGC	CAACCTTTAA	CGTCGGATGG	CCGCGAGACG	780
GCACCTTTAA	CCGAGACCTC	ATCACCCAGG	TTAAGATCAA	GGTCTTTTCA	CCTGGCCCGC	840
ATGGACACCC	AGACCAGGTC	CCCTACATCG	TGACCTGGGA	AGCCTTGGCT	TTTGACCCCC	900
CTCCCTGGGT	CAAGCCCTTT	GTACACCCTA	AGCCTCCGCC	TCCTCTTCCT	CCATCCGCCC	960
CGTCTCTCCC	CCTTGAACTT	CCTCGTTCGA	CCCCGCCTCG	ATCCTCCCTT	TATCCAGCCC	1020
TCACTCCTTC	TCTAGGCGCC	AAACCTAAAC	CTCAAGTTCT	TTCTGACAGT	GGGGGGCCGC	1080
TCATCGACCT	ACTTACAGAA	GACCCCCCGC	CTTATAGGGA	CCCAAGACCA	CCCCCTCCG	1140
ACAGGGACGG	AAATGGTGGA	GAAGCGACCC	CTGCGGGAGA	GGCACC GGAC	CCCTCCCCAA	1200
TGGCATCTCG	CCTACGTGGG	AGACGGGAGC	CCCTGTGGC	CGACTCCACT	ACCTCGCAGG	1260
CATTCCCCCT	CCGCGCAGGA	GGAAACGGAC	AGCTTCAATA	CTGGCCGTTC	TCCTCTTCTG	1320
ACCTTTACAA	CTGGAAAAAT	AATAACCCTT	CTTTTCTGA	AGATCCAGGT	AAACTGACAG	1380
CTCTGATCGA	GTCTGTTCTC	ATCACCCATC	AGCCACCTG	GGACGACTGT	CAGCAGCTGT	1440
TGGGGACTCT	GCTGACCGGA	GAAGAAAAAC	AACGGGTGCT	CTTAGAGGCT	AGAAAGGCGG	1500
TGCGGGGCGA	TGATGGGCGC	CCCACTCAAC	TGCCCAATGA	AGTCGATGCC	GCTTTTCCCC	1560
TCGAGCGCCC	AGACTGGGAT	TACACCACCC	AGGCAGGTAG	GAACCACCTA	GTCCACTATC	1620
GCCAGTTGCT	CCTAGCGGGT	CTCCAAAACG	CGGGCAGAAG	CCCCACCAAT	TGGCCAAGG	1680
TAAAAGGAAT	AACACAAGGG	CCCAATGAGT	CTCCCTCGGC	CTTCCTAGAG	AGACTTAAGG	1740
AAGCCTATCG	CAGGTACACT	CCTTATGACC	CTGAGGACCC	AGGGCAAGAA	ACTAATGTGT	1800
CTATGTCTTT	CATTTGGCAG	TCTGCCCCAG	ACATTGGGAG	AAAGTTAGAG	AGGTTAGAAG	1860
ATTTAAAAAA	CAAGACGCTT	GGAGATTGG	TTAGAGAGGC	AGAAAAGATC	TTAATAAAC	1920
GAGAAACCCC	GGAAGAAAGA	GAGGAACGTA	TCAGGAGAGA	AACAGAGGAA	AAAGAAGAAC	1980
GCCGTAGGAC	AGAGGATGAG	CAGAAAGAGA	AAGAAAGAGA	TCGTAGGAGA	CATAGAGAGA	2040
TGAGCAAGCT	ATTGGCCACT	GTCGTTAGTG	GACAGAAACA	GGATAGACAG	GGAGGAGAAC	2100
GAAGGAGGTC	CCAACTCGAT	CGCGACCAGT	GTGCCTACTG	CAAAGAAAAG	GGGCACTGGG	2160
CTAAAGATTG	TCCCAAGAAA	CCACGAGGAC	CTCGGGGACC	AAGACCCAG	ACCTCCCTCC	2220
TGACCCTAGA	TGACTAGGGA	GGTCAGGGTC	AGGAGCCCCC	CCCTGAACCC	AGGATAACCC	2280
TCAAAGTCGG	GGGGCAACCC	GTACACCTTC	TGGTAGATAC	TGGGGCCCAA	CACTCCGTGC	2340
TGACCCAAAA	TCCTGGACCC	CTAAGTGATA	AGTCTGCCTG	GGTCCAAGGG	GCTACTGGAG	2400
GAAAGCGGTA	TCGCTGGACC	ACGGATCGCA	AAGTACATCT	AGCTACCGGT	AAGGTCACCC	2460
ACTCTTTCCT	CCATGTACCA	GACTGTCCCT	ATCCTCTGTT	AGGAAGAGAT	TTGCTGACTA	2520
AACTAAAAGC	CCAAATCCAC	TTTGAGGGAT	CAGGAGCTCA	GGTTATGGGA	CCAATGGGGC	2580
AGCCCCTGCA	AGTGTTGACC	CTAAATATAG	AAGATGAGCA	TCGGCTACAT	GAGACCTCAA	2640

AAGAGCCAGA TGTTCCTCTA GGTCCACAT GGCTGTCTGA TTTTCCTCAG GCCTGGGCGG	2700
AAACCGGGGG CATGGGACTG GCAGTTCGCC AAGCTCCTCT GATCATACCT CTGAAAGCAA	2760
CCTCTACCCC CGTGTCCATA AAACAATACC CCATGTCACA AGAAGCCAGA CTGGGGATCA	2820
AGCCCCACAT ACAGAGACTG TTGGACCAGG GAATACTGGT ACCCTGCCAG TCCCCCTGGA	2880
ACACGCCCTT GCTACCCGTT AAGAAACCAG GGACTAATGA TTATAGGCCT GTCCAGGATC	2940
TGAGAGAAGT CAACAAGCGG GTGGAAGACA TCCACCCAC CGTGCCCAAC CCTTACAACC	3000
TCTTGAGCGG GCTCCACCG TCCCACCAGT GGTACACTGT GCTTGATTTA AAGGATGCCT	3060
TTTTCTGCCT GAGACTCCAC CCCACCAGTC AGCCTCTCTT CGCCTTTGAG TGGAGAGATC	3120
CAGAGATGGG AATCTCAGGA CAATTGACCT GGACCAGACT CCCACAGGGT TTCAAAAACA	3180
GTCCACCCCT GTTTGATGAG GCACTGCACA GAGACCTAGC AGACTTCCGG ATCCAGCACC	3240
CAGACTTGAT CCTGCTACAG TACGTGGATG ACTTACTGCT GGCCGCCACT TCTGAGCTAG	3300
ACTGCCAACA AGGTACTCGG GCCCTGTTAC AAACCTAGG GAACCTCGGG TATCGGGCCT	3360
CGGCCAAGAA AGCCCAAATT TGCCAGAAAC AGGTCAAGTA TCTGGGGTAT CTTCTAAAAG	3420
AGGGTCAGAG ATGGCTGACT GAGGCCAGAA AAGAGACTGT GATGGGGCAG CCTACTCCGA	3480
AGACCCCTCG ACAACTAAGG GAGTTCCTAG GGACGGCAGG CTTCTGTGCG CTCTGGATCC	3540
CTGGGTTTGC AGAAATGGCA GCCCCCTTGT ACCCTCTCAC CAAAACGGGG ACTCTGTTTA	3600
ATTGGGGCCC AGACCAACAA AAGGCCTATC AAGAAATCAA GCAAGCTCTT CTAAGTCCC	3660
CAGCCCTGGG GTTGCCAGAT TTGACTAAGC CCTTTGAACT CTTTGTGAC GAGAAGCAGG	3720
GCTACGCCAA AGGTGTCCTA ACGCAAAAAC TGGGACCTTG GCGTCGGCCG GTGGCCTACC	3780
TGTCCAAAAA GCTAGACCCA GTAGCAGCTG GGTGGCCCCC TTGCTACGG ATGGTAGCAG	3840
CCATTGCCGT ACTGACAAAG GATGCAGGCA AGCTAACCAT GGGACAGCCA CTAGTCATTC	3900
TGGCCCCCA TGCACTAGAG GCACTAGTCA AACAACCCCC CGACCGCTGG CTTTCCAACG	3960
CCCGGATGAC TCACTATCAG GCCTTGCTTT TGGACACGGA CCGGGTCCAG TTCGGACCGG	4020
TGGTAGCCCT GAACCCGGCT ACGCTGCTCC CACTGCCTGA GGAAGGGCTG CAACACAAC	4080
GCCTTGATAT CCTGGCCGAA GCCCACGGAA CCCGACCCGA CCTAACGGAC CAGCCGCTCC	4140
CAGACGCCGA CCACACCTGG TACACGGATG GAAGCAGTCT CTTACAAGAG GGACAGCGTA	4200
AGGCGGGAGC TGCGGTGACC ACCGAGACCG AGGTAATCTG GGCTAAAGCC CTGCCAGCCG	4260
GGACATCCGC TCAGCGGGCT GAACTGATAG CACTCACCCA GGCCCTAAAG ATGGCAGAAG	4320
GTAAGAAGCT AAATGTTTAT ACTGATAGCC GTTATGCTTT TGCTACTGCC CATATCCATG	4380
GAGAAATATA CAGAAGGCGT GGGTTGCTCA CATCAGAAG CAAAGAGATC AAAAATAAAG	4440
ACGAGATCTT GGCCCTACTA AAAGCCCTCT TTCTGCCCAA AAGACTTAGC ATAATCCATT	4500
GTCCAGGACA TCAAAGGGA CACAGCGCCG AGGCTAGAGG CAACCGGATG GCTGACCAAG	4560

CGGCCCCGAAA GGCAGCCATC ACAGAGACTC CAGACACCTC TACCCCTCCTC ATAGAAAATT 4620
CATCACCCCTA CACCTCAGAA CATTTTCATT ACACAGTGAC TGATATAAAG GACCTAACCA 4680
AGTTGGGGGC CATTTATGAT AAAACAAAGA AGTATTGGGT CTACCAAGGA AAACCTGTGA 4740
TGCTGACCA GTTTACTTTT GAATTATTAG ACTTTCTTCA TCAGCTGACT CACCTCAGCT 4800
TCTCAAAAT GAAGGCTCTC CTAGAGAGAA GCCACAGTCC CTACTACATG CTGAACCGGG 4860
ATCGAACACT CAAAAATATC ACTGAGACCT GCAAAGCTTG TGCACAAGTC AACGCCAGCA 4920
AGTCTGCCGT TAAACAGGGA ACTAGGGTCC GCGGGCATCG GCGCGGCACT CATTGGGAGA 4980
TCGATTTCAC CGAGATAAAG CCCGGATTGT ATGGCTATAA ATATCTTCTA GTTTTTATAG 5040
ATACCTTTTC TGGCTGGATA GAAGCCTTCC CAACCAAGAA AGAAACCGCC AAGGTCGTAA 5100
CCAAGAAGCT ACTAGAGGAG ATCTTCCCCA GGTTCGGCAT GCCTCAGGTA TTGGGAAGTG 5160
ACAATGGGCC TGCCCTTCGTC TCCAAGGTGA GTCAGACAGT GGCCGATCTG TTGGGGATTG 5220
ATTGGAAT ACATTGTGCA TACAGACCCC AAAGCTCAGG CCAGGTAGAA AGAATGAATA 5280
GAACCATCAA GGAGACTTTA ACTAAATTAA CGCTTGCAAC TGGCTCTAGA GACTGGGTGC 5340
TCCTACTCCC CTTAGCCCTG TACCGAGCCC GCAACACGCC GGGCCCCCAT GGCCTCACCC 5400
CATATGAGAT CTTATATGGG GCACCCCCGC CCCTTGTAAG CTTCCTGAC CCTGACATGA 5460
CAAGAGTTAC TAACAGCCCC TCTCTCCAAG CTCACTTACA GGCTCTCTAC TTAGTCCAGC 5520
ACGAAGTCTG GAGACCTCTG GCGGCAGCCT ACCAAGAACA ACTGGACCGA CCGGTGGTAC 5580
CTCACCTTA CCGAGTCGGC GACACAGTGT GGGTCCGCCG ACACCAGACT AAGAACCTAG 5640
AACCTCGCTG GAAAGGACCT TACACAGTCC TGCTGACCAC CCCCACCGCC CTCAAAGTAG 5700
ACGGCATCGC AGCTTGATA CACGCCGCC ACGTGAAGGC TGCCGACCCC GGGGTGGAC 5760
CATCCTCTAG ACTGACATGG CGCGTTCAAC GCTCTCAAAA CCCCTTAAAA ATAAGGTAA 5820
CCCGCAGGC CCCCTAATCC CCTTAATTCT TCTGATGCTC AGAGGGGTCA GTACTGCTTC 5880
GCGCGCTCC AGTCCTCATC AAGTCTATAA TATCACCTGG GAGGTAACCA ATGGAGATCG 5940
GGAGACGGTA TGGGCAACTT CTGGCAACCA CCCTCTGTGG ACCTGGTGGC CTGACCTTAC 6000
CCCAGATTTA TGTATGTTAG CCCACCATGG ACCATCTTAT TGGGGGCTAG AATATCAATC 6060
CCCTTTTCT TCTCCCCCGG GGCCCCCTTG TTGCTCAGG GGCAGCAGCC CAGGCTGTTC 6120
CAGAGACTGC GAAGAACCTT TAACCTCCCT CACCCCTCGG TGCAAACTG CCTGGAACAG 6180
ACTCAAGCTA GACCAGACAA CTCATAAATC AAATGAGGGA TTTTATGTTT GCGCGGGCC 6240
CCACCGCCCC CGAGAATCCA AGTCATGTGG GGGTCCAGAC TCCTTCTACT GTGCCTATTG 6300
GGGCTGTGAG ACAACCGGTA GAGCTTACTG GAAGCCCTCC TCATCATGGG ATTCATCAC 6360
AGTAAACAAC AATCTCACCT CTGACCAGGC TGTCCAGGTA TGCAAAGATA ATAAGTGGTG 6420
CAACCCCTTA GTTATTCGGT TTACAGACGC CGGGAGACGG GTTACTTCCT GGACCACAGG 6480

ACATTACTGG GGCTTACGTT TGTATGTCTC CGGACAAGAT CCAGGGCTTA CATTTGGGAT	6540
CCGACTCAGA TACCAAAATC TAGGACCCCG CGTCCCAATA GGGCCAAACC CCGTTCTGGC	6600
AGACCAACAG CCACTCTCCA AGCCCAAACC TGTTAAGTCG CCTTCAGTCA CCAAACCACC	6660
CAGTGGGACT CCTCTCTCCC CTACCCAACT TCCACCGGCG GGAACGGAAA ATAGGCTGCT	6720
AAACTTAGTA GACGGAGCCT ACCAAGCCCT CAACCTCACC AGTCCTGACA AAACCCAAGA	6780
GTGCTGGTTG TGTCTAGTAG CGGGACCCCG CTACTACGAA GGGGTTGCCG TCCTGGGTAC	6840
CTACTCCAAC CATACTCTG CTCCAGCCAA CTGCTCCGTG GCCTCCCAAC ACAAGTTGAC	6900
CCTGTCCGAA GTGACCGGAC AGGGACTCTG CATAGGAGCA GTTCCCAAAA CACATCAGGC	6960
CCTATGTAAT ACCACCCAGA CAAGCAGTCG AGGGTCCCTAT TATCTAGTTG CCCCTACAGG	7020
TACCATGTGG GCTTGTAGTA CCGGGCTTAC TCCATGCATC TCCACCACCA TACTGAACCT	7080
TACCACTGAT TATTGTGTTT TTGTCGAACT CTGGCCAAGA GTCACCTATC ATTCCCCCAG	7140
CTATGTTTAC GGCTGTGTTG AGAGATCCAA CCGACACAAA AGAGAACCGG TGTCGTTAAC	7200
CCTGGCCCTA TTATTGGGTG GACTAACCAT GGGGGGAATT GCCGCTGGAA TAGGAACAGG	7260
GACTACTGCT CTAATGGCCA CTCAGCAATT CCAGCAGCTC CAAGCCGAG TACAGGATGA	7320
TCTCAGGGAG GTTGAAAAAT CAATCTCTAA CCTAGAAAAG TCTCTCACTT CCCTGTCTGA	7380
AGTTGTCCTA CAGAATCGAA GGGGCTAGA CTTGTTATTT CTAAAAGAAG GAGGGCTGTG	7440
TGCTGCTCTA AAAGAAGAAT GTTGCTTCTA TGCGGACCAC ACAGGACTAG TGAGAGACAG	7500
CATGGCCAAA TTGAGAGAGA GGCTTAATCA GAGACAGAAA CTGTTTGAGT CAACTCAAGG	7560
ATGGTTTGAG GGACTGTTTA ACAGATCCCC TTGGTTTACC ACCTTGATAT CTACCATTAT	7620
GGGACCCCTC ATTGTACTCC TAATGATTTT GCTCTTCGGA CCCTGCATTC TTAATCGATT	7680
AGTCCAATTT GTTAAAGACA GGATATCAGT GGTCCAGGCT CTAGTTTGA CTCAACAATA	7740
TCACCAGCTG AAGCCTATAG AGTACGAGCC ATAGATAAAA TAAAAGATTT TATTTAGTCT	7800
CCAGAAAAAG GGGGGAATGA AAGACCCAC CTGTAGGTTT GGCAAGCTAG CTTAAGTAAC	7860
GCCATTTTGC AAGGCATGGA AAAATACATA ACTGAGAATA GAGAAGTTCA GATCAAGGTC	7920
AGGAACAGAT GGAACAGCTG AATATGGGCC AAACAGGATA TCTGTGGTAA GCAGTTCTG	7980
CCCCGGCTCA GGGCCAAGAA CAGATGGAAC AGCTGAATAT GGGCCAAACA GGATATCTGT	8040
GGTAAGCAGT TCCTGCCCGG GCTCAGGGCC AAGAACAGAT GGTCCCCAGA TGCGGTCCAG	8100
CCCTCAGCAG TTTCTAGAGA ACCATCAGAT GTTCCAGGG TGCCCCAAGG ACCTGAAATG	8160
ACCCTGTGCC TTATTGTAAC TAACCAATCA GTTCGCTTCT CGCTTCTGTT CGCGCGCTTC	8220
TGCTCCCCGA GCTCAATAAA AGAGCCCACA ACCCCTCACT CGGGGCGCCA GTCCTCCGAT	8280
TGACTGAGTC GCCCGGTAC CCGTGTATCC AATAAACCT CTTGCAGTTG CA	8332

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:
GGGAGTGGTA ACAGTCTGGC CTTAATTCTC AG 32
- (2) INFORMATION FOR SEQ ID NO:17:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:
CGGTCGACCT CGAGAATTAA TTC 23
- (2) INFORMATION FOR SEQ ID NO:18:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:
CTGGGAGACG TCCCAGGGAC TTC 23
- (2) INFORMATION FOR SEQ ID NO:19:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:
GGCCAGACTG TTACCACTCC CTGAAGTTTG AC 32
- (2) INFORMATION FOR SEQ ID NO:20:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
CATCGATAAA ATAAAAGATT TTATTAGTC 30
- (2) INFORMATION FOR SEQ ID NO:21:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:
CAAATGAAAG ACCCCGCTG AC 22
- (2) INFORMATION FOR SEQ ID NO:22:
(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:
GAAGCTTCTC CCAGAACCCA CCAGTCTTGA AACGCCATC 39
- (2) INFORMATION FOR SEQ ID NO:23:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:
GTACCAGCTT TTGGTCTCAT CAAAG 25
- (2) INFORMATION FOR SEQ ID NO:24:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 36 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:
CTCCTCGAGC TAAAGATATT TTAGAGAAGA ATTAAC 36
- (2) INFORMATION FOR SEQ ID NO:25:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:
TTCCTCTGGA CAGCTGTCTA CTTTG 25
- (2) INFORMATION FOR SEQ ID NO:26:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 51 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:
TCGAGGGGCC CAGATCTGCG GCCGCTCGCG AGTCGACAAG CTTGGATCCA T 51
- (2) INFORMATION FOR SEQ ID NO:27:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 49 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:
CGATGGATCC AAGCTTGTCG ACTCGCGAGC GGCCGCAGAT CTGGGCCCC 49
- (2) INFORMATION FOR SEQ ID NO:28:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:
CACCGTCGTC GACTTATGCT 20
- (2) INFORMATION FOR SEQ ID NO:29:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:
GACCGTCGAC TCAATTCTGG GAGAAGCTTC TTGG 34
- (2) INFORMATION FOR SEQ ID NO:30:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:
CACCGTCGTC GACTTATGCT 20
- (2) INFORMATION FOR SEQ ID NO:31:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:
CAACGCTCGA GAAGCAGAAT CGCAAAGGC 30
- (2) INFORMATION FOR SEQ ID NO:32:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 32 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: DNA (genomic)
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:
TCGGCTCGAG GCATCAACGG GAAATAACTC GT 32

- (2) INFORMATION FOR SEQ ID NO:33:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:
CCGACTCGAG TCAGTAGAGG TCCTGTGCCT C 31
- (2) INFORMATION FOR SEQ ID NO:34:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 46 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:
CGCGCCGGCG GCCGCTCTAC AATGGCCTTG ACCTTGCTT TACTGG 46
- (2) INFORMATION FOR SEQ ID NO:35:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 43 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:
CGGCCCATCG ATTCATTCCT TACTTCTTAA ACTTTCTTGC AAG 43
- (2) INFORMATION FOR SEQ ID NO:36:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 44 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:
CGCGCCGCTC GAGTCTACAA TGGCCTTGAC CTTTGCTTTA CTGG 44
- (2) INFORMATION FOR SEQ ID NO:37:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 43 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:
GCGGCCATCG ATTCATTCCT TACTTCTTAA ACTTTCTTGC AAG 43
- (2) INFORMATION FOR SEQ ID NO:38:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 47 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:
CGCGCCGCTC GAGCATCCAA TGGCCCTGTC CTTTCTTTA CTTATGG 47

- (2) INFORMATION FOR SEQ ID NO:39:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 39 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:
CCATCGATTC AATCCTTCCT CCTTAATCTT TTTTGCAAG 39
- (2) INFORMATION FOR SEQ ID NO:40:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 39 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:
CCGGATCCTC TACAATGGCC TTGACCTTG CTTTACTGG 39
- (2) INFORMATION FOR SEQ ID NO:41:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 46 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:
CGCGCCGGCG GCCGCTCATT CTTTACTTCT TAACTTTCT TGCAAG 46
- (2) INFORMATION FOR SEQ ID NO:42:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:
CCGGATCCCA TCCAATGGCC CTGTCCTTT CTTTACTTAT GG 42
- (2) INFORMATION FOR SEQ ID NO:43:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 46 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:
CGCGCCGGCG GCCGCTCAAT CCTTCCTCCT TAATCTTTT TGCAAG 46

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9080 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

GGGGGGGGGG	GGGGGGGGGG	GGGTGAGCAC	ATCCAGTGGG	TAAAGTTCCT	TAAAATGCTC	60
TGCAAAGAAA	TTGGGACTTT	TCATTAAATC	AGAAATTTTA	CTTTTTTCCC	CTCCTGGGAG	120
CTAAAGATAT	TTTAGAGAAG	AATTAACCTT	TTGCTTCTCC	AGTTGAACAT	TTGTAGCAAT	180
AAGTCATGCA	AATAGAGCTC	TCCACCTGCT	TCTTTCTGTG	CCTTTTGCGA	TTCTGCTTTA	240
GTGCCACCAG	AAGATACTAC	CTGGGTGCAG	TGGAACGTGC	ATGGGACTAT	ATGCAAAGTG	300
ATCTCCGTGA	GCTGCCCTGTG	GACGCAAGAT	TTCTCCTAG	AGTGCCAAAA	TCTTTTCCAT	360
TCAACACCTC	AGTCGTGTAC	AAAAAGACTC	TGTTTGTAGA	ATTACGGGAT	CACCTTTTCA	420
ACATCGCTAA	GCCAAGGCCA	CCCTGGATGG	GTCTGCTAGG	TCCTACCATC	CAGGCTGAGG	480
TTTATGATA	AGTGGTCATT	ACACTTAAGA	ACATGGCTTC	CCATCCTGTC	AGTCTTCATG	540
CTGTTGGTGT	ATCCTACTGG	AAAGCTTCTG	AGGGAGCTGA	ATATGATGAT	CAGACCGATC	600
AAAGGGAGAA	AGAAGATGAT	AAAGTCTTCC	CTGGTGGGAA	CCATACATAT	GTCTGGCAGG	660
TCCTGAAAGA	GAATGGTCCA	ATGGCCTCTG	ACCCACTGTG	CCTTACCTAC	TCATATCTTT	720
CTCATGTGGA	CCTGGTAAAA	GACTTGAATT	CAGGCCCTCAT	TGGAGCCCTA	CTAGTATGTA	780
GAGAAGGGAG	TCTGGCCAAG	GAAAAGACAC	AGACCTTGCA	CAAAATTATA	CTACTTTTTG	840
CTGTATTGTA	TGAAGGGAAA	AGTTGGCACT	CAGAAACAAA	GAACCTCCTG	ATGCAGGATA	900
GGGATGCTGC	ATCTGCTCGG	GCCTGGCCTA	AAATGCACAC	AGTCAATGGT	TATGTAAACA	960
GGTCTCTGCC	AGGTCTGATT	GGATGCCACA	GGAAATCAGT	CTATTGGCAT	GTGATTGGAA	1020
TGGGCACCAC	TCCTGAAGTG	CACTCAATAT	TCCTCGAAGG	TCACACATTT	CTTGTGAGGA	1080
ACCATCGCCA	GGCGTCCTTG	GAAATCTCGC	CAATAACTTT	CCTTACTGCT	CAAACACTCT	1140
TGATGGACCT	TGGACAGTTT	CTACTGTTTT	GTCAATATCTC	TTCCACCAA	CATGATGGCA	1200
TGGAAGCTTA	TGTCAAAGTA	GACAGCTGTC	CAGAGGAACC	CCAACACGA	ATGAAAAATA	1260
ATGAAGAAGC	GGAAGACTAT	GATGATGATC	TTACTGATTC	TGAAATGGAT	GTGGTCAGGT	1320
TTGATGATGA	CAACTCTCCT	TCCTTTATCC	AAATTCGCTC	AGTTGCCAAG	AAGCATCCTA	1380
AACTTTGGGT	ACATTACATT	GCTGCTGAAG	AGGAGGACTG	GGACTATGCT	CCCTAGTCC	1440
TCGCCCCCGA	TGACAGAAGT	TATAAAAGTC	AATATTTGAA	CAATGGCCCT	CAGCGGATTG	1500
GTAGGAAGTA	CAAAAAAGTC	CGATTTATGG	CATACACAGA	TGAAACCTTT	AAGACTCGTG	1560
AAGCTATTCA	GCATGAATCA	GGAATCTTGG	GACCTTTACT	TTATGGGGAA	GTGGAGACA	1620
CACCTGTTAG	TATATTTAAG	AATCAAGCAA	GCAGACCATA	TAACATCTAC	CCTCACGGAA	1680
TCACTGATGT	CCGTCCCTTG	TATTCAAGGA	GATTACCAAA	AGGTGTAAAA	CATTTGAAGG	1740
ATTTTCCAAT	TCTGCCAGGA	GAAATATTCA	AATATAAATG	GACAGTGACT	GTAGAAGATG	1800
GGCCAACTAA	ATCAGATCCT	CGGTGCCTGA	CCCGCTATTA	CTCTAGTTTC	GTAAATATGG	1860
AGAGAGATCT	AGCTTCAGGA	CTCATTGGCC	CTCTCCTCAT	CTGTACAAA	GAATCTGTAG	1920
ATCAAAGAGG	AAACCAGATA	ATGTCAGACA	AGAGGAATGT	CATCCTGTTT	TCTGTATTTG	1980
ATGAGAAACG	AAGCTGGTAC	CTCACAGAGA	ATATACAACG	CTTTCTCCCC	AATCCACTGT	2040
GAGTGCAGCT	TGAGGATCCA	GAGTTCCAAG	CCTCCAACAT	CATGCACAGC	ATCAATGGCT	2100
ATGTTTTTGA	TAGTTTGCAG	TTGTCAAGTT	GTTTGCATGA	GGTGGCATA	TGGTACATTC	2160
TAAGCATTTG	AGCACAGACT	GACTTCCTTT	CTGTCTTCTT	CTCTGGATAT	ACCTTCAAAC	2220
ACAAAATGGT	CTATGAAGAC	ACACTCACCC	TATTCCCAT	CTCAGGAGAA	ACTGTCTTCA	2280
TGTCGATGGA	AAACCCAGGT	CTATGGATTC	TGGGGTGCCA	CAACTCAGAC	TTTCGGAACA	2340
GAGGCATGAC	CGCCTTACTG	AAGGTTTCTA	GTTGTGACAA	GAACACTGGT	GATTATTACG	2400
AGGACAGTTA	TGAAGATATT	TCAGCATACT	TGCTGAGTAA	AAACAATGCC	ATTGAACCAA	2460
GAAGCTTCTC	CCAGAATTCA	AGACACCCTA	GCACCTAGGCA	AAAGCAATTT	AATGCCACCA	2520
CAATTCCAGA	AAATGACATA	GAGAAGACTG	ACCCTTGTTT	TGCACACAGA	ACACCTATGC	2580
CTAAAATACA	AAATGCTCTC	TCTAGTGATT	TGTTGATGCT	CTTGCAGACG	AGTCCTACTC	2640
CACATGGGCT	ATCCTTATCT	GATCTCCAAG	AAGCCAAATA	TGAGACTTTT	TCTGATGATC	2700
CATCACCTGG	AGCAATAGAC	AGTAATAACA	GCCTGTCTGA	AATGACACAC	TTCAGGCCAC	2760
AGCTCCATCA	CAGTGGGGAC	ATGGTATTTA	CCCCTGAGTC	AGGCCTCCAA	TTAAGATTAA	2820
ATGAGAAACT	GGGGACAAC	GCAGCAACAG	AGTTGAAGAA	ACTTGATTTT	AAAGTTTCTA	2880
GTACATCAAA	TAATCTGATT	TCAACAATTC	CATCAGACAA	TTTGGCAGCA	GGTACTGATA	2940
ATACAAGTTC	CTTAGGACCC	CCAAGTATGC	CAGTTCATTA	TGATAGTCAA	TTAGATACCA	3000
CTCTATTTGG	CAAAAAGTCA	TCTCCCTTTA	CTGAGTCTGG	TGGACCTCTG	AGCTTGAGTG	3060
AAGAAAATAA	TGATTCAAAG	TTGTTAGAAT	CAGGTTTAAT	GAATAGCCAA	GAAAGTTTAT	3120
GGGGAAAAAA	TGTATCGTCA	ACAGAGAGTG	GTAGTTTATT	TAAAGGGAAA	AGAGCTCATG	3180

GACCTGCTTT	GTTGACTAAA	GATAATGCCT	TATTCAAAGT	TAGCATCTCT	TTGTAAAGA	3240
CAAACAAAAC	TTCCAATAAT	TCAGCAACTA	ATAGAAAGAC	TCACATTGAT	GGCCCATCAT	3300
TATTAATTGA	GAATAGTCCA	TCAGTCTGGC	AAAATATATT	AGAAAGTGAC	ACTGAGTTTA	3360
AAAAAGTGAC	ACCTTTGATT	CATGACAGAA	TGCTTATGGA	CAAAAATGCT	ACAGCTTTGA	3420
GGCTAAATCA	TATGTCAAAT	AAAACACTTT	CATCAAAAAA	CATGGAAATG	GTCCAACAGA	3480
AAAAAGAGGG	CCCCATTCCA	CCAGATGCAC	AAAATCCAGA	TATGTCGTTT	TTTAAGATGC	3540
TATCTTGGCC	AGAATCAGCA	AGGTGGATAC	AAAGGACTCA	TGGAAAGAAC	TCTCTGAACT	3600
CTGGGCAAGG	CCCCAGTCCA	AAGCAATTAG	TATCCTTAGG	ACCAGAAAAA	TCTGTGGAAG	3660
GTCAGAATTT	CTTGCTGAG	AAAAACAAAG	TGGTAGTAGG	AAAGGGTGAA	TTTACAAAGG	3720
ACGTAGGACT	CAAAGAGATG	GTTTTTCCAA	GCAGCAGAAA	CCTATTTCTT	ACTAACTTGG	3780
ATAATTTACA	TGAAAATAAT	ACACACAATC	AAGAAAAAAA	AATTCAGGAA	GAAATAGAAA	3840
AGAAGGAAAC	ATTAATCCAA	GAGAATGTAG	TTTTGCCTCA	GATACATACA	GTGACTGGCA	3900
CTAAGAATTT	CATGAAGAAC	CTTTTCTTAC	TGAGCACTAG	GCAAAATGTA	GAAGGTTTAT	3960
ATGACGGGGC	ATATGCTCCA	GTACTTCAAG	ATTTTAGGTC	ATTAAATGAT	TCAACAAATA	4020
GAACAAAGAA	ACACACAGCT	CATTTCTCAA	AAAAAGGGGA	GGAAGAAAAC	TTGGAAGGCT	4080
TGGGAAATCA	AACCAAGCAA	ATTGTAGAGA	AATATGCATG	CACCACAAGG	ATATCTCCTA	4140
ATACAAGCCA	GCAGAATTTT	GTCAACGCAAC	GTAGTAAGAG	AGCTTTGAAA	CAATTCAGAC	4200
TCCCACTAGA	AGAAACAGAA	CTTGAAAAAA	GGATAATTGT	GGATGACACC	TCAACCCAGT	4260
GGTCCAAAAA	CATGAACAT	TTGACCCCGA	GCACCTCAC	ACAGATAGAC	TACAATGAGA	4320
AGGAGAAAGG	GGCCATTACT	CAGTCTCCCT	TATCAGATTG	CCTTACGAGG	AGTCATAGCA	4380
TCCCTCAAGC	AAATAGATCT	CCATTACCCA	TTGCAAAGGT	ATCATCATTT	CCATCTATTA	4440
GACCTATATA	TCTGACCAGG	GTCCATTATC	AAGACAACCT	TTCTCATCTT	CCAGCAGCAT	4500
CTTATAGAAA	GAAAGATTCT	GGGGTCCAAG	AAAGCAGTCA	TTTCTTACAA	GGAGCCAAAA	4560
AAAAATAACCT	TTCTTTAGCC	ATTCTAACCT	TGGAGATGAC	TGGTGATCAA	AGAGAGGTTG	4620
GCTCCCTGGG	GACAAGTGCC	ACAAATTCAG	TCACATACAA	GAAAGTTGAG	AACACTGTTC	4680
TCCCGAAACC	AGACTTGCCC	AAAACATCTG	GCAAGATTGA	ATTGCTTCCA	AAAGTTTACA	4740
TTTATCAGAA	GGACCTATTC	CCTACGGAAA	CTAGCAATGG	GTCTCCTGGC	CATCTGGATC	4800
TCGTGGAAGG	GAGCCTTCTT	CAGGGAACAG	AGGGAGCGAT	TAAGTGGAAT	GAAGCAAACA	4860
GACCTGGAAA	AGTTCCCTTT	CTGAGAGTAG	CAACAGAAAG	CTCTGCAAAG	ACTCCCTCCA	4920
AGCTATTGGA	TCCTCTTGCT	TGGGATAACC	ACTATGGTAC	TCAGATACCA	AAAGAAGAGT	4980
GGAAATCCCA	AGAGAAGTCA	CCAGAAAAAA	CAGCTTTTAA	GAAAAAGGAT	ACCATTTTGT	5040
CCCtGAACGC	TTGTGAAAGC	AATCATGCAA	TAGCAGCAAT	AAATGAGGGA	CAAAATAAGC	5100
CCGAATAGTA	AGTCACCTGG	GCAAAGCAAG	GTAGGACTGA	AAGGCTGTGC	TCTCAAAACC	5160
CACCACTCTT	GAAACGCCAT	CAACGGGAAA	TAACCTGTAC	TACTCTTCAG	TCAGATCAAG	5220
AGGAAATTGA	CTATGATGAT	ACCATATCAG	TTGAAATGAA	GAAGGAAGAT	TTTGACATTT	5280
ATGATGAGGA	TGAAAATCAG	AGCCCCCGCA	GCTTTCAAAA	GAAAAACAGA	CACTATTTTA	5340
TTGCTGCAGT	GGAGAGGCTC	TGGGATTATG	GGATGAGTAG	CTCCCCACAT	GTTCTAAGAA	5400
ACAGGGCTCA	GAGTGGCAGT	GTCCCTCAGT	TCAGAAAGT	TGTTTTCCAG	GAATTTACTG	5460
ATGGCTCCTT	TACTCAGCCC	TTATACCGTG	GAGAACTAAA	TGAACATTTG	GGACTCCTGG	5520
GGCCATATAT	AAGAGCAGAA	GTTGAAGATA	ATATCATGGT	AACTTTCAGA	AATCAGGCCT	5580
CTCGTCCCTA	TTCTTCTAT	TCTAGCCTTA	TTTCTTATGA	GGAAGATCAG	AGGCAAGGAG	5640
CAGAACCTAG	AAAAAACTTT	GTCAAGCCTA	ATGAAACCAA	AACTTACTTT	TGGAAGATGC	5700
AACATCATAT	GGCACCCTAT	AAAGATGAGT	TTGACTGCAA	AGCCTGGGCT	TATTTCTCTG	5760
ATGTTGACCT	GGAAAAAGAT	GTGCACTCAG	GCCTGATTGG	ACCCCTTCTG	GTCTGCCACA	5820
CTAACACACT	GAACCTTGCT	CATGGGAGAC	AAGTGACAGT	ACAGGAATTT	GCTCTGTTTT	5880
TCACCATCTT	TGATGAGACC	AAAAGCTGGT	ACTTCACTGA	AAATATGGAA	AGAAACTGCA	5940
GGGCTCCCTG	CAATATCCAG	ATGGAAGATC	CCACTTTTAA	AGAGAATTAT	CGCTTCCATG	6000
CAATCAATGG	CTACATAATG	GATACACTAC	CTGGCTTAGT	AATGGCTCAG	GATCAAAGGA	6060
TTCGATGGTA	TCTGCTCAGC	ATGGGCAGCA	ATGAAAACAT	CCATTCTATT	CATTTTCAGT	6120
GACATGTGTT	CAGTGTACGA	AAAAAAGAGG	AGTATAAAAT	GGCACTGTAC	AATCTCTATC	6180
CAGGTGTTTT	TGAGACAGTG	GAAATGTTAC	CATCCAAAGC	TGGAATTTGG	CGGGTGGAA	6240
GCCTTATTGG	CGAGCATCTA	CATGCTGGGA	TGACACACAT	TTTCTGTTG	TACAGAAATA	6300
AGTGTGAGAC	TCCCCTGGGA	ATGGCTTCTG	GACACATTAG	AGATTTTCAG	ATTACAGCTT	6360
CAGGACAATA	TGGACAGTGG	GCCCCAAAGC	TGGCCAGACT	TCATTATTCC	GGATCAATCA	6420
ATGCCCTGGAG	CACCAAGGAG	CCCTTTTCTT	GGATCAAGGT	GGATCTGTTG	GCACCAATGA	6480
TTATTCACGG	CATCAAGACC	CAGGGTGCCC	GTGAGAAGTT	CTCCAGCCTC	TACATCTCTC	6540
AGTTTATCAT	CATGTATAGT	CTTGATGGGA	AGAAAGTGGA	GACTTATCGA	GGAAATTTCCA	6600
CTGGAACCTT	AATGGTCTTC	TTTGGCAATG	TGGATTCAATC	TGGGATAAAA	CACAATATTT	6660
TTAACCTTCC	AATTATTGCT	CGATACATCC	GTTTGCACCC	AACTCATTAT	AGCATTTCGA	6720
GCACTCTTCG	CATGGAGTTG	ATGGGCTGTG	ATTTAAATAG	TTGCAGCATG	CCATTGGGAA	6780
TGGAGAGTAA	AGCAATATCA	GATGCACAGA	TTAGTGCTTC	ATCCTACTTT	ACCAATATGT	6840

```

TTGCCACCTG GTCTCCTTCA AAAGCTCGAC TTCACCTCCA AGGGAGGAGT AATGCCTGGA 6900
GACCTCAGGT GAATAATCCA AAAGAGTGGC TGCAGTGGA CTTCCAGAAG ACAATGAAAG 6960
TCACAGGAGT AACTACTCAG GGAGTAAAT CTCTGCTTAC CAGCATGTAT GTGAAGGAGT 7020
TCCTCATCTC CAGCAGTCAA GATGGCCATC AGTGGACTCT CTTTTTTCAG AATGGCAAAG 7080
TAAAGGTTTT TCAGGGAAAT CAAGACTCCT TCACACCTGT GGTGAACTCT CTAGACCCAC 7140
CGTTACTGAC TCGCTACCTT CGAATTCACC CCCAGAGTTG GGTGCACCAG ATTGCCCTGA 7200
GGATGGAGGT TCTGGGCTGC GAGGCACAGG ACCTCTACTG AGGGTGGCCA CTGCAGCACC 7260
TGCCACTGCC GTCACCTCTC CCTCCTCAGC TCCAGGGCAG TGTCCTCCC TGGCTTGCTT 7320
TCTACCTTTG TGCTAAATCC TAGCAGACAC TGCCTTGAAG CCTCCTGAAT TAACTATCAT 7380
CAGTCCTGCA TTTCTTTGGT GGGGGGCCAG GAGGGTGCAT CCAATTTAAC TTAACCTTTA 7440
CCTATTTTCT GCAGCTGCTC CCAGATTACT CCTTCCTTCC AATATAACTA GGCAAAAAGA 7500
AGTGAGGAGA AACCTGCATG AAAGCATTCT TCCCTGAAAA GTTAGGCCTC TCAGAGTCAC 7560
CACTTCCTCT GTTGTAGAAA AACTATGTGA TGAACCTTTG AAAAGATAT TTATGATGTT 7620
AACATTTGAG GTTAAAGCCTC ATACGTTTAA AATAAACTC TCAGTTGTTT ATTATCCTGA 7680
TCAAGCATGG AACAAAGCAT GTTTCAGGAT CAGATCAATA CAATCTTGGA GTCAAAAGGC 7740
AAATCATTTG GACAATCTGC AAAATGGAGA GAATACAATA ACTACTACAG TAAAGTCTGT 7800
TTCTGCTTCC TTACACATAG ATATAATTAT GTTATTTAGT CATTATGAGG GGCACATTCT 7860
TATCTCCAAA ACTAGCATTC TTAACTGAG AATTATAGAT GGGGTTCAAG AATCCCTAAG 7920
TCCCTGAAA TTATATAAGG CATTCTGTAT AAATGCAAAT GTGCATTTTT CTGACGAGTG 7980
TCCATAGATA TAAAGCCATT TGGTCTTAAT TCTGACCAAT AAAAAATAA GTCAGGAGGA 8040
TGCAATTGTT GAAAGCTTTG AAATAAAATA ACAATGTCTT CTTGAAATTT GTGATGGCCA 8100
AGAAAGAAAA TGATGATGAC ATTAGCTTTC TAAAGGACAT ACATTTAATA TTTCTGTGGA 8160
AATATGAGGA AAATCCATGG TTATCTGAGA TAGGAGATAC AAACCTTGTA ATTCTAATAA 8220
TGCCTCAGT TTAATCTCTC CCTCTACTAA TTTCTGCTG AAAATAACAC AACAAAAATG 8280
TAACAGGGGA AATTATATAC CGTGAAGTAA AACTAGAGTC CTACTTACAT AGTTGAAATA 8340
TCAAGGAGGT CAGAAGAAA TTGGACTGGT GAAAACAGAA AAAACACTCC AGTCTGCCAT 8400
ATCACCACAC AATAGGATCC CCCTCTTGC CCTCCACCCC CATAAGATTG TGAAGGGTTT 8460
ACTGCTCCTT CCATCTGCCT GACCCCTTCA CTATGACTAC ACAGAATCTC CTGATAGTAA 8520
AGGGGGCTGG AGGCAAGGAT AAGTTATAGA GCAGTTGGAG GAAGCATCCA AAGATTGCAA 8580
CCCAGGGCAA ATGGAAACA GGAGATCCTA ATATGAAAGA AAAATGGATC CCAATCTGAG 8640
AAAAGGCAAA AGAATGGCTA CTTTTTCTA TGCTGGAGTA TTTTCTAATA ATCCTGCTTG 8700
ACCCTTATCT GACCTCTTTG GAAACTATAA CATAGCTGTC ACAGTATAGT CACAATCCAC 8760
AAATGATGCA GGTGCAATG GTTTATAGCC CTGTGAAGTT CTTAAAGTTT AGAGGCTAAC 8820
TTACAGAAAT GAATAAGTTG TTTTGTTTA TAGCCCGGTA GAGGAGTTAA CCCCAAAGGT 8880
GATATGGTTT TATTTCTGT TATGTTAAC TTAATAATCT TATTTGGCA TTCTTTCCC 8940
ATTGACTATA TACATCTCTA TTTCTCAAAT GTTCATGGAA CTAGCTCTTT TATTTTCTG 9000
CTGGTTTCTT CAGTAATGAG TTAAATAAAA CATTGACACA TACAAAAAAA AAAAAAAA 9060
AAAAAAA AAAA

```

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2351 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

```

Met Gln Ile Glu Leu Ser Thr Cys Phe Phe Leu Cys Leu Leu Arg Phe
1           5           10           15

```

```

Cys Phe Ser Ala Thr Arg Arg Tyr Tyr Leu Gly Ala Val Glu Leu Ser
20           25           30

```

```

Trp Asp Tyr Met Gln Ser Asp Leu Gly Glu Leu Pro Val Asp Ala Arg
35           40           45

```

```

Phe Pro Pro Arg Val Pro Lys Ser Phe Pro Phe Asn Thr Ser Val Val
50           55           60

```

```

Tyr Lys Lys Thr Leu Phe Val Glu Phe Thr Asp His Leu Phe Asn Ile

```

65	70	75	80
Ala Lys Pro Arg	Pro 85 Pro Trp Met Gly	Leu 90 Leu Gly Pro Thr	Ile Gln 95
Ala Glu Val Tyr	Asp 100 Thr Val Val Ile	Thr 105 Leu Lys Asn Met	Ala Ser 110
His Pro Val Ser	Leu 115 His Ala Val Gly	Val 120 Ser Tyr Trp Lys	Ala Ser 125
Glu Gly Ala Glu	Tyr 130 Asp Asp Gln Thr	Ser 135 Gln Arg Glu Lys	Glu Asp 140
Asp Lys Val Phe	Pro 150 Gly Gly Ser His Thr	Tyr 155 Val Trp Gln Val	Leu 160
Lys Glu Asn Gly	Pro 165 Met Ala Ser Asp	Pro 170 Leu Cys Leu Thr	Tyr Ser 175
Tyr Leu Ser His	Val 180 Asp Leu Val Lys	Asp 185 Leu Asn Ser Gly	Leu Ile 190
Gly Ala Leu Leu	Val 195 Cys Arg Glu Gly	Ser 200 Leu Ala Lys Glu	Lys Thr 205
Gln Thr Leu His	Lys 210 Phe Ile Leu Leu Phe	Ala 220 Val Phe Asp Glu Gly	
Lys Ser Trp His	Ser 225 Glu Thr Lys Asn Ser	Leu 235 Met Gln Asp Arg Asp	240
Ala Ala Ser Ala	Arg 245 Ala Trp Pro Lys	Met 250 His Thr Val Asn	Gly Tyr 255
Val Asn Arg Ser	Leu 260 Pro Gly Leu Ile	Gly 265 Cys His Arg Lys	Ser Val 270
Tyr Trp His Val	Ile 275 Gly Met Gly Thr	Thr 280 Pro Glu Val His	Ser Ile 285
Phe Leu Glu Gly	His 290 Thr Phe Leu Val Arg	Asn 300 His Arg Gln Ala	Ser 305
Leu Glu Ile Ser	Pro 310 Ile Thr Phe Leu Thr	Ala 315 Gln Thr Leu Leu	Met 320
Asp Leu Gly Gln	Phe 325 Leu Leu Phe Cys	His 330 Ile Ser Ser His	Gln His 335
Asp Gly Met Glu	Ala 340 Tyr Val Lys Val	Asp 345 Ser Cys Pro Glu	Glu Pro 350
Gln Leu Arg Met	Lys 355 Asn Asn Glu Glu	Ala 360 Glu Asp Tyr Asp	Asp Asp 365
Leu Thr Asp Ser	Glu 370 Met Asp Val Val	Arg 380 Phe Asp Asp Asp	Asn Ser 385
Pro Ser Phe Ile	Gln 390 Ile Arg Ser Val	Ala 395 Lys Lys His Pro	Lys Thr 400

Trp Val His Tyr Ile Ala Ala Glu Glu Glu Asp Trp Asp Tyr Ala Pro
 405 410 415
 Leu Val Leu Ala Pro Asp Asp Arg Ser Tyr Lys Ser Gln Tyr Leu Asn
 420 425 430
 Asn Gly Pro Gln Arg Ile Gly Arg Lys Tyr Lys Lys Val Arg Phe Met
 435 440 445
 Ala Tyr Thr Asp Glu Thr Phe Lys Thr Arg Glu Ala Ile Gln His Glu
 450 455 460
 Ser Gly Ile Leu Gly Pro Leu Leu Tyr Gly Glu Val Gly Asp Thr Leu
 465 470 475 480
 Leu Ile Ile Phe Lys Asn Gln Ala Ser Arg Pro Tyr Asn Ile Tyr Pro
 485 490 495
 His Gly Ile Thr Asp Val Arg Pro Leu Tyr Ser Arg Arg Leu Pro Lys
 500 505 510
 Gly Val Lys His Leu Lys Asp Phe Pro Ile Leu Pro Gly Glu Ile Phe
 515 520 525
 Lys Tyr Lys Trp Thr Val Thr Val Glu Asp Gly Pro Thr Lys Ser Asp
 530 535 540
 Pro Arg Cys Leu Thr Arg Tyr Tyr Ser Ser Phe Val Asn Met Glu Arg
 545 550 555 560
 Asp Leu Ala Ser Gly Leu Ile Gly Pro Leu Leu Ile Cys Tyr Lys Glu
 565 570 575
 Ser Val Asp Gln Arg Gly Asn Gln Ile Met Ser Asp Lys Arg Asn Val
 580 585 590
 Ile Leu Phe Ser Val Phe Asp Glu Asn Arg Ser Trp Tyr Leu Thr Glu
 595 600 605
 Asn Ile Gln Arg Phe Leu Pro Asn Pro Ala Gly Val Gln Leu Glu Asp
 610 615 620
 Pro Glu Phe Gln Ala Ser Asn Ile Met His Ser Ile Asn Gly Tyr Val
 625 630 635 640
 Phe Asp Ser Leu Gln Leu Ser Val Cys Leu His Glu Val Ala Tyr Trp
 645 650 655
 Tyr Ile Leu Ser Ile Gly Ala Gln Thr Asp Phe Leu Ser Val Phe Phe
 660 665 670
 Ser Gly Tyr Thr Phe Lys His Lys Met Val Tyr Glu Asp Thr Leu Thr
 675 680 685
 Leu Phe Pro Phe Ser Gly Glu Thr Val Phe Met Ser Met Glu Asn Pro
 690 695 700
 Gly Leu Trp Ile Leu Gly Cys His Asn Ser Asp Phe Arg Asn Arg Gly
 705 710 715 720

Met Thr Ala Leu Leu Lys Val Ser Ser Cys Asp Lys Asn Thr Gly Asp
 725 730 735
 Tyr Tyr Glu Asp Ser Tyr Glu Asp Ile Ser Ala Tyr Leu Leu Ser Lys
 740 745 750
 Asn Asn Ala Ile Glu Pro Arg Ser Phe Ser Gln Asn Ser Arg His Pro
 755 760 765
 Ser Thr Arg Gln Lys Gln Phe Asn Ala Thr Thr Ile Pro Glu Asn Asp
 770 775 780
 Ile Glu Lys Thr Asp Pro Trp Phe Ala His Arg Thr Pro Met Pro Lys
 785 790 795 800
 Ile Gln Asn Val Ser Ser Ser Asp Leu Leu Met Leu Leu Arg Gln Ser
 805 810 815
 Pro Thr Pro His Gly Leu Ser Leu Ser Asp Leu Gln Glu Ala Lys Tyr
 820 825 830
 Glu Thr Phe Ser Asp Asp Pro Ser Pro Gly Ala Ile Asp Ser Asn Asn
 835 840 845
 Ser Leu Ser Glu Met Thr His Phe Arg Pro Gln Leu His His Ser Gly
 850 855 860
 Asp Met Val Phe Thr Pro Glu Ser Gly Leu Gln Leu Arg Leu Asn Glu
 865 870 875 880
 Lys Leu Gly Thr Thr Ala Ala Thr Glu Leu Lys Lys Leu Asp Phe Lys
 885 890 895
 Val Ser Ser Thr Ser Asn Asn Leu Ile Ser Thr Ile Pro Ser Asp Asn
 900 905 910
 Leu Ala Ala Gly Thr Asp Asn Thr Ser Ser Leu Gly Pro Pro Ser Met
 915 920 925
 Pro Val His Tyr Asp Ser Gln Leu Asp Thr Thr Leu Phe Gly Lys Lys
 930 935 940
 Ser Ser Pro Leu Thr Glu Ser Gly Gly Pro Leu Ser Leu Ser Glu Glu
 945 950 955 960
 Asn Asn Asp Ser Lys Leu Leu Glu Ser Gly Leu Met Asn Ser Gln Glu
 965 970 975
 Ser Ser Trp Gly Lys Asn Val Ser Ser Thr Glu Ser Gly Arg Leu Phe
 980 985 990
 Lys Gly Lys Arg Ala His Gly Pro Ala Leu Leu Thr Lys Asp Asn Ala
 995 1000 1005
 Leu Phe Lys Val Ser Ile Ser Leu Leu Lys Thr Asn Lys Thr Ser Asn
 1010 1015 1020
 Asn Ser Ala Thr Asn Arg Lys Thr His Ile Asp Gly Pro Ser Leu Leu
 1025 1030 1035 1040
 Ile Glu Asn Ser Pro Ser Val Trp Gln Asn Ile Leu Glu Ser Asp Thr

1045					1050					1055					
Glu	Phe	Lys	Lys	Val	Thr	Pro	Leu	Ile	His	Asp	Arg	Met	Leu	Met	Asp
			1060					1065					1070		
Lys	Asn	Ala	Thr	Ala	Leu	Arg	Leu	Asn	His	Met	Ser	Asn	Lys	Thr	Thr
		1075					1080					1085			
Ser	Ser	Lys	Asn	Met	Glu	Met	Val	Gln	Gln	Lys	Lys	Glu	Gly	Pro	Ile
		1090				1095					1100				
Pro	Pro	Asp	Ala	Gln	Asn	Pro	Asp	Met	Ser	Phe	Phe	Lys	Met	Leu	Phe
		1105				1110					1115				1120
Leu	Pro	Glu	Ser	Ala	Arg	Trp	Ile	Gln	Arg	Thr	His	Gly	Lys	Asn	Ser
				1125					1130					1135	
Leu	Asn	Ser	Gly	Gln	Gly	Pro	Ser	Pro	Lys	Gln	Leu	Val	Ser	Leu	Gly
			1140					1145					1150		
Pro	Glu	Lys	Ser	Val	Glu	Gly	Gln	Asn	Phe	Leu	Ser	Glu	Lys	Asn	Lys
		1155					1160					1165			
Val	Val	Val	Gly	Lys	Gly	Glu	Phe	Thr	Lys	Asp	Val	Gly	Leu	Lys	Glu
		1170					1175					1180			
Met	Val	Phe	Pro	Ser	Ser	Arg	Asn	Leu	Phe	Leu	Thr	Asn	Leu	Asp	Asn
				1190						1195					1200
Leu	His	Glu	Asn	Asn	Thr	His	Asn	Gln	Glu	Lys	Lys	Ile	Gln	Glu	Glu
				1205					1210					1215	
Ile	Glu	Lys	Lys	Glu	Thr	Leu	Ile	Gln	Glu	Asn	Val	Val	Leu	Pro	Gln
			1220					1225					1230		
Ile	His	Thr	Val	Thr	Gly	Thr	Lys	Asn	Phe	Met	Lys	Asn	Leu	Phe	Leu
		1235					1240					1245			
Leu	Ser	Thr	Arg	Gln	Asn	Val	Glu	Gly	Ser	Tyr	Asp	Gly	Ala	Tyr	Ala
		1250				1255					1260				
Pro	Val	Leu	Gln	Asp	Phe	Arg	Ser	Leu	Asn	Asp	Ser	Thr	Asn	Arg	Thr
				1270						1275					1280
Lys	Lys	His	Thr	Ala	His	Phe	Ser	Lys	Lys	Gly	Glu	Glu	Glu	Asn	Leu
				1285					1290					1295	
Glu	Gly	Leu	Gly	Asn	Gln	Thr	Lys	Gln	Ile	Val	Glu	Lys	Tyr	Ala	Cys
			1300					1305					1310		
Thr	Thr	Arg	Ile	Ser	Pro	Asn	Thr	Ser	Gln	Gln	Asn	Phe	Val	Thr	Gln
		1315					1320					1325			
Arg	Ser	Lys	Arg	Ala	Leu	Lys	Gln	Phe	Arg	Leu	Pro	Leu	Glu	Glu	Thr
		1330				1335					1340				
Glu	Leu	Glu	Lys	Arg	Ile	Ile	Val	Asp	Asp	Thr	Ser	Thr	Gln	Trp	Ser
				1350						1355					1360
Lys	Asn	Met	Lys	His	Leu	Thr	Pro	Ser	Thr	Leu	Thr	Gln	Ile	Asp	Tyr
				1365					1370					1375	

Asn Glu Lys Glu Lys Gly Ala Ile Thr Gln Ser Pro Leu Ser Asp Cys
 1380 1385 1390
 Leu Thr Arg Ser His Ser Ile Pro Gln Ala Asn Arg Ser Pro Leu Pro
 1395 1400 1405
 Ile Ala Lys Val Ser Ser Phe Pro Ser Ile Arg Pro Ile Tyr Leu Thr
 1410 1415 1420
 Arg Val Leu Phe Gln Asp Asn Ser Ser His Leu Pro Ala Ala Ser Tyr
 1425 1430 1435 1440
 Arg Lys Lys Asp Ser Gly Val Gln Glu Ser Ser His Phe Leu Gln Gly
 1445 1450 1455
 Ala Lys Lys Asn Asn Leu Ser Leu Ala Ile Leu Thr Leu Glu Met Thr
 1460 1465 1470
 Gly Asp Gln Arg Glu Val Gly Ser Leu Gly Thr Ser Ala Thr Asn Ser
 1475 1480 1485
 Val Thr Tyr Lys Lys Val Glu Asn Thr Val Leu Pro Lys Pro Asp Leu
 1490 1495 1500
 Pro Lys Thr Ser Gly Lys Val Glu Leu Leu Pro Lys Val His Ile Tyr
 1505 1510 1515 1520
 Gln Lys Asp Leu Phe Pro Thr Glu Thr Ser Asn Gly Ser Pro Gly His
 1525 1530 1535
 Leu Asp Leu Val Glu Gly Ser Leu Leu Gln Gly Thr Glu Gly Ala Ile
 1540 1545 1550
 Lys Trp Asn Glu Ala Asn Arg Pro Gly Lys Val Pro Phe Leu Arg Val
 1555 1560 1565
 Ala Thr Glu Ser Ser Ala Lys Thr Pro Ser Lys Leu Leu Asp Pro Leu
 1570 1575 1580
 Ala Trp Asp Asn His Tyr Gly Thr Gln Ile Pro Lys Glu Glu Trp Lys
 1585 1590 1595 1600
 Ser Gln Glu Lys Ser Pro Glu Lys Thr Ala Phe Lys Lys Lys Asp Thr
 1605 1610 1615
 Ile Leu Ser Leu Asn Ala Cys Glu Ser Asn His Ala Ile Ala Ala Ile
 1620 1625 1630
 Asn Glu Gly Gln Asn Lys Pro Glu Ile Glu Val Thr Trp Ala Lys Gln
 1635 1640 1645
 Gly Arg Thr Glu Arg Leu Cys Ser Gln Asn Pro Pro Val Leu Lys Arg
 1650 1655 1660
 His Gln Arg Glu Ile Thr Arg Thr Thr Leu Gln Ser Asp Gln Glu Glu
 1665 1670 1675 1680
 Ile Asp Tyr Asp Asp Thr Ile Ser Val Glu Met Lys Lys Glu Asp Phe
 1685 1690 1695

Asp Ile Tyr Asp Glu Asp Glu Asn Gln Ser Pro Arg Ser Phe Gln Lys
 1700 1705 1710
 Lys Thr Arg His Tyr Phe Ile Ala Ala Val Glu Arg Leu Trp Asp Tyr
 1715 1720 1725
 Gly Met Ser Ser Ser Pro His Val Leu Arg Asn Arg Ala Gln Ser Gly
 1730 1735 1740
 Ser Val Pro Gln Phe Lys Lys Val Val Phe Gln Glu Phe Thr Asp Gly
 1745 1750 1755 1760
 Ser Phe Thr Gln Pro Leu Tyr Arg Gly Glu Leu Asn Glu His Leu Gly
 1765 1770 1775
 Leu Leu Gly Pro Tyr Ile Arg Ala Glu Val Glu Asp Asn Ile Met Val
 1780 1785 1790
 Thr Phe Arg Asn Gln Ala Ser Arg Pro Tyr Ser Phe Tyr Ser Ser Leu
 1795 1800 1805
 Ile Ser Tyr Glu Glu Asp Gln Arg Gln Gly Ala Glu Pro Arg Lys Asn
 1810 1815 1820
 Phe Val Lys Pro Asn Glu Thr Lys Thr Tyr Phe Trp Lys Val Gln His
 1825 1830 1835 1840
 His Met Ala Pro Thr Lys Asp Glu Phe Asp Cys Lys Ala Trp Ala Tyr
 1845 1850 1855
 Phe Ser Asp Val Asp Leu Glu Lys Asp Val His Ser Gly Leu Ile Gly
 1860 1865 1870
 Pro Leu Leu Val Cys His Thr Asn Thr Leu Asn Pro Ala His Gly Arg
 1875 1880 1885
 Gln Val Thr Val Gln Glu Phe Ala Leu Phe Phe Thr Ile Phe Asp Glu
 1890 1895 1900
 Thr Lys Ser Trp Tyr Phe Thr Glu Asn Met Glu Arg Asn Cys Arg Ala
 1905 1910 1915 1920
 Pro Cys Asn Ile Gln Met Glu Asp Pro Thr Phe Lys Glu Asn Tyr Arg
 1925 1930 1935
 Phe His Ala Ile Asn Gly Tyr Ile Met Asp Thr Leu Pro Gly Leu Val
 1940 1945 1950
 Met Ala Gln Asp Gln Arg Ile Arg Trp Tyr Leu Leu Ser Met Gly Ser
 1955 1960 1965
 Asn Glu Asn Ile His Ser Ile His Phe Ser Gly His Val Phe Thr Val
 1970 1975 1980
 Arg Lys Lys Glu Glu Tyr Lys Met Ala Leu Tyr Asn Leu Tyr Pro Gly
 1985 1990 1995 2000
 Val Phe Glu Thr Val Glu Met Leu Pro Ser Lys Ala Gly Ile Trp Arg
 2005 2010 2015
 Val Glu Cys Leu Ile Gly Glu His Leu His Ala Gly Met Ser Thr Leu

2020					2025					2030						
Phe	Leu	Val	Tyr	Ser	Asn	Lys	Cys	Gln	Thr	Pro	Leu	Gly	Met	Ala	Ser	
2035					2040					2045						
Gly	His	Ile	Arg	Asp	Phe	Gln	Ile	Thr	Ala	Ser	Gly	Gln	Tyr	Gly	Gln	
2050					2055					2060						
Trp	Ala	Pro	Lys	Leu	Ala	Arg	Leu	His	Tyr	Ser	Gly	Ser	Ile	Asn	Ala	
2065					2070					2075					2080	
Trp	Ser	Thr	Lys	Glu	Pro	Phe	Ser	Trp	Ile	Lys	Val	Asp	Leu	Leu	Ala	
2085					2090					2095						
Pro	Met	Ile	Ile	His	Gly	Ile	Lys	Thr	Gln	Gly	Ala	Arg	Gln	Lys	Phe	
2100					2105					2110						
Ser	Ser	Leu	Tyr	Ile	Ser	Gln	Phe	Ile	Ile	Met	Tyr	Ser	Leu	Asp	Gly	
2115					2120					2125						
Lys	Lys	Trp	Gln	Thr	Tyr	Arg	Gly	Asn	Ser	Thr	Gly	Thr	Leu	Met	Val	
2130					2135					2140						
Phe	Phe	Gly	Asn	Val	Asp	Ser	Ser	Gly	Ile	Lys	His	Asn	Ile	Phe	Asn	
2145					2150					2155					2160	
Pro	Pro	Ile	Ile	Ala	Arg	Tyr	Ile	Arg	Leu	His	Pro	Thr	His	Tyr	Ser	
2165					2170					2175						
Ile	Arg	Ser	Thr	Leu	Arg	Met	Glu	Leu	Met	Gly	Cys	Asp	Leu	Asn	Ser	
2180					2185					2190						
Cys	Ser	Met	Pro	Leu	Gly	Met	Glu	Ser	Lys	Ala	Ile	Ser	Asp	Ala	Gln	
2195					2200					2205						
Ile	Thr	Ala	Ser	Ser	Tyr	Phe	Thr	Asn	Met	Phe	Ala	Thr	Trp	Ser	Pro	
2210					2215					2220						
Ser	Lys	Ala	Arg	Leu	His	Leu	Gln	Gly	Arg	Ser	Asn	Ala	Trp	Arg	Pro	
2225					2230					2235					2240	
Gln	Val	Asn	Asn	Pro	Lys	Glu	Trp	Leu	Gln	Val	Asp	Phe	Gln	Lys	Thr	
2245					2250					2255						
Met	Lys	Val	Thr	Gly	Val	Thr	Thr	Gln	Gly	Val	Lys	Ser	Leu	Leu	Thr	
2260					2265					2270						
Ser	Met	Tyr	Val	Lys	Glu	Phe	Leu	Ile	Ser	Ser	Ser	Ser	Gln	Asp	Gly	
2275					2280					2285						
Gln	Trp	Thr	Leu	Phe	Phe	Gln	Asn	Gly	Lys	Val	Lys	Val	Phe	Gln	Gly	
2290					2295					2300						
Asn	Gln	Asp	Ser	Phe	Thr	Pro	Val	Val	Asn	Ser	Leu	Asp	Pro	Pro	Leu	
2305					2310					2315					2320	
Leu	Thr	Arg	Tyr	Leu	Arg	Ile	His	Pro	Gln	Ser	Trp	Val	His	Gln	Ile	
2325					2330					2335						
Ala	Leu	Arg	Met	Glu	Val	Leu	Gly	Cys	Glu	Ala	Gln	Asp	Leu	Tyr		
2340					2345					2350						

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4832 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

CTCGAGCTAA	AGATATTTTA	GAGAAGAATT	AACCTTTTGC	TTCTCCAGTT	GAACATTTGT	60
AGCAATAAGT	CATGCAAATA	GAGCTCTCCA	CCTGCTTCTT	TCTGTGCCTT	TTGCGATTCT	120
GCTTTAGTGC	CACCAGAAGA	TACTACCTGG	GTGCAGTGGA	ACTGTCATGG	GACTATATGC	180
AAAGTGATCT	CGGTGAGCTG	CCTGTGGACG	CAAGATTTC	TCCTAGAGTG	CCAAAATCTT	240
TTCCATTCAA	CACCTCAGTC	GTGTACAAAA	AGACTCTGTT	TGTAGAATTC	ACGGATCACC	300
TTTTCAACAT	CGCTAAGCCA	AGGCCACCCT	GGATGGGTCT	GCTAGGTCCCT	ACCATCCAGG	360
CTGAGGTTTA	TGATACAGTG	GTCATTACAC	TTAAGAACAT	GGCTTCCCAT	CCTGTGAGTC	420
TTCATGCTGT	TGGTGTATCC	TACTGGAAAG	CTTCTGAGGG	AGCTGAATAT	GATGATCAGA	480
CCAGTCAAAG	GGAGAAAGAA	GATGATAAAG	TCTTCCCTGG	TGGAAGCCAT	ACATATGTCT	540
GGCAGGTCCT	GAAAGAGAAT	GGTCCAATGG	CCTCTGACCC	ACTGTGCCTT	ACCTACTCAT	600
ATCTTTCTCA	TGTGGACCTG	GTAAGAGACT	TGAATTCAGG	CCTCATTGGA	GCCCTACTAG	660
TATGTAGAGA	AGGGAGTCTG	GCCAAGGAAA	AGACACAGAC	CTTGCCACAAA	TTTATACTAC	720
TTTTTGCTGT	ATTTGATGAA	GGGAAAAGTT	GGCACTCAGA	AACAAAGAAC	TCCTTGATGC	780
AGGATAGGGA	TGCTGCATCT	GCTCGGGCCT	GGCCTAAAAT	GCACACAGTC	AATGGTTATG	840
TAAACAGGTC	TCTGCCAGGT	CTGATTGGAT	GCCACAGGAA	ATCAGTCTAT	TGGCATGTGA	900
TTGGAATGGG	CACCACTCCT	GAAGTGCAC	CAATATTCCCT	CGAAGGTCAC	ACATTTCTTG	960
TGAGGAACCA	TCGCCAGGCG	TCCTTGGA	TCTCGCCAAT	AACTTTCCTT	ACTGCTCAAA	1020
CACCTTTGAT	GGACCTTGGA	CAGTTTCTAC	TGTTTTGTCA	TATCTCTTCC	CACCAACATG	1080
ATGGCATGGA	AGCTTATGTC	AAAGTAGACA	GCTGTCCAGA	GGAACCCCAA	CTACGAATGA	1140
AAAATAATGA	AGAAGCGGAA	GACTATGATG	ATGATCTTAC	TGATTCTGAA	ATGGATGTGG	1200
TCAGGTTTGA	TGATGACAAC	TCTCCTTCCT	TTATCCAAAT	TCGCTCAGTT	GCCAAGAAGC	1260
ATCCTAAAC	TTGGGTACAT	TACATTGCTG	CTGAAGAGGA	GGACTGGGAC	TATGCTCCCT	1320
TAGTCCTCGC	CCCCGATGAC	AGAAGTTATA	AAAGTCAATA	TTTGAACAAT	GGCCCTCAGC	1380
GGATTGGTAG	GAAGTACAAA	AAAGTCCGAT	TTATGGCATA	CACAGATGAA	ACCTTTAAGA	1440
CTCGTGAAGC	TATTCAGCAT	GAATCAGGAA	TCTTGGGACC	TTTACTTTAT	GGGGAAGTTG	1500
GAGACACACT	GTTGATTATA	TTAAGAATC	AAGCAAGCAG	ACCATATAAC	ATCTACCCTC	1560
ACGGAATCAC	TGATGTCCGT	CCTTTGTATT	CAAGGAGATT	ACCAAAAGGT	GTAACACATT	1620
TGAAGGATTT	TCCAATTCTG	CCAGGAGAAA	TATTCAAATA	TAAATGGACA	GTGACTGTAG	1680
AAGATGGGCC	AACTAAATCA	GATCCTCGGT	GCCTGACCCG	CTATTACTCT	AGTTTCGTTA	1740
ATATGGAGAG	AGATCTAGCT	TCAGGACTCA	TTGGCCCTCT	CCTCATCTGC	TACAAAGAA	1800
CTGTAGATCA	AAGAGGAAAC	CAGATAATGT	CAGACAAGAG	GAATGTCATC	CTGTTTTCTG	1860
TATTTGATGA	GAACCGAAGC	TGGTACCTCA	CAGAGAATAT	ACAACGCTTT	CTCCCCAATC	1920
CAGCTGGAGT	GCAGCTTGAG	GATCCAGAGT	TCCAAGCCTC	CAACATCATG	CACAGCATCA	1980
ATGGCTATGT	TTTTGATAGT	TTGCAGTTGT	CAGTTTGTGT	GCATGAGGTG	GCATACTGGT	2040
ACATTCTAAG	CATTGGAGCA	CAGACTGACT	TCCTTTCTGT	CTTCTTCTCT	GGATATACCT	2100
TCAAACACAA	AATGGTCTAT	GAAGACACAC	TCACCCTATT	CCCATTCTCA	GGAGAACTG	2160
TCTTCATGTC	GATGGAAAAC	CCAGGTCTAT	GGATTCTGGG	GTGCCACAAC	TCAGACTTTC	2220
GGAAACAGAG	CATGACCGCC	TTACTGAAGG	TTTCTAGTTG	TGACAAGAAC	ACTGGTGATT	2280
ATTACGAGGA	CAGTTATGAA	GATATTTTCA	CATACTTGCT	GAGTAAAAAC	AATGCCATTG	2340
AACCAAGAAG	CTTCTCCCAG	AACCCACCAG	TCTTGAAACG	CCATCAACGG	GAAATAACTC	2400
GTACTACTCT	TCAGTCAGAT	CAAGAGGAAA	TTGACTATGA	TGATACCATA	TCAGTTGAAA	2460
TGAAGAAGGA	AGATTTTGAC	ATTTATGATG	AGGATGAAAA	TCAGAGCCCC	CGCAGGTTTC	2520
AAAAGAAAAC	ACGACACTAT	TTTATTGCTG	CAGTGGAGAG	GCTCTGGGAT	TATGGGATGA	2580
GTAGCTCCCC	ACATGTTCTA	AGAAACAGGG	CTCAGAGTGG	CAGTGTCCCT	CAGTTCAAGA	2640
AAGTTGTTTT	CCAGGAATTT	ACTGATGGCT	CCTTTACTCA	GCCCTTATAC	CGTGGAGAAC	2700
TAAATGAACA	TTTGGGACTC	CTGGGGCCAT	ATATAAGAGC	AGAAGTTGAA	GATAATATCA	2760
TGGTAACTTT	CAGAAATCAG	GCCTCTCGTC	CCTATTCTTT	CTATTCTAGC	CTTATTTCTT	2820
ATGAGGAAGA	TCAGAGGCAA	GGAGCAGAAC	CTAGAAAAAA	CTTTGTCAAG	CCTAATGAAA	2880
CCAAAACCTA	CTTTTGAAAA	GTGCAACATC	ATATGGCACC	CACTAAAGAT	GAGTTTGACT	2940
GCAAAGCCTG	GGCTTATTTT	TCTGATGTTG	ACCTGGAAAA	AGATGTGCAC	TCAGGCCTGA	3000
TTGGACCCCT	TCTGGTCTGC	CACACTAACA	CACTGAACCC	TGCTCATGGG	AGACCAATGA	3060

```

CAGTACAGGA ATTTGCTCTG TTTTTCACCA TCTTTGATGA GACCAAAAGC TGGTACTTCA 3120
CTGAAAATAT GGAAAGAAAC TGCAGGGCTC CCTGCAATAT CCAGATGGAA GATCCCCTT 3180
TTAAAGAGAA TTATCGCTTC CATGCAATCA ATGGCTACAT AATGGATACA CTACCTGGCT 3240
TAGTAATGGC TCAGGATCAA AGGATTTCGAT GGTATCTGCT CAGCATGGGC AGCAATGAAA 3300
ACATCCATTC TATTCATTTT AGTGGACATG TGTTCACCTG ACGAAAAAAA GAGGAGTATA 3360
AAATGGCACT GTACAATCTC TATCCAGGTG TTTTGTAGAC AGTGGAAATG TTACCATCCA 3420
AAGCTGGAAT TTGGCGGGTG GAATGCCTTA TTGGCGAGCA TCTACATGCT GGGATGAGCA 3480
CACTTTTTCT GGTGTACAGC AATAAGTGTC AGACTCCCTT GGAATGGCT TCTGGACACA 3540
TTAGAGATTT TCAGATTACA GCTTCAGGAC AATATGGACA GTGGGCCCCA AAGCTGGCCA 3600
GACTTCATTA TTCCGGATCA ATCAATGCCT GGAGCACCAA GGAGCCCTTT TCTTGGATCA 3660
AGGTGGATCT GTTGGCACCA ATGATTATTC ACGGCATCAA GACCCAGGGT GCCCGTCAGA 3720
AGTTCTCCAG CCTCTACATC TCTCAGTTTA TCATCATGTA TAGTCTTGAT GGGAAGAAGT 3780
GGCAGACTTA TCGAGGAAAT TCCACTGGAA CCTTAATGGT CTTCTTTGGC AATGTGGATT 3840
CATCTGGGAT AAAACACAAT ATTTTAAACC CTCCAATTAT TGCTCGATAC ATCCGTTTGC 3900
ACCCAACTCA TTATAGCATT CGCAGCACTC TTCGCATGGA GTTGATGGGC TGTGATTTAA 3960
ATAGTTGCAG CATGCCATTG GGAATGGAGA GTAAAGCAAT ATCAGATGCA CAGATTACTG 4020
CTTCATCCTA CTTTACCAAT ATGTTTGCCA CCTGGTCTCC TTCAAAAGCT CGACTTCACC 4080
TCCAAGGGAG GAGTAATGCC TGGAGACCTC AGGTGAATAA TCCAAAAGAG TGGCTGCAAG 4140
TGGACTTCCA GAAGACAATG AAAGTCACAG GAGTAACTAC TCAGGGAGTA AAATCTCTGC 4200
TTACCAGCAT GTATGTGAAG GAGTTCCTCA TCTCCAGCAG TCAAGATGGC CATCAGTGGA 4260
CTCTCTTTT TCAGAATGGC AAAGTAAAGG TTTTTCAGGG AAATCAAGAC TCCTTCACAC 4320
CTGTGGTGAA CTCTCTAGAC CCACCGTTAC TGACTCGCTA CCTTCGAATT CACCCCAAGA 4380
GTTGGGTGCA CCAGATTGCC CTGAGGATGG AGGTTCTGGG CTGCGAGGCA CAGGACCTCT 4440
ACTGAGGGTG GCCACTGCAG CACCTGCCAC TGCCGTCAAC TCTCCCTCCT CAGCTCCAGG 4500
GCAGTGTCCC TCCCTGGCTT GCCTTCTACC TTTGTCTAA ATCCTAGCAG ACCTGCCTT 4560
GAAGCCTCCT GAATTAATA TCATCAGTCC TGCATTCTT TGGTGGGGGG CCAGGAGGGT 4620
GCATCCAATT TAACTTAAT CTACCTTATT TTCTGCAGCT GCTCCAGAT TACTCCTTC 4680
TTCCAATATA ACTAGGCAAA AAGAAGTGAG GAGAAACCTG CATGAAAGCA TTCTTCCTG 4740
AAAAGTTAGG CCTCTCAGAG TCACCACTTC CTCTGTTGTA GAAAACTAT GTGATGAAAC 4800
TTTGAAAAAG ATATTTATGA TGTGCGGCC GC 4832

```

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1457 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

```

Met Gln Ile Glu Leu Ser Thr Cys Phe Phe Leu Cys Leu Leu Arg Phe
1           5           10           15

Cys Phe Ser Ala Thr Arg Arg Tyr Tyr Leu Gly Ala Val Glu Leu Ser
20           25           30

Trp Asp Tyr Met Gln Ser Asp Leu Gly Glu Leu Pro Val Asp Ala Arg
35           40           45

Phe Pro Pro Arg Val Pro Lys Ser Phe Pro Phe Asn Thr Ser Val Val
50           55           60

Tyr Lys Lys Thr Leu Phe Val Glu Phe Thr Asp His Leu Phe Asn Ile
65           70           75           80

Ala Lys Pro Arg Pro Pro Trp Met Gly Leu Leu Gly Pro Thr Ile Gln
85           90           95

Ala Glu Val Tyr Asp Thr Val Val Ile Thr Leu Lys Asn Met Ala Ser
100          105          110

```

176

His Pro Val Ser Leu His Ala Val Gly Val Ser Tyr Trp Lys Ala Ser
 115 120 125
 Glu Gly Ala Glu Tyr Asp Asp Gln Thr Ser Gln Arg Glu Lys Glu Asp
 130 135 140
 Asp Lys Val Phe Pro Gly Gly Ser His Thr Tyr Val Trp Gln Val Leu
 145 150 155 160
 Lys Glu Asn Gly Pro Met Ala Ser Asp Pro Leu Cys Leu Thr Tyr Ser
 165 170 175
 Tyr Leu Ser His Val Asp Leu Val Lys Asp Leu Asn Ser Gly Leu Ile
 180 185 190
 Gly Ala Leu Leu Val Cys Arg Glu Gly Ser Leu Ala Lys Glu Lys Thr
 195 200 205
 Gln Thr Leu His Lys Phe Ile Leu Leu Phe Ala Val Phe Asp Glu Gly
 210 215 220
 Lys Ser Trp His Ser Glu Thr Lys Asn Ser Leu Met Gln Asp Arg Asp
 225 230 235 240
 Ala Ala Ser Ala Arg Ala Trp Pro Lys Met His Thr Val Asn Gly Tyr
 245 250 255
 Val Asn Arg Ser Leu Pro Gly Leu Ile Gly Cys His Arg Lys Ser Val
 260 265 270
 Tyr Trp His Val Ile Gly Met Gly Thr Thr Pro Glu Val His Ser Ile
 275 280 285
 Phe Leu Glu Gly His Thr Phe Leu Val Arg Asn His Arg Gln Ala Ser
 290 295 300
 Leu Glu Ile Ser Pro Ile Thr Phe Leu Thr Ala Gln Thr Leu Leu Met
 305 310 315 320
 Asp Leu Gly Gln Phe Leu Leu Phe Cys His Ile Ser Ser His Gln His
 325 330 335
 Asp Gly Met Glu Ala Tyr Val Lys Val Asp Ser Cys Pro Glu Glu Pro
 340 345 350
 Gln Leu Arg Met Lys Asn Asn Glu Glu Ala Glu Asp Tyr Asp Asp Asp
 355 360 365
 Leu Thr Asp Ser Glu Met Asp Val Val Arg Phe Asp Asp Asp Asn Ser
 370 375 380
 Pro Ser Phe Ile Gln Ile Arg Ser Val Ala Lys Lys His Pro Lys Thr
 385 390 395 400
 Trp Val His Tyr Ile Ala Ala Glu Glu Glu Asp Trp Asp Tyr Ala Pro
 405 410 415
 Leu Val Leu Ala Pro Asp Asp Arg Ser Tyr Lys Ser Gln Tyr Leu Asn
 420 425 430
 Asn Gly Pro Gln Arg Ile Gly Arg Lys Tyr Lys Lys Val Arg Phe Met

435					440					445					
Ala	Tyr	Thr	Asp	Glu	Thr	Phe	Lys	Thr	Arg	Glu	Ala	Ile	Gln	His	Glu
450						455					460				
Ser	Gly	Ile	Leu	Gly	Pro	Leu	Leu	Tyr	Gly	Glu	Val	Gly	Asp	Thr	Leu
465					470					475					480
Leu	Ile	Ile	Phe	Lys	Asn	Gln	Ala	Ser	Arg	Pro	Tyr	Asn	Ile	Tyr	Pro
				485					490					495	
His	Gly	Ile	Thr	Asp	Val	Arg	Pro	Leu	Tyr	Ser	Arg	Arg	Leu	Pro	Lys
			500					505					510		
Gly	Val	Lys	His	Leu	Lys	Asp	Phe	Pro	Ile	Leu	Pro	Gly	Glu	Ile	Phe
		515					520					525			
Lys	Tyr	Lys	Trp	Thr	Val	Thr	Val	Glu	Asp	Gly	Pro	Thr	Lys	Ser	Asp
	530					535					540				
Pro	Arg	Cys	Leu	Thr	Arg	Tyr	Tyr	Ser	Ser	Phe	Val	Asn	Met	Glu	Arg
545					550					555					560
Asp	Leu	Ala	Ser	Gly	Leu	Ile	Gly	Pro	Leu	Leu	Ile	Cys	Tyr	Lys	Glu
				565					570					575	
Ser	Val	Asp	Gln	Arg	Gly	Asn	Gln	Ile	Met	Ser	Asp	Lys	Arg	Asn	Val
			580					585					590		
Ile	Leu	Phe	Ser	Val	Phe	Asp	Glu	Asn	Arg	Ser	Trp	Tyr	Leu	Thr	Glu
		595					600					605			
Asn	Ile	Gln	Arg	Phe	Leu	Pro	Asn	Pro	Ala	Gly	Val	Gln	Leu	Glu	Asp
	610					615					620				
Pro	Glu	Phe	Gln	Ala	Ser	Asn	Ile	Met	His	Ser	Ile	Asn	Gly	Tyr	Val
625					630					635					640
Phe	Asp	Ser	Leu	Gln	Leu	Ser	Val	Cys	Leu	His	Glu	Val	Ala	Tyr	Trp
				645					650					655	
Tyr	Ile	Leu	Ser	Ile	Gly	Ala	Gln	Thr	Asp	Phe	Leu	Ser	Val	Phe	Phe
			660					665					670		
Ser	Gly	Tyr	Thr	Phe	Lys	His	Lys	Met	Val	Tyr	Glu	Asp	Thr	Leu	Thr
		675					680					685			
Leu	Phe	Pro	Phe	Ser	Gly	Glu	Thr	Val	Phe	Met	Ser	Met	Glu	Asn	Pro
	690					695					700				
Gly	Leu	Trp	Ile	Leu	Gly	Cys	His	Asn	Ser	Asp	Phe	Arg	Asn	Arg	Gly
705					710					715					720
Met	Thr	Ala	Leu	Leu	Lys	Val	Ser	Ser	Cys	Asp	Lys	Asn	Thr	Gly	Asp
				725					730					735	
Tyr	Tyr	Glu	Asp	Ser	Tyr	Glu	Asp	Ile	Ser	Ala	Tyr	Leu	Leu	Ser	Lys
			740					745					750		
Asn	Asn	Ala	Ile	Glu	Pro	Arg	Ser	Phe	Ser	Gln	Asn	Pro	Pro	Val	Leu
		755					760					765			

Lys Arg His Gln Arg Glu Ile Thr Arg Thr Thr Leu Gln Ser Asp Gln
 770 775 780
 Glu Glu Ile Asp Tyr Asp Asp Thr Ile Ser Val Glu Met Lys Lys Glu
 785 790 795 800
 Asp Phe Asp Ile Tyr Asp Glu Asp Glu Asn Gln Ser Pro Arg Ser Phe
 805 810 815
 Gln Lys Lys Thr Arg His Tyr Phe Ile Ala Ala Val Glu Arg Leu Trp
 820 825 830
 Asp Tyr Gly Met Ser Ser Ser Pro His Val Leu Arg Asn Arg Ala Gln
 835 840 845
 Ser Gly Ser Val Pro Gln Phe Lys Lys Val Val Phe Gln Glu Phe Thr
 850 855 860
 Asp Gly Ser Phe Thr Gln Pro Leu Tyr Arg Gly Glu Leu Asn Glu His
 865 870 875 880
 Leu Gly Leu Leu Gly Pro Tyr Ile Arg Ala Glu Val Glu Asp Asn Ile
 885 890 895
 Met Val Thr Phe Arg Asn Gln Ala Ser Arg Pro Tyr Ser Phe Tyr Ser
 900 905 910
 Ser Leu Ile Ser Tyr Glu Glu Asp Gln Arg Gln Gly Ala Glu Pro Arg
 915 920 925
 Lys Asn Phe Val Lys Pro Asn Glu Thr Lys Thr Tyr Phe Trp Lys Val
 930 935 940
 Gln His His Met Ala Pro Thr Lys Asp Glu Phe Asp Cys Lys Ala Trp
 945 950 955 960
 Ala Tyr Phe Ser Asp Val Asp Leu Glu Lys Asp Val His Ser Gly Leu
 965 970 975
 Ile Gly Pro Leu Leu Val Cys His Thr Asn Thr Leu Asn Pro Ala His
 980 985 990
 Gly Arg Gln Val Thr Val Gln Glu Phe Ala Leu Phe Phe Thr Ile Phe
 995 1000 1005
 Asp Glu Thr Lys Ser Trp Tyr Phe Thr Glu Asn Met Glu Arg Asn Cys
 1010 1015 1020
 Arg Ala Pro Cys Asn Ile Gln Met Glu Asp Pro Thr Phe Lys Glu Asn
 1025 1030 1035 1040
 Tyr Arg Phe His Ala Ile Asn Gly Tyr Ile Met Asp Thr Leu Pro Gly
 1045 1050 1055
 Leu Val Met Ala Gln Asp Gln Arg Ile Arg Trp Tyr Leu Leu Ser Met
 1060 1065 1070
 Gly Ser Asn Glu Asn Ile His Ser Ile His Phe Ser Gly His Val Phe
 1075 1080 1085

Thr Val Arg Lys Lys Glu Glu Tyr Lys Met Ala Leu Tyr Asn Leu Tyr
 1090 1095 1100
 Pro Gly Val Phe Glu Thr Val Glu Met Leu Pro Ser Lys Ala Gly Ile
 1105 1110 1115 1120
 Trp Arg Val Glu Cys Leu Ile Gly Glu His Leu His Ala Gly Met Ser
 1125 1130 1135
 Thr Leu Phe Leu Val Tyr Ser Asn Lys Cys Gln Thr Pro Leu Gly Met
 1140 1145 1150
 Ala Ser Gly His Ile Arg Asp Phe Gln Ile Thr Ala Ser Gly Gln Tyr
 1155 1160 1165
 Gly Gln Trp Ala Pro Lys Leu Ala Arg Leu His Tyr Ser Gly Ser Ile
 1170 1175 1180
 Asn Ala Trp Ser Thr Lys Glu Pro Phe Ser Trp Ile Lys Val Asp Leu
 1185 1190 1195 1200
 Leu Ala Pro Met Ile Ile His Gly Ile Lys Thr Gln Gly Ala Arg Gln
 1205 1210 1215
 Lys Phe Ser Ser Leu Tyr Ile Ser Gln Phe Ile Ile Met Tyr Ser Leu
 1220 1225 1230
 Asp Gly Lys Lys Trp Gln Thr Tyr Arg Gly Asn Ser Thr Gly Thr Leu
 1235 1240 1245
 Met Val Phe Phe Gly Asn Val Asp Ser Ser Gly Ile Lys His Asn Ile
 1250 1255 1260
 Phe Asn Pro Pro Ile Ile Ala Arg Tyr Ile Arg Leu His Pro Thr His
 1265 1270 1275 1280
 Tyr Ser Ile Arg Ser Thr Leu Arg Met Glu Leu Met Gly Cys Asp Leu
 1285 1290 1295
 Asn Ser Cys Ser Met Pro Leu Gly Met Glu Ser Lys Ala Ile Ser Asp
 1300 1305 1310
 Ala Gln Ile Thr Ala Ser Ser Tyr Phe Thr Asn Met Phe Ala Thr Trp
 1315 1320 1325
 Ser Pro Ser Lys Ala Arg Leu His Leu Gln Gly Arg Ser Asn Ala Trp
 1330 1335 1340
 Arg Pro Gln Val Asn Asn Pro Lys Glu Trp Leu Gln Val Asp Phe Gln
 1345 1350 1355 1360
 Lys Thr Met Lys Val Thr Gly Val Thr Thr Gln Gly Val Lys Ser Leu
 1365 1370 1375
 Leu Thr Ser Met Tyr Val Lys Glu Phe Leu Ile Ser Ser Ser Gln Asp
 1380 1385 1390
 Gly His Gln Trp Thr Leu Phe Phe Gln Asn Gly Lys Val Lys Val Phe
 1395 1400 1405
 Gln Gly Asn Gln Asp Ser Phe Thr Pro Val Val Asn Ser Leu Asp Pro

1410 1415 1420

Pro Leu Leu Thr Arg Tyr Leu Arg Ile His Pro Gln Ser Trp Val His
 1425 1430 1435 1440

Gln Ile Ala Leu Arg Met Glu Val Leu Gly Cys Glu Ala Gln Asp Leu
 1445 1450 1455

Tyr

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 100 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

Arg Gly Met Thr Ala Leu Leu Lys Val Ser Ser Cys Asp Lys Asn Thr
 1 5 10 15

Gly Asp Tyr Tyr Glu Asp Ser Tyr Glu Asp Ile Ser Ala Tyr Leu Leu
 20 25 30

Ser Lys Asn Asn Ala Ile Glu Pro Arg Ser Phe Ser Gln Asn Ser Arg
 35 40 45

His Pro Ser Thr Arg Gln Lys Gln Phe Asn Ala Thr Pro Pro Thr Pro
 50 55 60

Pro Thr Pro Pro Val Leu Lys Arg His Gln Arg Glu Ile Thr Arg Thr
 65 70 75 80

Thr Leu Gln Ser Asp Gln Glu Glu Ile Asp Tyr Asp Asp Thr Ile Ser
 85 90 95

Val Glu Met Lys
 100

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 300 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:
- | | | | | | | |
|------------|------------|------------|------------|------------|-------------|-----|
| AGAGGCATGA | CCGCCTTACT | GAAGGTTTCT | AGTTGTGACA | AGAACACTGG | TGATTATTAC | 60 |
| GAGGACAGTT | ATGAAGATAT | TTCAGCATAC | TTGCTGAGTA | AAAACAATGC | CATTGAACCA | 120 |
| AGAAGCTTCT | CCCAGAATTC | TAGACACCCT | AGCACTAGGC | AAAAGCAATT | TAATGCCACC | 180 |
| CCTCCTACAC | CACCAACCCC | ACCAGTACTG | AAACGCCATC | AACGGGAAAT | AACCTCGTACT | 240 |
| ACTCTTCAGT | CTGATCAAGA | GGAAATTGAC | TATGATGATA | CCATATCAGT | TGAAATGAAG | 300 |

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

Ser Arg His Pro Ser Thr Arg Gln Lys Gln Phe Asn Ala Thr Pro Pro
 1 5 10 15

Val Leu Lys Arg His Gln Arg Glu Ile Thr Arg
 20 25

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 81 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

TCGCGACACC CTAGCACTAG GCAAAAGCAA TTAAATGCCA CCCCACCAGT CCTGAAACGC 60
 CATCAACGGG AAATAACGCG T 81

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

ACTACTCTTC AATCTGATCA AGAGGAA 27

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 43 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

CGCGCCGCTC GAGTCTACAA TGGCTTTGCC TTTTGCTTTA CTG 43

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 43 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GCGCCCATCG ATTTATTCCT TCCTCCTTAA CCTTTCTGC AAG 43

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 48 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

- (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:
CGCGCCGCTC GAGCATCCCA ATGGCCCTGT CCTTTCTTT ACTGATGG 48
- (2) INFORMATION FOR SEQ ID NO:56:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:
CCATCGATT C AATCCTTCCT CCTTAATCTT TTTTGCAAG 39
- (2) INFORMATION FOR SEQ ID NO:57:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:
CCGGATCCTC TACAATGGCT TTGCCTTTTG CTTTACTG 38
- (2) INFORMATION FOR SEQ ID NO:58:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 46 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:
CGCGCCGGCG GCCGCTTATT CCTTCCTCCT TAACCTTTCT TGCAAG 46
- (2) INFORMATION FOR SEQ ID NO:59:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:
CCGGATCCCA TCCCAATGGC CCTGTCCTTT TCTTTACTGA TGG 43
- (2) INFORMATION FOR SEQ ID NO:60:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 46 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:
CGCGCCGGCG GCCGCTCAAT CCTTCCTCCT TAATCTTTT TGCAAG 46
- (2) INFORMATION FOR SEQ ID NO:61:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 90 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:
AGCTTGCTGT TTGTGTGCTG CCTCTGAAGT CCACACTGAA CAAACTTCAG CCTACTCATG 60
TCCCTAAAAT GGGCAAACAT TGCAAGCAGC 90

(2) INFORMATION FOR SEQ ID NO:62:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:
AAACAGCAAA CACACAGCCC TCCCTGCCTG CTGACCTTGG AGCTGGGGCA GAGGTCAGAG 60
ACCTCTCTGA 70

(2) INFORMATION FOR SEQ ID NO:63:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 82 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:
AGCTTCAGAG AGGTCTCTGA CCTCTGCCCC AGCTCCAAGG TCAGCAGGCA GGGAGGGCTG 60
TGTGTTTGCT GTTTGCTGCT TG 82

(2) INFORMATION FOR SEQ ID NO:64:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 78 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:
CAATGTTTGC CCATTTTAGG GACATGAGTA GGCTGAAGT TGTTCAGTGT GGACTTCAGA 60
GGCAGCACAC AAACAGCA 78

(2) INFORMATION FOR SEQ ID NO:65:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 90 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:
AATTCGCTGT TTGTGTGCTG CCTCTGAAGT CCACACTGAA CAAACTTCAG CCTACTCATG 60
TCCCTAAAAT GGGCAAACAT TGCAAGCAGC 90

(2) INFORMATION FOR SEQ ID NO:66:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:
AAACAGCAAA CACACAGCCC TCCCTGCCTG CTGACCTTGG AGCTGGGGCA GAGGTCAGAG 60
ACCTCTCTGG 70

(2) INFORMATION FOR SEQ ID NO:67:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 82 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

AATCCAGAG AGGTCTCTGA CCTCTGCCCC AGCTCCAAGG TCAGCAGGCA GGGAGGGCTG 60
TGTGTTTGCT GTTTGCTGCT TG 82

(2) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 78 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

CAATGTTTGC CCATTTTAGG GACATGAGTA GGCTGAAGTT TGTTCAGTGT GGA CTTCAGA 60
GGCAGCACAC AAACAGCG 78

(2) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

CGCGCCGCC GGGGTAGATC TTGCTACCAG TGG 33

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

GCGCCCGCGG CCGCCACTGT CCCAGGTCAG TGGTGGTGCC 40

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

CGCGCCGGCG GCCGCTCTAC AATGGCCTTG ACCTTTGCTT TACTGG 46

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

GCGCCCATCG ATTCATTCCT TACTTCTTAA ACTTTCTTGC AAG 43

- (2) INFORMATION FOR SEQ ID NO:73:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: peptide
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:
 Asn Ser Arg His Pro Ser
 1 5
- (2) INFORMATION FOR SEQ ID NO:74:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:
 Gln Asn Pro Pro Val Leu Lys Arg His Gln Arg Glu Ile Thr
 1 5 10
- (2) INFORMATION FOR SEQ ID NO:75:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:
AATTCGCGAC ACCCTAGC 18
- (2) INFORMATION FOR SEQ ID NO:76:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:
CAAAACCCAC CAGTCTTGAA ACGCCATCAA CGGGAAATAA CG 42
- (2) INFORMATION FOR SEQ ID NO:77:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:
GCGCTGTGGG ATCGGTTTTG GGTGGTCAGA AC 32

(2) INFORMATION FOR SEQ ID NO:78:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:
TTTGCGGTAG TTGCCCTTA TTGC

24

(2) INFORMATION FOR SEQ ID NO:79:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: peptide
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:
Arg Thr Leu Gln Ser Asp
1 5

(2) INFORMATION FOR SEQ ID NO:80:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:
CGTACTCTTC AGTCT

15

(2) INFORMATION FOR SEQ ID NO:81:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: DNA (genomic)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:
GCATGAGAAG TCAGACTAG

19

Claims

WHAT IS CLAIMED

1. A replication defective recombinant retrovirus expressing a B domain-deleted human factor VIII protein, wherein said recombinant retrovirus is capable of infecting human cells.
2. The recombinant retrovirus of claim 1, wherein said recombinant retrovirus preparation has a titer on HT1080 cells of greater than 10^6 cfu/ml.
3. The recombinant retrovirus of claim 3, wherein said retrovirus has a titer on HT1080 cells of greater than 10^7 cfu/ml.
4. The recombinant retrovirus of claim 1, wherein said retrovirus is resistant to degradation by human complement.
5. The recombinant retrovirus of claim 1, wherein said factor VIII protein comprises the amino acid sequence depicted in Seq. ID No. 47.
6. The recombinant retrovirus of claim 6, wherein said factor VIII protein is encoded by the nucleic acid sequence depicted in Seq. ID No. 46.
7. A recombinant retrovirus expressing a factor VIII protein, said recombinant retrovirus comprising retroviral structural proteins, a retroviral vector nucleic acid backbone, and a nucleic acid sequence encoding a factor VIII protein operably linked to a promoter, wherein said retroviral vector nucleic acid backbone is selected from the group consisting of pBA-5a, pBA-5b and pBA-5c.
8. The recombinant retrovirus of claim 7 wherein said factor VIII protein is a B domain-deleted factor VIII protein.

9. The retroviral vector of claim 8 wherein said B domain-deleted factor VIII protein is the factor VIII SQN deletion.
10. The retroviral vector of claim 9 wherein said factor VIII protein comprises the amino acid sequence depicted in Seq. ID No. 47.
11. The retroviral vector of claim 10 wherein said factor VIII protein is encoded by the nucleic acid sequence depicted in Seq. ID No. 46.
12. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a replication defective recombinant retrovirus expressing a B domain-deleted human factor VIII protein, wherein said recombinant retrovirus is capable of infecting human cells.
13. The pharmaceutical composition of claim 12, wherein said recombinant retrovirus preparation has a titer on HT1080 cells of greater than 10^6 cfu/ml.
14. The pharmaceutical composition of claim 13, wherein said retrovirus has a titer on HT1080 cells of greater than 10^7 cfu/ml.
15. The pharmaceutical composition of claim 12, wherein said retrovirus is resistant to degradation by human complement.
16. The pharmaceutical composition of claim 12, wherein said factor VIII protein comprises the amino acid sequence depicted in Seq. ID No. 47.
17. The pharmaceutical composition of claim 16, wherein said factor VIII protein is encoded by the nucleic acid sequence depicted in Seq. ID No. 46.

18. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a recombinant retrovirus expressing a factor VIII protein, said retrovirus comprising retroviral structural proteins, a retroviral vector nucleic acid backbone, and a nucleic acid sequence encoding a factor VIII protein operably linked to a promoter, wherein said retroviral vector nucleic acid backbone is selected from the group consisting of pBA-5a, pBA-5b and pBA-5c.

19. The pharmaceutical composition of claim 18 wherein said factor VIII protein is a B domain-deleted factor VIII protein.

20. The pharmaceutical composition of claim 19 wherein said B domain-deleted factor VIII protein is the factor VIII SQN deletion.

21. The pharmaceutical composition of claim 20 wherein said factor VIII protein comprises the amino acid sequence depicted in Seq. ID No. 47.

22. The pharmaceutical composition of claim 21 wherein said factor VIII protein is encoded by the nucleic acid sequence depicted in Seq. ID No. 46.

23. A method of obtaining long term systemic expression of recombinant factor VIII protein in a patient afflicted with hemophilia A comprising injecting said a preparation of recombinant retrovirus expressing factor VIII protein into a blood vessel of said patient, wherein said recombinant retrovirus preparation is resistant to human complement and expresses human factor VIII protein in the blood of said patient resulting in a measurable level of human factor VIII protein being produced in said patient for a period of at least 30 days after the injection of said recombinant retrovirus preparation.

24. The method of claim 23, wherein said recombinant retrovirus expresses a B domain-deleted form of factor VIII.

25. The method of claim 24, wherein said B domain-deleted form of factor VIII is the SQN mutation.

26. The method of claim 23, wherein said recombinant retrovirus has a titer of greater than 10^6 on HT1080 cells.

27. The method of claim 26, wherein said retrovirus has a titer of greater than 10^7 on HT1080 cells.

28. The method of claim 23, wherein said long term systemic expression results in measurable levels of human factor VIII protein being produced in the blood of said human for a period of at least six months after the administration of said recombinant retroviral vector.

29. The method of claim 23 wherein said blood vessel is a peripheral vein.

30. A replication defective recombinant retrovirus expressing human factor IX protein, wherein said recombinant retrovirus is capable of infecting human cells and is resistant to degradation by human complement.

31. The replication defective recombinant retrovirus of claim 30, wherein said recombinant retrovirus has a titer of greater than 10^6 on HT1080 cells.

32. The replication defective recombinant retrovirus of claim 31, wherein said retrovirus has a titer of greater than 10^7 on HT1080 cells.

33. A replication defective recombinant retrovirus expressing human factor IX protein, wherein said recombinant retrovirus preparation is resistant to degradation by human complement and is capable of inducing long term systemic expression of human factor IX when administered intravenously to a human afflicted with hemophilia B, wherein

said long term systemic expression results in a measurable level of recombinant human factor IX protein being produced in the blood of said human for a period of at least 30 days after the administration of said recombinant retrovirus.

34. The replication defective recombinant retrovirus of claim 33, wherein said recombinant retrovirus has a titer of greater than 10^6 on HT1080 cells.

35. The replication defective recombinant retrovirus of claim 34, wherein said retrovirus has a titer of greater than 10^7 on HT1080 cells.

36. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a replication defective recombinant retrovirus expressing human factor IX protein, wherein said recombinant retrovirus is resistant to degradation by human complement and is capable of inducing long term systemic expression of human factor IX when administered intravenously to a human afflicted with hemophilia B, wherein said long term systemic expression results in a measurable level of recombinant human factor IX protein being produced in the blood of said human for a period of at least 30 days after the administration of said recombinant retrovirus.

37. The pharmaceutical composition of claim 36 wherein said recombinant retrovirus has a titer of greater than 10^6 on HT1080 cells.

38. The pharmaceutical composition of claim 37, wherein said retrovirus has a titer of greater than 10^7 on HT1080 cells.

39. A method of administering a preparation of replication defective recombinant retrovirus expressing human factor IX protein to a patient afflicted with hemophilia B comprising injecting said preparation of recombinant retrovirus into a blood vessel of said patient, wherein said recombinant retrovirus preparation is resistant to human complement and expresses human factor IX protein into the blood of said patient resulting in a

measurable level of human factor IX protein being produced in said patient for a period of at least 30 days after the injection of said recombinant retrovirus preparation.

40. The method of claim 39 wherein said recombinant retrovirus has a titer of greater than 10^6 on HT1080 cells.

41. The method of claim 40, wherein said retrovirus has a titer of greater than 10^7 on HT1080 cells.

1/43





FIG. I




FIG. 2





FIG. 3




FIG. 4




FIG. 5

6/43





FIG. 6

7/43





FIG. 7




FIG. 8

9/43




FIG. 9

10/43




FIG. 10

11/43




FIG. II

12/43





Fig 12

13/43

**Formulation:**

25mM Tris pH 7.2
60mM NaCl
1 mg/ml Arginine
5 mg/ml HSA
50 mg/ml Lactose

- -80°C Liquid
- -20°C Liquid
- ▽— -20°C Lyoph
- Refrig. Lyoph
- △--- R.T. Lyoph

FIG. 13





FIG. 14

15/43

**Formulation:**

25mM Tris pH 7.2

60mM NaCl

1mg/ml Arginine

5mg/ml HSA

50mg/ml Trehalose

—■— -80°C Liquid

·····▲····· -20°C Liquid

-◆- -20°C Lyoph

—●— Refrig. Lyoph

-★- R.T. Lyoph

FIG 15

16/43




FIG. 16A




FIG. 16B

17/43




FIG. 16C




FIG 16D

18/43




FIG. 17

19/43




FIG. 18




FIG. 19




FIG. 20B




FIG. 20A




FIG. 20D




FIG. 20C

23/43





FIG. 21

24/43

Fig 22



25/43




Fig 23

26/43





Fig 24

27/43

Time Course of hGH Expression in Mice Injected with DA 827




Time Point 1 (Day 14) is an average of each pool.

Fig 25

28/43

Fig 26



29/43




Fig 27

30/43




Fig 28

31/43

EXPRESSION OF HGH PROTEIN IN MOUSE ORGAN LYSATES




Fig 29

32/43

P888 NORMAL DOGS




Fig 30

Transfer of hGH Expression by Spleenocytes




Fig 31

34/43

HEMOPHILIAC DOGS




Fig 32

35/43

pKT1





Fig 33

Fig 34

RETROVIRAL BACKBONE (N2-derived)



CROSS-LESS RETROVIRAL BACKBONE: pBA-5




Fig 35




Fig 36





Fig 37



2341 ArgGlyMetThrAlaLeuLeuLysValSerSerCysAspLysAsnThrGlyAspTyrTyr Seq ID No. 48
 AGAGGGATGACCGCCTTACTGAAGGTTTCTAGTTGTGACAAGAACACTGGTGATTATTAC Seq ID No. 49
 TCTCCGTACTGGCGGAATGACTTCCAAGATCAACACTGTTCTTGTGACCACTAATAATG

2401 GluAspSerTyrGluAspIleSerAlaTyrLeuLeuSerLysAsnAsnAlaIleGluPro
 GAGGACAGTTATGAAGATATTTCAAGCATCTTGCTGAGTAAAAACAATGCCATTGAACCA
 CTCCTGTCAATACTTCTATAAAGTCGTATGAACGACTCATTTTTGTACGGTAACTTGGT

<----- N-terminus of beta domain ----->

2461 ArgSerPheSerGlnAsnSerArgHisProSerThrArgGlnLysGlnPheAsnAlaThr
 AGAAGCTTCTCCAGAAATTCTAGACACCCTAGCACTAGGCAAAAGCAATTTAATGCCACC
 TCTTCGAAGAGGGTCTTAAGATCTGTGGGATCGTGATCGTTTTCGTTAAATTACGGTGG

2463 HIND3, 2475 ECOR1, 2479 XBAI,

<-- IgA hinge ---><-- C-term. beta domain -->

2521 ProProThrProProThrProProValLeuLysArgHisGlnArgGluIleThrArgThr
 CCTCCTACACCACCAACCCACCACTACTGAAACGCCATCAACGGGAAATAACTCGTACT
 GGAGGATGTGGTGGTTGGGGTGGTCATGACTTTGCGGTAGTTGCCCTTTATTGAGCATGA

2544 SCAI,

2581 ThrLeuGlnSerAspGlnGluGluIleAspTyrAspAspThrIleSerValGluMetLys
 ACTCTTCAGTCTGATCAAGAGGAAATTGACTATGATGATAACCATATCAGTTGAAATGAAG
 TGAGAAGTCAGACTAGTTCTCCTTTAACTGATACTACTATGGTATAGTCAACTTTACTTC

2592 BCL1,

Fig 38

Fig 39

ECOR1
 NRU1
 MLU1
 BCL1



Seq ID No. 73 AsnSerArgHisProSer GlnAsnProProValLeuLysArgHisGlnArgGluIleThr
 F8-14E
 Seq ID No. 75 2 AATTCGGACACCCTAGC CAAAACCCACCACTCTTGAAACGCCATCAACGGGAAATAACG
 F8-16E
 Seq ID No. 77 GCGCTGTGGGATCGGTTTTGGGTGGTCAGAAC TTTGCGGTAGTTGCCCTTTATTGC
 F8-15E
 F8-17E

Seq ID No. 79 ArgThrLeuGlnSerAsp
 F8-16E
 Seq ID No. 80 62 CGTACTCTTCAGTCT
 Seq ID No. 81 GCATGAGAAGTCAGACTAG
 F8-17E
 75 BCL1.

1 ECOR1, 5 NRU1, 59 MLU1.

42/43

Fig 40



Linkers for pSVF8-500B

end 92 19aa C terminal
to thrombin cleavage at 740

SerArgHisProSerThrArgGlnLysGlnPheAspAlaThrProProValLeuLysArg ~Seq ID No. 50
TGGCGACACCCTAGCACTAGGCAAAAGCAATTAAATGCCACCCCAACAGTCCTGAAACGC ~Seq ID No. 51
mutant ACGGCTGTGGGATCGTGATCCGTTTTCGTAAATTACGGTGGGCTGCTCATGACTTTGCC
wild type (TT) (CT)
NRU1

Start 80K
HisGlnArgGluIleThrArg
CATCAACGGCAAAATAACGGCT
GTAGTTGCCCTTTATTGGCA

MLU1
9aa N terminal to 80K

Fig 41




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/12, 15/57, 15/86, 7/01, 9/64, C07K 14/755, A61K 38/17, 38/36, 38/37, 48/00		A3	(11) International Publication Number: WO 98/00542
(21) International Application Number: PCT/US97/11785		(43) International Publication Date: 8 January 1998 (08.01.98)	
(22) International Filing Date: 2 July 1997 (02.07.97)		(72) Inventors: JOLLY, Douglas, J.; 277 Hillcrest Drive, Leucadia, CA 92024 (US). BARBER, Jack, R.; 1168 Carlotta Street, San Diego, CA 92129 (US). CHANG, Stephen, M., W.; 12912 Camino del Valle, Poway, CA 92064 (US). RESPESS, James, G.; 4966 Lamont Street, San Diego, CA 92109 (US). ALLEN, John, R.; 8998 Adobe Buffs Drive, San Diego, CA 92129 (US). BODER, Mordechai; 3480 Juniper Street, San Diego, CA 92104 (US). CHONG, Kimberly; 12014 Dappie Way, San Diego, CA 92128 (US). DE LA VEGA, Dan, Jr.; 4220 Ampudia Street, San Diego, CA 92103 (US). DePOLO, Nicholas, J.; 964 Santa Estella, Solano Beach, CA 92075 (US). HSU, David, Chi-Tang; 7387 Mannix Court, San Diego, CA 92122 (US). IBANEZ, Carlos, E.; 13592 Millpond Way, San Diego, CA 92129 (US). MITTELSTAEDT, Denice, M.; 13687 Quinton Road, San Diego, CA 92129 (US). PRUSSAK, Charles, E.; 5121 Camino Playa Malaya, San Diego, CA 92124 (US). GREENGARD, Judith; 9053 Mesa Woods Avenue, San Diego, CA 92126 (US).	
(30) Priority Data: 08/645,601 3 July 1996 (03.07.96) US 08/696,381 13 August 1996 (13.08.96) US Not furnished 4 June 1997 (04.06.97) US		(74) Agents: KRUSE, Norman, J. et al.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US).	
(71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
		(88) Date of publication of the international search report: 19 March 1998 (19.03.98)	

(54) Title: METHODS FOR ADMINISTRATION OF RECOMBINANT GENE DELIVERY VEHICLES FOR TREATMENT OF HEMOPHILIA

(57) Abstract

Methods are provided for obtaining measurable levels of a protein, nucleic acid molecule, or enzymatic product in a bodily fluid or cells of a human, comprising the step of administering to a human a recombinant retroviral preparation having a titer on HT1080 cells of greater than 10^5 cfu/ml, wherein the recombinant retroviral preparation is capable of directing the expression of a protein, nucleic acid molecule, or enzyme which generates an enzymatic product, such that measurable levels of the protein, nucleic acid molecule, or enzymatic product may be obtained in the bodily fluid or cells of the human.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/US 97/11785

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C12N15/57 C12N15/86 C12N7/01 C12N9/64
 C07K14/755 A61K38/17 A61K38/36 A61K38/37 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 04934 A (GENETIC THERAPY INC ;CANCER RES INST ROYAL (GB)) 22 February 1996	30-32
Y	see page 9, line 10	2-6, 13-17
X	--- C.M. LYNCH ET AL.: "Sequences in the coding region of clotting factor VIII act as dominant inhibitors of RNA accumulation and protein production" HUMAN GENE THERAPY, vol. 4, no. 3, June 1993, MARY ANN LIEBERT, INC. PUBLISHERS, NEW YORK, US, pages 259-272, XP002051708 cited in the application	1,12
Y	see the whole document	2-6, 13-17
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

12 January 1998

Date of mailing of the international search report

03.02.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Hornig, H

INTERNATIONAL SEARCH REPORT

Intern 1st Application No

PCT/US 97/11785

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 91 09122 A (KABIVITRUM AB) 27 June 1991 cited in the application see the whole document ---	2-6, 13-17
A	COSSET F -L ET AL: "HIGH-TITER PACKAGING CELLS PRODUCING RECOMBINANT RETROVIRUSES RESISTANT TO HUMAN SERUM" JOURNAL OF VIROLOGY, vol. 69, no. 12, 1 December 1995, pages 7430-7436, XP000569527 see the whole document ---	1-41
A	WO 95 03834 A (US HEALTH) 9 February 1995 see the whole document ---	1-26
A	WO 93 25698 A (US GOV AS REPRESENTED BY THE) 23 December 1993 cited in the application see the whole document ---	1-41
A	WO 91 02805 A (VIAGENE INC) 7 March 1991 cited in the application see the whole document ---	1-41
P,X	WO 96 21014 A (VIAGENE INC) 11 July 1996 see page 3, line 8 - line 14; examples 2A,7 see page 3, line 17 - page 4, line 29; claims 2-19 see page 15, line 6 - page 36, line 19 ---	1-29
P,X	WO 96 33282 A (VIAGENE INC) 24 October 1996 see page 57, line 23 - page 58, line 26 ---	1,12
P,X	GREENGARD J S ET AL: "Sustained high level factor VIII antigen expression in immune-competent rabbits following peripheral intravenous injection of retroviral vector." THIRTY-EIGHTH ANNUAL MEETING OF THE AMERICAN SOCIETY OF HEMATOLOGY, ORLANDO, FLORIDA, USA, DECEMBER 6-10, 1996. BLOOD 88 (10 SUPPL. 1 PART 1-2). 1996. 273A. ISSN: 0006-4971, XP002051681 Abstract no. 1080 see abstract ---	1-29
P,A	WO 96 21035 A (VIAGENE INC) 11 July 1996 see page 20, line 32 - page 21, line 5 see page 29, line 18 - line 25 -----	1-29

INTERNATIONAL SEARCH REPORT

In international application No.
PCT/US 97/11785

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Remark : Although claims 23-29, 39-41 are directed to a method of treatment of the human/animal body , the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/11785

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9604934 A	22-02-96	CA 2196208 A EP 0769968 A	22-02-96 02-05-97
WO 9109122 A	27-06-91	SE 465222 B AU 645539 B AU 7039391 A CA 2071875 A EP 0506757 A EP 0786474 A HU 9500489 A JP 2644084 B JP 5502161 T SE 8904239 A	12-08-91 20-01-94 18-07-91 16-06-91 07-10-92 30-07-97 28-11-95 25-08-97 22-04-93 16-06-91
WO 9503834 A	09-02-95	AU 7516594 A	28-02-95
WO 9325698 A	23-12-93	CA 2137361 A EP 0644946 A JP 9507741 T	23-12-93 29-03-95 12-08-97
WO 9102805 A	07-03-91	AU 5915394 A AU 648261 B AU 6185390 A AU 7058496 A CA 2066053 A EP 0487587 A JP 4507196 T US 5662896 A US 5591624 A US 5691177 A	16-06-94 21-04-94 03-04-91 16-01-97 19-02-91 03-06-92 17-12-92 02-09-97 07-01-97 25-11-97
WO 9621014 A	11-07-96	AU 4608096 A EP 0796331 A	24-07-96 24-09-97
WO 9633282 A	24-10-96	AU 5568296 A	07-11-96
WO 9621035 A	11-07-96	US 5681746 A AU 4602596 A EP 0795021 A	28-10-97 24-07-96 17-09-97